

AMERICAN MUSEUM NOVITATES

Number 942

Published by
THE AMERICAN MUSEUM OF NATURAL HISTORY
New York City

Sept. 10, 1937

AN EARLY PLEISTOCENE FAUNA FROM NEBRASKA

BY ERWIN H. BARBOUR¹ AND C. BERTRAND SCHULTZ²

A lower Pleistocene deposit³ containing fossil vertebrates was discovered in 1936, four miles east and one mile north of Broadwater, in NE¹/₂ of sec. 20 and NW¹/₄ of sec. 21, T. 19 N., R. 47 W., Morrill County, Nebraska. This new locality is on land owned by Mr. Dan J. Boman and Mrs. Mary A. Boman of Broadwater, and was reported to the writers by Messrs. S. R. Sweet and T. C. Middleswart of Bridgeport. Four quarries were opened during 1936.

The fossilized remains were found in a white, sandy, argillaceous, diatomaceous earth, and also in lenses of fine, bog-iron stained sand. The bones are light chocolate-brown in color and are often crushed, warped and bent. Sands and gravels occur both below and above the fossil-bearing horizon. These deposits belong to the "upper terrace" of the north side of the North Platte Valley. The fossil-bearing stratum extends some forty miles to the east and several miles to the west.

The character of the deposits at the various quarry sites, together with the fauna, suggests the former presence of swamps. Imprints of fossil reed-like plants occur frequently in the diatomaceous layer and remains of a number of aquatic animals have been found in the quarries. Among the latter are otter, beaver-like rodents, muskrat, and also fishes and frogs.

Twenty-eight mammals are represented. The horse is most abundant. The absence of mammoth, bison and musk-oxen is noted, but the writers believe these forms to be later migrants into that section during the Pleistocene.

The geology, and also the fauna, indicate that the quarries are of lower Pleistocene age, possibly Aftonian. Fossils have already been attributed to the "Aftonian" from Nebraska and near-by localities, but it is very doubtful if many of these specimens are actually of Aftonian age. See chart (Fig. 1) for suggested stratigraphic relationship of the

¹ Professor of Geology and Paleontology, University of Nebraska and Director, Nebraska State Museum.

² Assistant Curator in Paleontology and Senior Field Man, Nebraska State Museum.

³ Barbour, Erwin H., and Schultz, C. Bertrand, 1936, p. 450.

fauna of the Broadwater Quarries and other known faunas from the Nebraska Region.

The American Museum of Natural History and the Frick collections from Hay Springs have aided much in a better understanding of the middle Pleistocene fauna of Nebraska. Until recently very little has been known about the lower Pleistocene. In 1934, the Frick expedition, under the leadership of Morris F. Skinner, opened a new fossil quarry¹ of




Fig. 1. Chart showing approximate stratigraphic position of fauna from Broadwater Quarries in relation to other known Pleistocene faunas from Nebraska.

lower Pleistocene age in Brown County, Nebraska. In this peat and diatomaceous earth bed were found the remains of a number of stegomastodonts (*Stegomastodon primitivus* Osborn) and a few bones of other Pleistocene mammals. The fauna of the Broadwater Quarries adds to the list of known lower Pleistocene forms and it is hoped that future work in this locality will contribute much more information.

¹ Osborn, Henry Fairfield, 1936, pp. 726-728.

The field work at Broadwater was carried on under the direction of the junior writer for the Nebraska State Museum. Members of the parties who collected at the quarries include Messrs. S. R. Sweet, T. C. Middleswart, W. F. Chaloupka, and John Ochoa of Bridgeport, and Messrs. Thompson M. Stout, E. L. Blue, Frank Crabill, Gordon Graham, David Abbott, Jack Graham, Dean Kerl, and Mrs. Marian Schultz of the Nebraska State Museum expedition, and Mr. Fred Schall of Princeton University.

The writers are deeply grateful to Mr. Childs Frick and Dr. Walter Granger of The American Museum of Natural History and Dr. C. Lewis Gazin of the National Museum for helpful suggestions and allowing the examination of specimens. Mr. Thompson M. Stout has liberally cooperated with the writers in the study of the rodents from the quarries. The drawings for figures 2 to 4 were made by Mr. Ralph Mefferd.

LIST OF MAMMALS FROM BROADWATER AND
HAY SPRINGS QUARRIES

Broadwater¹
(lower Pleistocene)

Hay Springs²
(middle Pleistocene)

INSECTIVORA

Sorex sp.

EDENTATA

Mylodon sp.

Mylodon garmani Allen

Mylodon nebrascensis (Brown)

Megalonyx leidy Lindahl

LAGOMORPHA

Lepus sp.

Sylvilagus sp.

RODENTIA

Sciurid

Geomys sp.

Thomomys?

Procastoroides sweeti, new genus and
species

Cynomys niobrarius Hay

Geomys sp.

Thomomys sp.

Castoroides ohioensis nebrascensis
Barbour

Castor sp.

¹ A preliminary list.

² Matthew, W. D., 1918, pp. 226-229.

Frick, Childs, 1930, pp. 70-80; 1937, pp. 199-202, 521-537

Schultz, C. Bertrand, 1934, pp. 357-393.

Broadwater
(lower Pleistocene)

Dipoides?

Peromyscus sp. [near *P. maniculatus*
(Wagner)]

Ondatra sp.

Mimomys?

Microtinid indet.

Zapus sp. [near *Z. hudsonius*
(Zimmerman)]

Neotoma?

CARNIVORA

Canis sp. (near *C. latrans* Say)

Canis sp. [near *Canis* (*Aenocyon*)
dirus Leidy]

Satherium piscinaria middleswarti,
new variety

Felis?

PROBOSCIDEA

Stegomastodon mirificus primitivus
Osborn

PERISSODACTYLA

Equus sp. (near *E. excelsus* Leidy)

Equus sp. (more primitive species)
Equid (very light-limbed form;
known only from a single meta-
podial)

ARTIODACTYLA

Camelops sp. (much larger form than
C. kansanus Leidy)

Tanupolama sp. [much larger form
than *Tanupolama americanus*
(Wortman)]

Camelid, possibly *Titanotylopus*
nebrascensis Barbour and Schultz

Hay Springs
(middle Pleistocene)

Ondatra nebrascensis (Hollister)

Microtus?

Canis latrans? Say

Canis (*Aenocyon*) *dirus nebrascensis*
Frick

Arctodus simus nebrascensis Frick

Mustela vision? Schreber

Smilodon nebrascensis Matthew

Archidiskodon imperator (Leidy)

Equus excelsus Leidy

Equus excelsus niobrarensis Hay

Equus calabatus nebrascensis Frick

Platygonus vetus Leidy

Camelops kansanus Leidy

Camelops vitakerianus? (Cope)

Tanupolama americanus (Wortman)





Fig. 2. *Procastoroides sweeti*. Type. N.S.M. 100-12-6-36S.P., skull, top and right side; N.S.M. 164-12-6-36S.P., ramus, right side. One-half natural size.

Broadwater (lower Pleistocene)	Hay Springs (middle Pleistocene)
<i>Capromeryx</i> sp. (larger form than <i>C. furcifer</i> Matthew)	<i>Odocoileus sheridanus</i> Frick <i>Capromeryx furcifer</i> Matthew
Antilocaprid, probably <i>Tetrameryx</i>	<i>Tetrameryx</i> (<i>Hayoceros</i>) <i>falkenbachii</i> Frick Bovid

The collection from the Broadwater Quarries also contains the following unidentified vertebrates:

PISCES

Various forms

AMPHIBIA

A very large collection of frog material

REPTILIA

Turtle, perhaps two forms

Snake, at least one form

AVES

Various forms

DESCRIPTION OF NEW MATERIAL

CASTOROIDIDAE

Procastoroides sweeti, new genus and species

TYPE.—Nebr. State Mus. Nos. 100-12-6-36S.P., skull, and 164-12-6-36S.P., left ramus.

REFERRED.—4 skull fragments, 7 rami, 33 molars and premolars, 20 incisors and incisor fragments, 2 scapulae, 4 humeri, 5 ulnae, 2 partial radii, 5 portions of pelvi, 2 femora, 1 tibia, 3 astragali, 4 misc. foot bones, 1 distal phalanx, and misc. ribs and vertebrae from the Broadwater Quarries.

LOCALITY.—Broadwater Quarry 3, NW¹/₄ of sec. 21, T. 19 N., R. 47 W., Morrill County, Nebraska.

HORIZON.—Lower Pleistocene, (?) Aftonian.

GENERIC AND SPECIFIC CHARACTERS.—Form and proportions very beaver-like, approaching those of *Castoroides*; approximately three-fourths of the size of *Castoroides*; incisors without longitudinal grooves but cheek-tooth pattern as in *Castoroides*; skull brachycephalic and more flattened than that of *Castoroides*; width of rear of skull less than one-half its length while in *Castoroides*, rear of skull much wider proportionately; narrowest part of brain-case at the middle of length of skull and just behind orbits as in *Castoroides*; limb elements like *Castoroides*.

DISCUSSION.—The size variation of the referred specimens is great. Upper incisors vary from 8 mm. wide in immature specimens to as much as 15 mm. in old individuals. Perhaps when more material is secured it may be determined that the largest forms represent another variety or species.



N.S.M. 100-12-6-36S.P



N.S.M. 164-12-6-36S.P

Fig. 3. *Procacstoroides sweeti*. Type. N.S.M. 100-12-6-36S.P., upper right dentition, reversed; N.S.M. 164-12-6-36S.P., lower left dentition. Natural size.

Thanks are due to Professor William Berryman Scott for suggestions concerning the description of this material.

Measurements of type skull and ramus of *Procacstoroides sweeti*, N.S.M. 100- and 164-12-6-36S.P.

	mm.
Length (max., including supraoccipital crest and incisors).....	200.
Basal length.....	167.
Condyllo-basal length.....	177.
Width (max., across zygomatic arches).....	145.
Length of dental series (P ⁴ -M ³) on triturating.....	43.5
Length of P ⁴	12.5
Width of P ⁴	10.5
Length of M ¹	10.
Width of M ¹	10.
Length of M ²	10.
Width of M ²	9.
Length of M ³	11.5
Width of M ³	8.
Max. width of incisor.....	11.5
Length (max., including incisors, measured parallel to grinding surface of premolar-molar series).....	170.
Length from posterior of condyle to tip of incisor.....	156.
Depth of jaw below anterior of P ₄	46.
Length of dental series (P ₄ -M ₃) on triturating surface.....	49.
Length of P ₄	15.
Width of P ₄	10.5
Length of M ₁	13.
Width of M ₁	10.
Length of M ₂	12.
Width of M ₂	10.5

Length of M_3	10.
Width of M_3	9.
Max. width of incisor.....	11.5

MUSTELIDAE

Satherium piscinaria middleswarti, new variety

TYPE.—Nebr. State Mus. Nos. 250-12-6-36S.P., incomplete skull, and 251-12-6-36S.P., right ramus.

LOCALITY.—Broadwater Quarry 3, NW $\frac{1}{4}$ of sec. 21, T. 19 N., R. 47 W., Morrill County, Nebraska.




Fig. 4. *Satherium piscinaria middleswarti*. Type. N.S.M. 250-12-6-36S.P., crown and side views of right upper dentition; N.S.M. 251-12-6-36S.P., right ramus and crown view of right lower dentition. Natural size.

HORIZON.—Lower Pleistocene, (?) Aftonian.

DIAGNOSIS.—Very close to *Satherium piscinaria* from Hagerman, Idaho, but slightly larger; lower teeth similar to those of the Hagerman species but carnassial heavier, especially posteriorly; upper carnassial from Hagerman with a much more definite anterior cusp.

Measurements of type skull and ramus of *Satherium piscinaria middleswarti*, N.S.M. 250- and 251-12-6-36S.P.

	mm.
Width of skull between orbits.....	28.
Length of dental series (C-M ¹).....	43.5
Length of premolar series.....	32.
Length of P ³	9.3
Width of P ³	6.1
Length of P ⁴	15.5
Width of P ⁴	11.5
Length of M ¹	9.5
Transverse diameter of M ¹	15.
Width across upper incisors.....	16.
Length of ramus from posterior of condyle to anterior of P ₃	72.5
Depth of jaw below anterior of M ₁	18.5
Length of P ₃ -M ₁	35.
Length of P ₃	8.5
Width of P ₃	5.2
Length of P ₄	10.
Width of P ₄	6.2
Length of M ₁	17.
Width of M ₁	9.3

REFERENCES

- BARBOUR, ERWIN HINCKLEY. 1931. 'The giant beaver, *Castoroides*, and the common beaver, *Castor*, in Nebraska.' Bull. Nebr. State Mus., I, No. 20, June, pp. 171-186.
- BARBOUR, ERWIN HINCKLEY, AND SCHULTZ, C. BERTRAND. 1936. 'Palaeontologic and geologic consideration of early man in Nebraska (with notice of a new bone bed in the early Pleistocene of Morrill County, Nebraska).' Bull. Nebr. State Mus., I, No. 45, April, p. 450.
- CALVIN, SAMUEL. 1909. 'Aftonian mammalian fauna.' Bull. Geol. Soc. Am., XX, pp. 341-356, Pls. xvi-xxviii.
1911. 'Aftonian mammalian fauna, II.' Bull. Geol. Soc. Am., XXII, pp. 207-216, Pls. xviii-xxiv.
- FOSTER, J. W. 1838. 'Organic remains.' Ohio Geol. Surv., 2nd Ann. Report, pp. 79-83, figs. A-D.
- FRICK, CHILDS. 1930. 'Alaska's frozen fauna.' Natural History, XXX, No. 1, Jan.-Feb., pp. 70-80 (list of Hay Springs fauna included).
1937. 'Horned ruminants of North America.' Bull. Am. Mus., LXIX, pp. 196-202, 521-537, figs. 20, 54.
- GAZIN, C. LEWIS. 1934. 'Upper Pliocene mustelids from the Snake River Basin of Idaho.' Jour. Mam., XV, No. 2, May, pp. 137-149, figs. 3-5.
1936. 'A study of the fossil horse remains from the upper Pliocene of Idaho.' Proc. U. S. Nat. Mus., LXXXIII, No. 2985, pp. 281-320, Pls. xxiii-xxxiii.

- HAY, OLIVER P. 1912. 'The Pleistocene mammals of Iowa.' *Iowa Geol. Surv.*, XXIII, pp. 1-662, figs. 1-142, Pls. I-LXXV.
- KAY, GEORGE FREDERICK. 1931. 'Classification and duration of the Pleistocene period.' *Bull. Geol. Soc. Am.*, XLII, No. 1, pp. 425-466, 9 figs.
- LUGN, A. L. 1934. 'Outline of Pleistocene geology of Nebraska.' *Bull. Nebr. State Mus.*, I, No. 41, Pt. I, Oct., pp. 319-356.
1935. 'The Pleistocene geology of Nebraska.' *Bull. Nebr. Geol. Surv.* (2nd Ser.), No. 10, pp. 1-223.
- MATTHEW, W. D. 1918. 'Contributions to the Snake Creek fauna (with notes upon the Pleistocene of western Nebraska.)' *Bull. Am. Mus.*, XXXVIII, Art. 7, Apr. 18, pp. 226-229.
- OSBORN, HENRY FAIRFIELD. 1936. 'Proboscidea.' *Monograph Am. Mus.*, pp. 726-728.
- SCHULTZ, C. BERTRAND. 1934. 'The Pleistocene mammals of Nebraska.' *Bull. Nebr. State Mus.*, I, No. 41, Pt. II, Oct., pp. 357-393, table A.
- SHIMEK, B. 1908. 'Aftonian sands and gravels in western Iowa.' *Science* (n.s.), XXVIII, p. 923.
1909. 'Aftonian sands and gravels in Iowa.' *Bull. Geol. Soc. Am.* XX, pp. 399-408, Pls. XXXIII-XXXVII.
- WILSON, ROBERT W. 1933. 'A rodent fauna from later Cenozoic beds of southwestern Idaho.' *Carn. Inst. Wash. Pub.*, No. 440, pp. 117-135, figs. 1-8, Pls. I-II.

