AMERICAN MUSEUM NOVITATES

Number 3934, 22 pp.

August 9, 2019

New fossil birds from the earliest Eocene of Mongolia

SARAH C. HOOD,¹ CHRIS R. TORRES,^{1, 2} MARK A. NORELL,^{3, 4} AND JULIA A. CLARKE^{1, 2, 4}

ABSTRACT

Understanding of the Asian early Paleogene avifauna is limited relative to that of North American and European avifauna of the same period. While major patterns of mammalian faunal exchange among these three regions across the Paleocene/Eocene boundary have been described, much less is known about the dynamics of bird diversity over the same time interval. Here, we report bird fossils from the earliest Eocene Bumban Member of the Naranbulag Formation in central Mongolia that add to the known record from Asia from just after this boundary. Most of this material, collected by the joint American Museum of Natural History/ Mongolia Academy of Sciences expeditions, is referable to a previously described taxon in Presbyornithidae (Anseriformes). However, five isolated elements are identified as comprising at least four species from at least three other major avian clades. While further inclusive phylogenetic analyses of each of these clades are necessary, the new remains represent possible earliest occurrences in Asia of these clades. The material includes a humerus and a furcula from shorebirds (Pan-Charadriiformes), a quadrate from a stem member of the flamingogrebe lineage (Pan-Mirandornithes), and a coracoid from a stem galliform (Pangalliformes). We also report a humerus with uncertain phylogenetic affinities but with similarities to core Gruiformes. These new fossils expand our knowledge of the Asian avifauna during this time and have the potential to further inform our understanding of the early biogeography of these clades. The shorebird and flamingo-grebe material indicate that both these lineages were present in Asia by the earliest Eocene. The pan-mirandornithine quadrate provides insight into the early feeding ecology of the flamingo-grebe clade.

¹ Jackson School of Geosciences, University of Texas at Austin, Austin.

² Department of Integrative Biology, University of Texas at Austin, Austin.

³ Richard Gilder Graduate School, American Museum of Natural History, New York, NY.

⁴ Division of Paleontology, American Museum of Natural History, New York, NY.

Copyright © American Museum of Natural History 2019

INTRODUCTION

The mammalian fossil record of Asia during the earliest Eocene is relatively well characterized and marks a major period of faunal turnover, including the first appearances on the continent of primates, perissodactyls, and artiodactyls (Dashzeveg, 1988; Meng and McKenna, 1998; Bowen et al., 2002; Kondrashov et al., 2004; Smith et al., 2006; Bajpai et al., 2008). By contrast, the Asian avifauna from this interval remains poorly understood, and bird remains from key mammal-bearing deposits are poorly sampled (Mayr, 2009; Hwang et al., 2010). In Asia, only nine bird taxa are currently described from the Paleocene and at least 33 from the Eocene (table 1). Many of these taxa are described from fragmentary single elements.

The Eocene Naranbulag Formation of Mongolia holds potential to inform our understanding of avian diversity at this time. Naranbulag Formation fossils have greatly contributed to our understanding of early Paleogene mammal diversity in Asia (Dashzeveg et al., 1998, and citations therein). The lower bound of the Bumban Member of this formation is placed at or just after the Paleocene-Eocene boundary based on biostratigraphic correlation of the Bumban fauna with other faunal assemblages (Dashzeveg, 1988; Meng and McKenna, 1998, Bowen et al., 2002), and was constrained to between 55.7–54.97 Ma by Bowen et al. (2002) using isotope stratigraphy, magnetostratigraphy, and quantitative biochronology. Several fossiliferous mammalian sites are located in the Bumban Member, including the Tsagaan Khushuu "Quarry 1" locality described by Dashzeveg et al. (1998: 194) as containing, "one of the most diverse early Eocene faunas known from Asia." Avian remains from the Naranbulag Formation are comparatively rare. The Bumban Member has produced abundant isolated elements of the extinct anseriform *Presbyornis* (Kurochkin and Dyke, 2010), as well as isolated records of proposed stem owls (Strigiformes [Kurochkin and Dyke, 2011]) and the earliest Asian record of a stemmember landfowl (Pangalliformes [Hwang et al., 2010]).

Here, we report new avian remains recovered from the Bumban Member by joint American Museum of Natural History/Mongolia Academy of Sciences expeditions between 2000 and 2002. All material was recovered from in and around "Quarry 2" of Dashzeveg et al. (1998) at Tsagaan Khushuu in the Nemegt Basin, Mongolia (fig. 1). This site lies at the base of the Bumban Member, suggesting an age of ~55 Ma. Out of the 86 collected avian elements, ~20% (19 specimens) are readily identifiable as presbyornithid remains, and much of the rest is fragmentary and nondiagnostic. Five of the remaining specimens can be unambiguously distinguished from presbyornithids. Of these, three represent proposed earliest records of their respective lineages in Asia.

MATERIALS AND METHODS

Quadrate terminology follows Elzanowski and Stidham (2010). All other osteological terminology follows Baumel and Witmer (1993). All terms have been anglicized. All measurements are in mm. IGM 100/1268 and IGM 100/1418 were scanned using high-resolution 3D X-ray computed tomography (CT) at the University of Texas High-Resolution X-ray Computed Tomography Facility, Austin, Texas (UTCT) on 22–23 May 2018, on a NSI scanner with a Fein Focus

lable I. A summary of the Pale	eogene fossil bird record of Asia. Also see	earlier reviews in Ness	ov (1992), Mayr (2	.009) and Wang et al. (2012a).
Species	Clade	Age	Locality	Reference
Qinornis paleocenica*	Ornithurae	Early Paleocene	China	Xue, 1995
Protoplotus beauforti	Protoplotidae, Suliformes	Paleocene	Sumatra	Lambrecht, 1931; van Tets et al., 1989; Stidham et al., 2005
Wanshuina lii	?Rallidae, 'Gruiformes'	Paleocene	China	Hou, 1994
Qianshanornis rapax	Qianshanornithidae	Middle Paleocene	China	Mayr et al., 2013a
Naranbulagornis khun	Anseriformes	Late Paleocene	Mongolia	Zelenkov, 2019
Odontopteryx "Pseudontornis" tsulensis	Pelagornithidae, Odontopterygiformes	Late Paleocene	Kazakhstan	Averianov et al., 1991
Eopuffinus kazakhstanensis	Procellariiformes	Late Paleocene	Kazakhstan	Nessov, 1986
Tshulia litorea**	Prophaethontidae, Phaethontiformes	Late Paleocene	Kazakhstan	Nessov, 1988; Mayr and Scofield, 2016
Zhylgaia aestiflua	Prophaethontidae, Phaethontiformes	Late Paleocene	Kazakhstan	Nessov, 1988; Bourdon et al., 2008; Mayr and Scofield 2015
?Berruornis sp.	Strigiformes	Late Paleocene	Kazakhstan	Nessov, 1992
Unnamed	Galliformes	Early Eocene	Mongolia	Hwang et al., 2010
"Zhongyuanus" Gastornis xichuanensis	Gastornithidae, Anseriformes	Early Eocene	China	Hou, 1980; Buffetaut, 2013
Songzia heidangkouensis	'Gruiformes'	Early Eocene	China	Hou, 1990; Wang et al., 2012a
Unnamed	Indet.	Early Eocene	Mongolia	
1 or 2 Unnamed	Pan-Charadriiformes	Early Eocene	Mongolia	
Unnamed	Pan-Mirandornithes	Early Eocene	Mongolia	
Unnamed	?Quercymegapodiidae, Pangalliformes	Early Eocene	Mongolia	
Presybornis mongoliensis	Presbyornithidae, Anseriformes	Early Eocene	Mongolia	Kurochkin and Dyke, 2010
Presbyornis sp. indet.	Presbyornithidae, Anseriformes	Early Eocene	Mongolia	Kurochkin and Dyke, 2010
Eostrix tsaganica	Protostrigidae, Strigiformes	Early Eocene	Mongolia	Kurochkin and Dyke, 2011
Romainvillia kazakhstanensis	Romainvilliidae, Anseriformes	Early Eocene	Kazakhstan	Zelenkov, 2018
Unnamed	cathartid-like	Early Eocene	India	Mayr et al., 2010

Table 1. Continued				
Species	Clade	Age	Locality	Reference
Unnamed	Indet.	Early Eocene	India	Mayr et al., 2013b
Vastanavis cambayensis	Vastanavidae, ?Psittaciformes	Early Eocene	India	Mayr et al., 2010
Vastanavis eocaena	Vastanavidae, ?Psittaciformes	Early Eocene	India	Mayr et al., 2007; Mayr et al., 2010
Vastanavis sp.	Vastanavidae, ?Psittaciformes	Early Eocene	India	Mayr et al., 2010
Jiliniornis huadianensis	?Charadriidae, Charadriiformes	Middle Eocene	China	Hou and Ericson, 2002
Eociconia sangequanensis	Ciconiidae, Ciconiiformes	Middle Eocene	China	Hou, 1989
Sanshuiornis zhangi	?Ciconiiformes	Middle Eocene	China	Wang et al., 2012b
Unnamed	galliform-like	Middle Eocene	Ukraine	Mayr and Zvonok, 2011
Colymbiculus udovinchenkoi	Gaviidae, Gaviiformes	Middle Eocene	Ukraine	Mayr and Zvonok, 2011
Unnamed	gruiform-like	Middle Eocene	Ukraine	Mayr and Zvonok, 2011
Dasornis sp.	Pelagornithidae, Odontopterygiformes	Middle Eocene	Ukraine	Mayr and Zvonok, 2011
Lutetodontopteryx tethyensis	Pelagornithidae, Odontopterygiformes	Middle Eocene	Ukraine	Mayr and Zvonok, 2012
Kievornis rogovitshi	?Procellariformes	Middle Eocene	Ukraine	Averianov et al. 1990; Zvonok et al., 2015
Mangystania humilicristata	?Suliformes	Middle Eocene	Kazakhstan	Zvonok et al., 2016
Murunkus subitus	?Procellariformes	Middle Eocene	Uzbekistan	Panteleyev and Nessov, 1993
Unnamed	?Threskiornithidae, Ciconiiformes	Middle Eocene	Myanmar	Stidham et al., 2005
Xorazmortyx turkestanensis	Paraortygidae, Galliformes	Middle Eocene	Uzbekistan	Zelenkov and Panteleyev, in press
Zheroia kurochkini	'Gruiformes'	Middle Eocene	Uzbekistan	Nessov, 1988
Eogrus aeola	Eogruidae, Gruiformes	Mid-Late Eocene	Kazakhstan, Mongolia	Wetmore, 1934
2 Unnamed	Anatidae, ?Romainvilliinae, Anseriformes	Late Eocene	China	Stidham and Ni, 2014
Eogrus crudus	Eogruidae, Gruiformes	Late Eocene	Mongolia	Kurochkin, 1981
Eogrus turanicus	Eogruidae, Gruiformes	Late Eocene	Kazakhstan	Kurochkin, 1981; Bendukidze 1971
Telecrex grangeri	?Phasianidae, Galliformes	Late Eocene	Mongolia	Wetmore, 1934; Olson, 1974
Minggangia changgouensis	Threskiornithidae, Ciconiiformes	Late Eocene	China	Hou, 1982
Cygnavus formosus	Anatidae, Anseriformes	Early Oligocene	Kazakhstan	Kurochkin, 1968

Table 1. Continued				
Species	Clade	Age	Locality	Reference
Ergilornis rapidus	Eogruidae, Gruiformes	Early Oligocene	Mongolia	Kozlova, 1960
Ergilornis minor	Eogruidae, Gruiformes	Early Oligocene	Mongolia	Kozlova, 1960
Sonogrus gregalis	Eogruidae, Gruiformes	Early Oligocene	Mongolia	Kurochkin, 1981
Heterostrix tatsinensis	Heterostrigidae, Strigiformes	Early Oligocene	Mongolia	Kurochkin and Dyke, 2011
Buteo circoides	Accipitridae, Accipitriformes	Oligocene	Mongolia	Kurochkin, 1968
Venerator dementjevi	Accipitridae, Accipitriformes	Oligocene	Mongolia	Kurochkin, 1968
Gobihierax edax	Accipitridae, Accipitriformes	Oligocene	Mongolia	Kurochkin, 1968
?Somateria sp.	Anatidae, Anseriformes	Late Oligocene	Kazakhstan	Kurochkin, 1968
Colymboides sp.	Gaviiformes	Late Oligocene	Kazakhstan	Kurochkin, 1976
Palaeorallus alienus	Galliformes	Oligocene	Mongolia	Kurochkin, 1968; Cracraft, 1973
Limicorallus saiensis	Phalacrocoracidae, Suliformes	Late Oligocene	Kazakhstan	Kurochkin, 1968
Cygnopterus lambrechti	Phoenicopteriformes	Late Oligocene	Kazakhstan	Kurochkin, 1968; Zelenkov, 2013
Agnopterus turgaiensis	Phoenicopteridae, Phoenicopteriformes	Late Oligocene	Kazakhstan	Tugarinov, 1940; Mlikovsky and Svec, 1986; Zelenkov, 2013
Unnamed	Plotopteridae, Suliformes	Oligocene	Japan	Hasegawa et al., 1979
Copepteryx hexeris	Plotopteridae, Suliformes	Late Oligocene	Japan	Olson and Hasegawa, 1996
Copepteryx titan	Plotopteridae, Suliformes	Late Oligocene	Japan	Olson and Hasegawa, 1996
Hokkaidornis abashiriensis	Plotopteridae, Suliformes	Late Oligocene	Japan	Sakurai et al., 2008
Megagallinula harundinea	?Rallidae, Gruiformes	Late Oligocene	Kazakhstan	Kurochkin, 1968
*Oinornis holocconico is libely a n	M_{max} neornithing (Y_{max} 1005)			

**Qinornis paleocenica* is likely a non-neornithine (Xue, 1995). ** *Tshulia litorea* is proposed to be a synonym of *Zhylgaia aestiflua* by Mayr and Scofield (2015).



FIGURE 1. Map showing the location of Tsagaan Khushuu locality in the central Gobi Desert of Mongolia, where the fossils were recovered (modified from Clarke and Norell, 2004).

Microfocal X-ray source. IGM 100/1268 was scanned at 130 kV/0.14 mA with voxel size = 15.6 μ m, resulting in 1981 total slices. IGM 100/1418 was scanned at 130 kV/0.14 mA with voxel size = 9.7 μ m, resulting in 983 total slices. Data was visualized and segmented in Avizo 9.3 (FEI).

INSTITUTIONAL ABBREVIATIONS: IGM, Mongolian Institute of Geology, Ulaan Bataar, Mongolia; GPIM, Institute of Geosciences of Johannes-Gutenberg-University Mainz, Germany; TMM, Texas Memorial Museum, Austin, Texas; USTL, Laboratoire de Paléontologie de l'Université des Sciences et Techniques du Languedoc, Montpellier, France.

SYSTEMATIC PALEONTOLOGY

AVES Linnaeus, 1758

PAN-CHARADRIIFORMES, new clade

DEFINITION: "Pan-Charadriiformes" refers to the clade containing all birds more closely related to *Charadrius hiaticula* and *Sterna hirundo* than to flamingos and grebes (Mirandornithes) based on the Prum et al. (2015) reference phylogeny.

Gen. et sp. indet.

Figure 2

REFERRED SPECIMEN: IGM 100/1435 (distal end of left humerus; fig. 2).

MEASUREMENTS: Dorsoventral width of distal end, 4.7 mm.

REMARKS: IGM 100/1435 is broken immediately proximal to the dorsal supracondylar process. This humerus is from a bird slightly smaller than the smallest extant sandpiper, *Calidris minutilla* (Least Sandpiper). The fossil exhibits a well-developed dorsal supracondylar process, which projects from the shaft well proximal to the dorsal condyle. The tip is rounded and is directed proximally, and it has a slight sigmoidal curve along the base of its dorsal margin. The fossa for the m. brachialis is preserved as a shallow scar with a well-demarcated ventral



FIGURE 2. Photographs and line drawings of the left distal humerus of a fossil shorebird specimen, IGM 100/1435, in **A** and **B**, cranial; and **D** and **E**, caudal views, compared to the left distal humerus of a Western Sandpiper *Calidris mauri* (TMM M-2742) in **C**, cranial; and **F**, caudal views. Abbreviations: **dc**, dorsal condyle; **dsp**, dorsal supracondylar process; **fp**, flexor process; **ii**, intercondylar incisure; **fmb**, fossa for the m. brachia-lis; **of**, olecranon fossa; **sh**, sulcus for the m. humerotricipitalis; **ss**, sulcus for the m. scapulotricipitalis; **vc**, ventral condyle; **vst**, ventral supracondylar tubercle. Scale bar = 2 mm.

terminus and fainter margins proximally and dorsally (fig. 2A, B). The dorsal and ventral condyles are of similar maximum length (fig. 2A, B, D, E), and the flexor process is short and broad. The olecranon fossa is deep and undercuts the edge of the flexor process, giving that process a dorsally hooked margin in caudal view. A ridge separates the sulcus for the m. scapulotricipitalis from the sulcus for the m. humerotricipitalis. The sulcus for the m. scapulotricipitalis is narrower than the sulcus for the m. humerotricipitalis (fig. 2D, E).

COMPARISONS: A well-developed dorsal supracondylar process as in IGM 100/1435 is observed in shorebirds, tubenoses (Procellariiformes), swifts and hummingbirds (Apodiformes), and songbirds (Passeriformes) among extant birds. However, the new fossil humerus is otherwise dissimilar in size and proportion to those found in swifts, hummingbirds, and songbirds; specifically, the dorsal supracondylar process is more strongly projected as well as being angled more cranially than dorsally. In the new fossil, this process is narrow and projects proximally, consistent with shorebirds (fig. 2C); by contrast, in tubenoses this process is broad and projects more cranially (Hou and Ericson, 2002). Additionally, this fossil shows a distinct sulcus for the m. scapulotricipitalis, which is observed in shorebirds but not in tubenoses (Zvonok et al., 2015). The new fossil can be further differentiated from most tubenoses by the presence of a shallow fossa for the m. brachialis; among nonalbatross tubenoses, this fossa is deep. IGM 100/1435 also exhibits a ventral condyle that projects distally past the dorsal condyle, consistent with shorebirds but not tubenoses, in which these two condyles show similar distal projections (Zvonok et al., 2015). Features of the distal humerus alone are not diagnostic for most lineages within shorebirds (Zvonok et al., 2015; Mayr, 2016), so further taxonomic assignment for the new specimen is not possible. Because this fossil cannot be assigned to crown-group charadriiforms, we place this new fossil within Pan-Charadriiformes.

A slightly younger ($\sim 2-3$ Ma) distal humerus (SMF Av 619) was reported from the early Eocene Nanjemoy Formation of Virginia (Mayr, 2016). This specimen is similar in size to IGM 100/1435 but has a shallower fossa for the m. brachialis and the dorsal supracondylar process has a slightly narrower base.

Gen. et sp. indet.

Figure 3

REFERRED SPECIMEN: IGM 100/1268 (furcula missing both omal extremities; fig. 3).

MEASUREMENTS: Maximum preserved length of right ramus, 13.7 mm; maximum length of furcular apophysis, 3 mm; maximum preserved interclavicular width from lateral margins of rami, 7.5 mm.

REMARKS: The apophysis of IGM 100/1268 is slightly crushed but retains its shape in lateral view. Mechanical preparation of the fragile rami and apophysis is impossible, and so these parts of the specimen remain encased in rock. Digitally "prepped" renderings derived from high-resolution CT data were used for comparison with other birds. The fossil is from an individual slightly smaller than the extant Least Sandpiper. The furcula is a narrow U-shape in cranial view. Toward the apophysis, the rami mediolaterally broaden and dorsoventrally flatten (fig. 3A, C). The furcular apophysis is bladelike and sharply projects dorsally and perpendicularly to the rami (fig. 3A). The ventral margin is curved in lateral view (fig. 3A).

COMPARISONS: IGM 100/1268 has a well-developed, bladelike furcular apophysis, similar to those of charadriiforms, passeriforms, and galliforms (e.g., Fürbringer, 1888; Nesbitt et al., 2009; Smith, 2014). Among these groups, the new fossil is most similar to small extant charadriiforms (e.g., *Calidris, Charadrius, Turnix*), which also exhibit furculae with dorsally directed apophyses as well as dorsoventrally compressed and gently ventrally curved rami (Smith, 2014; fig. 3B, D). By contrast, the apophysis in galliforms is elongate and projects sternodorsally, well past the sternal margin of the rami, resulting in more V-shaped furculae (Fürbringer, 1888; Nesbitt et al., 2009). Galliforms also do not exhibit the curvature of the rami observed in IGM 100/1268 in lateral view. Passeriforms have furculae with rami that are variably arched or curved in lateral view and are highly dorsoventrally compressed, with no appreciable broadening toward the sternal extremity, unlike in IGM 100/1268. Based on these comparisons we assign IGM 100/1268 to Pan-Charadriiformes.



FIGURE 3. Digital renderings from CT data of the furcula of a second fossil shorebird specimen, IGM 100/1268, in **A**, right lateral, and **C**, dorsal views, compared to the furcula of a Killdeer *Charadrius vociferous* (TMM M-3245) in **B**, right lateral, and **D**, dorsal views. Abbreviation: **a**, furcular apophysis. Scale bars = 2 mm.

PAN-MIRANDORNITHES, new clade

DEFINITION: "Pan-Mirandornithes" refers to the clade including all birds more closely related to *Phoenicopterus ruber* and *Podiceps cristatus* than to shorebirds (Charadriiformes) based on the reference phylogeny in Prum et al. (2015).

REFERRED TAXA: In addition to crown-group Mirandornithes (flamingos and grebes), Pan-Mirandornithes also includes the proposed stem mirandornithine *Juncitarsus* (Olson and Feduccia, 1980; Mayr, 2014).

Gen. et sp. indet.

Figure 4

REFERRED SPECIMEN: IGM 100/1418 (left quadrate lacking most of the orbital process).

MEASUREMENTS: Lateromedial width of mandibular articulation, 6 mm; maximum dorsoventral height, 9.4 mm.

REMARKS: IGM 100/1418 shows slight abrasion of the otic process and body. It is similar in size to the quadrate of the extant grebe *Podiceps grisegena* (Red-Necked Grebe). We used digital renderings derived from high-resolution CT scan data to aid comparison with other taxa.

The otic and squamosal facets are both slightly abraded, and the intercapitular incisure is wide and shallow (fig. 4A, B, K, L). A distinct tympanic crest runs ventromedially from the lateral margin of the squamosal facet to the medial face of the body where it meets the medial crest (fig. 4A, B, F,



FIGURE 4. Comparison of the left quadrates of Pan-Mirandornithes, including digital rendering from CT data and line drawings of the proposed Mongolian pan-mirandornithine, IGM 100/1418, in **A**, **B**, lateral; **F**, **G**, medial; **K**, **L**, caudal; and **P**, **Q**, ventral views; Chilean Flamingo *Phoenocopterus chilensis* (TMM M-5325) in **C**, lateral; **H**, medial; **M**, caudal; and **R**, ventral views; Eared Grebe *Podiceps nigricollis* (TMM M-1394) in **D**, lateral; **I**, medial: **N**, caudal; and **S**, ventral views; and line drawings of *Palaelodus ambiguus* (GPIM Op 395) based on photographs in Mayr (2015) in **E**, lateral; **J**, medial; **O**, caudal, and **T**, ventral views. Abbreviations: **bf**, basiorbital fossa; **bpf**, basiorbital pneumatic foramina; **c**, caudal condyle; **cfm**, convex articular facet of medial condyle; **i**, intercapitular incisure; **l**, lateral condyle; **m**, medial condyle; **mc**, medial crest; **mp**, medial process; **o**, otic facet; **or**, orbital process; **ot**, otic process; **paf**, pterygoid articular facet; **pc**, pterygoid condyle; **qc**, quadratojugal cotyle; **rpf**, rostromedial pneumatic foramina; **s**, squamosal facet; **tc**, tympanic crest; **vqf**, ventral quadratojugal facet. Scale bars = 2 mm.

G, L, K). The medial crest is also well defined and runs from the medial margin of the otic facet down to the medial condyle (fig. 4F, G). Between these crests lies a deep furrow, containing four small dorsoventrally aligned pneumatic foramina (fig. 3B). The basiorbital fossa, bounded by the medial crest and the base of the orbital process, contains three additional basiorbital pneumatic foramina also arranged in a line (fig. 4F, G). No caudomedial pneumatic foramina are present.

The pterygoid articulation is conformed as a facet on the medial base of the orbital process (fig. 4F, G). The specimen lacks a pterygoid condyle on the medial process, but it is unclear whether it is missing due to breakage (fig. 4F, G, K, L). The mandibular articulation consists of distinct lateral, caudal, and medial condyles. The medial condyle is convex mediorostrally with a small lip and is developed laterally as a concave articular facet (fig. 4K, L, P, Q). The lateral and caudal condyles are nearly confluent, separated by a shallow depression (fig. 4K, L, P, Q). The lateral process is moderately projected and bears a deeply concave quadratojugal cotyle with a complete, rounded margin (fig. 4A, B). The ventral margin of this cotyle is markedly expanded into a rostroventrally oriented lip in lateral view (fig. 4A, B, K, L).

COMPARISONS: We assign IMG 100/1418 to Pan-Mirandornithes based on the following suite of characters: a wide and shallow intercapitular incisure; a gracile quadrate body; a deep furrow bounded by the tympanic and medial crests; a concave articular facet lateral of the medial condyle; a rostrally projecting lip formed by the medial condyle; and a deeply concave quadratojugal socket. The combined presence of rostromedial and basiorbital pneumatic foramina is observed in both extant and stem flamingos (e.g., *Palaelodus*). These foramina are absent in most grebes (fig. 4I); however, a loss of pneumatic foramina is commonly observed in diving birds (Witmer, 1990; Mayr, 2015). This same combination of foramina placement has otherwise been reported only in some gulls (Laridae; Samejima and Otsuka, 1987; Mayr, 2015). However, in gulls, these foramina are relatively much larger than those observed in the new fossil and in stem and crown flamingos.

IMG 100/1418 exhibits additional similarities with the much younger (by ~20 Ma) stem flamingo Palaelodus ambiguus (Milne-Edwards, 1863; Cheneval and Escuillie, 1992), known from the Oligo-Miocene of Europe, which are not observed in extant flamingos and grebes. These characters include a distinct furrow bounded by the medial crest and the base of the orbital process (fig. 4J), as well as a pterygoid articular facet located on the medial base of the orbital process (Mayr, 2015; fig. 4J). It differs from *Palaelodus* by exhibiting a concave caudal margin in lateral and medial views, whereas in Palaelodus this margin is straight (fig. 4E, J). Palaelodus also has a more ventrally projecting lip on the quadratojugal facet (fig. 4E) and a more rostrally projecting lip on the medial condyle (fig. 4J) than in the new fossil. The new fossil can be differentiated from Presbyornis spp., the most abundant avian taxon from this locality, by its trituberculate mandibular articulation (presence of a caudal condyle). In contrast, Presbyornithidae bears only two mandibular condyles (lateral and medial), consistent with other galloanserines (e.g., Cracraft and Clarke, 2001; Mayr and Clarke, 2003). The isolated nature of the new fossil and the lack of relevant comparative material available for other proposed pan-mirandornithines outside the crown group (e.g., Juncitarsus; Olson and Feduccia, 1980; Peters, 1987; Mayr, 2014) prevent a confident taxonomic assignment within Mirandornithes.



FIGURE 5. Comparison of the omal ends of the coracoids of Pangalliformes, including the new fossil, IGM 100/1371, in **A**, ventral; and **B**, dorsal views; line drawings of **C**, *Quercymegapodius depereti* (USTL SNB 700) (Quercymegapodiidae); and **D**, *Paraortyx lorteti* (USTL MGB 1547) (Paraortygidae), in dorsal view, based on photographs in Mourer-Chauviré (1992). Abbreviations: **ap**, acrocoracoid process; **caf**, clavicular articular face; **g**, glenoid facet; **p**, procoracoid; **sc**, scapular cotyla. Scale bars = 2 mm.

PANGALLIFORMES Clarke, 2002

cf. QUERCYMEGAPODIIDAE Mourer-Chauviré, 1992

Gen. et sp. indet.

Figure 5

REFERRED SPECIMEN: IGM 100/1371 (omal end of left coracoid).

MEASUREMENTS: Maximum length as preserved, 13.6 mm; maximum mediolateral width of omal end, 4.7 mm.

REMARKS: The dimensions of IGM 100/1371indicate that it is from a bird significantly smaller than the smallest extant galliform, *Excalfactoria chinensis* (King Quail). The shaft is narrow with negligible sternal expansion. It bears a deeply concave scapular cotyle with a rounded, slightly teardrop shape (fig. 4B). There is no foramen associated with the passage of the n. supracoracoideus. The acrocoracoid process is hooked with its lip forming a near-per-pendicular (~100°) angle to the shaft (fig. A, B). The clavicular articular face is flat. The well-projected glenoid facet terminates at the omal margin of the scapular cotyla (fig. 5B). The procoracoid process is reduced, appearing only as a slight ridge along the medial margin mediosternal to the scapular cotyla (fig. 5B).

COMPARISONS: The morphology of IGM 100/1371 is consistent with Galliformes in its narrow shaft, hooked acrocoracoid, greatly reduced procoracoid process, and lack of a supracondylar nerve foramen. A concave scapular cotyle like that of the new fossil is not observed among crown-group galliforms but is present in stem members of the clade and considered the ancestral condition in birds (Mayr, 2000; Clarke, 2002, Mayr and Weidig, 2004). Based on these traits we assign IGM 100/1371 to the clade Pangalliformes.

Among stem galliforms, the coracoid is most like the middle-late Eocene Quercymegapodiidae with respect to the subcircular scapular cotyle, parallel medial and lateral margins of the shaft, and a near-perpendicular angle formed by the hooked acrocoracoid and the shaft. By contrast, other stem galliforms (e.g., the early Eocene Gallinuloididae and the middle-late Eocene Paraortygidae) exhibit more mediolaterally broad scapular cotylae, sternally diverging lateral and medial margins of the midshaft, and more obtuse acrocoracoid-to-shaft angles (Mourer-Chauviré, 1992; Mayr, 2000, 2006; Mayr, and Weidig, 2004; fig. 5D). The new specimen can be differentiated from previously reported Quercymegapodiidae by its subplanar clavicular articular face, which in other quercymegapodiids is slightly concave (e.g., *Quercymegapodius*; fig. 5C) or convex (e.g., *Ameripodius*) (Mourer-Chauviré, 1992, Mourer-Chauviré, 2000). The acrocoracoid process is also more hooked in IGM 1371 than in these taxa or in *Taubacrex* (Alvarenga, 1988; Mourer-Chauviré, 1992, Mourer-Chauviré, 2000).

Aves, gen. et sp. indet.

Figure 6

REFERRED SPECIMEN: IGM 100/1360 (distal end of left humerus).

MEASUREMENTS: Dorsoventral width of distal end, 5 mm.

REMARKS: IGM 100/1360 is broken proximal to the dorsal supracondylar tubercle. It is from a bird slightly larger than *Porzana carolina* (Sora Rail). The dorsal supracondylar tubercle is a proximodistally elongate rectangular projection with a slight depression on its dorsal margin (fig. 6A, B). The fossa for the m. brachialis is elongate and ventrally positioned, with clearly demarcated ventral and distal edges (fig. 6A, B). Both the ventral and dorsal condyles are weakly cranially projected. The ventral condyle is roughly half the length of the dorsal condyle and extends distal to the dorsal condyle (fig. 6A, B). The flexor process is similar in distal extent to the dorsal condyle (fig. 6A, B, D, E). Both the m. scapulotricipitalis sulcus and the olecranon fossa are broad and shallow (fig. 6D, E).

COMPARISONS: IGM 100/1360 exhibits a combination of conditions consistent with a position within the clade Messelornithidae + Ralloideae (rails and allies) (Mayr, 2004), including: craniocaudal compression, a proximodistally elongate dorsal supracondylar tubercle bearing a concavity on its margin, a narrow and ventrally situated fossa for the m. brachialis, and a well-developed flexor process (fig. 6C, F). These characters can be observed in several fossil and extant taxa of this group; the Paleocene "messel rail" *Walbeckornis creber* (Messelornithidae), the early Oligocene *Bellgirallus* (Rallidae), the early



FIGURE 6. Photographs and interpretive line drawings of the Mongolian left distal humerus, IGM 100/1360, in **A**, **B**, cranial; and **D**, **E**, caudal views, and the left distal humerus of Sora Rail *Porzana carolina* (TMM M-10382) (Rallidae) in **C**, cranial; and **F**, caudal views. Abbreviations: **dc**, dorsal condyle; **dst**, dorsal supracondylar tubercle; **fp**, flexor process; **ii**, intercondylar incisure; **fmb**, fossa for the m. brachialis; **of**, olecranon fossa; **sh**, sulcus for the m. humerotricipitalis; **ss**, sulcus for the m. scapulotricipitalis; **vc**, ventral condyle; **vst**, ventral supracondylar tubercle. Scale bars = 2 mm.

Miocene *Palaeoaramides* (Rallidae), and in the extant *Aramides saracura* (Rallidae) and *Podica senegalensis* (Heliornithidae) (Mayr, 2007; Mayr and Smith, 2002; Louchart et al., 2005). However, many of these characters are not apomorphic to the clade and are also seen outside this group. Therefore, while this new fossil may show affinities to a clade formed by messelornithids, rails, and finfoots (Mayr, 2004), we do not feel it can be assigned confidently at this time. For example, a weakly projected but elongate dorsal supracondylar tubercle with a concavity along the margin is found in the taxa listed above but is not found in all rails; in the Miocene *Baselrallus intermedius* and *Porzana carolina* the tubercle is entirely convex, and this feature shows variability in form throughout the clade (De Pietri and Mayr, 2014). An elongate, concave dorsal supracondylar tubercle is also present in the Sunbittern *Eurypyga helias* (Eurypygidae), and even in charadriiforms like *Burhinus* and members of the Alcidae (Mayr, 2007; Smith, 2014). Traits such as the development of the flexor process and the situation of the fossa for the m. brachialis are also variable.

15

DISCUSSION

The fossils we report here add between three and five new taxa to the known avifauna of the earliest Eocene Naranbulag Formation of Mongolia. The distal humerus and partial furcula from shorebirds are the oldest definitive records of Pan-Charadriiformes. These specimens were not found in association and may represent different species; however, they likely belonged to similarly sized individuals, and both bones most resemble those of small plovers (Charadriidae) and sandpipers (Scolopacidae). The previous earliest-known and confidently assigned member of the charadriiform lineage is from a slightly younger deposit of the Nanjemoy Formation in North America (53.6-52.8 Ma; Mayr, 2016). Like IGM 100/1360, this North American record is composed only of a distal humerus displaying characters similar to small extant plovers (Mayr, 2016). The previous oldest Asian charadriiform, Jiliniornis huadianensis, is from the middle Eocene of the Ukraine, and was tentatively placed in Charadriidae by Hou and Ericson (2002). Remains referred to "Graculavidae" were reported from the late Cretaceousearly Paleocene Navesink and Hornerstown Formations of New Jersey, by Olson and Parris (1987) as a potential stem lineage of shorebirds. However, these remains lack apomorphic support for placement within Pan-Charadriiformes and may not represent members of crowngroup birds (Norell and Clarke, 2001; Clarke and Norell, 2002; Mayr, 2009; Smith, 2015). Several fossil birds from the earliest Eocene of Europe have been described as "charadriiformlike" (e.g., Vanolimicola longihallucis from Germany: Mayr, 2017; Scandiavis mikkelseni from Denmark: Bertelli et al., 2013), but have not been unambiguously referred to Pan-Charadriiformes. None of these European specimens preserve the pectoral girdle, so they cannot be directly compared to the humerus or furcula reported here. Without new material and phylogenetic analyses, whether the Bumban remains are crown- or stem-clade Charadriiformes cannot be assessed, hence our referral to Pan-Charadriiformes. No preserved morphologies exclude these remains from part of the crown clade.

The new quadrate would comprise the earliest-known worldwide record of the flamingo/ grebe lineage, displacing Juncitarsus from slightly younger deposits in North America and Europe (Olson and Feduccia, 1980; Peters, 1987; Ericson, 1999). The faceted pterygoid articulation in IGM 100/1418 is markedly different from the projecting knob condition observed in both extant flamingos and grebes, possibly shedding light on the evolution of feeding style in both groups. In birds, the pterygoid articulation of the quadrate plays a crucial role in cranial kinesis, anchoring the pterygoid-palate complex so that forward motion of the quadrate results in an upward movement of the upper mandible relative to the braincase (e.g., Bout and Zweers, 2001, and references therein). Although the precise functional consequences of the difference in shape between IGM 100/1418 and living mirandornithines are unclear without more fossil material and new phylogenetic optimization of these traits, the morphology in the fossil is consistent with that observed in birds that feed by pecking on a hard substrate (Samejima and Otsuka, 1987; Gussekloo, 2005). This form is also seen in the stem flamingo Palaelodus, which exhibited a straight, blunt beak proposed to be adapted for pecking (Mayr, 2015), and in Presbyornis, which is proposed to have been only weakly adapted for filter feeding (Zelenkov and Stidham, 2018). By contrast, extant flamingos are highly specialized filter feeders, and grebes AMERICAN MUSEUM NOVITATES

exhibit a range of beak shapes adapted for crushing small crustaceans and catching fish. The apparent presence of a pecking-consistent morphology prior to the divergence of flamingos and grebes (the new quadrate), as well as early in the flamingo lineage (*Palaelodus*), may provide insight into the feeding ecology ancestral for Mirandornithes.

The fossils described here newly suggest that the shorebird and flamingo/grebe lineages were established in Asia by the beginning of the Eocene. Although the higher-order phylogenetic affinities of both groups remain controversial (Hackett et al., 2008; Jarvis et al., 2014; Prum et al., 2015), the recent investigation of avian phylogenomics by Prum et al. (2015) recovered shorebirds and flamingos/grebes in a clade diverging in the middle Paleocene (~62–58 Ma). This divergence may have occurred in Asia, followed by dispersal of both groups across the rest of the northern hemisphere. However, further material is necessary to assess this hypothesis.

The additional material described here indicates other new taxa from the Bumban Member. The humerus, IGM 100/1360, is most like ralloids (rails and finfoots), and may represent a new gruiform record. While this new humerus cannot be confidently referred to any specific avian clade, it is not consistent with any previously described taxa from Asia and thus likely represents a new species record. The coracoid IGM 100/1371 is the second record of Pangalliformes from this locality. Though the humerus described by Hwang et al. (2010) likely came from a similarly sized individual to that of the new coracoid, a direct comparison cannot be made between these nonoverlapping elements.

These new fossils expand known species richness in Asia in the early Eocene. However, known taxonomic diversity remains relatively low compared to Europe and North America. Like the early Eocene Green River Formation of North America, the Bumbanian fossil record remains dominated by *Presbyornis mongoliensis* (Kurochkin and Dyke, 2010). The fossils reported here are consistent with the "transitional from fluvial to lacustrine" setting proposed for the Bumban Member exposure at Tsagaan Khushuu (Kondrashov, 2004:165), as shore-birds and stem flamingo-grebes, as well as the waterfowl *Presbyornis*, suggest the presence of standing water. Unambiguous presbyornithid remains are not known from Europe (e.g., Mayr, 2009), possibly indicating that the Bumbanian avifauna is more like North American than European faunas. Also like in North America, presbyornithid-like taxa at Naranbulag are known from both the late Cretaceous and the Paleogene (Kurochkin et al., 2002; Clarke and Norell, 2004; Kurochkin and Dyke, 2010), suggesting that this clade survived the K-Pg extinction event in both regions. Recovery of further avian material may inform a fuller reconstruction of both the paleogene.

ACKNOWLEDGMENTS

We thank the members of the joint American Museum of Natural History/Mongolia Academy of Sciences expeditions (2000–2002) for fossil collection; Carl Mehling (AMNH) and Chris Sagabiel (TMM) for specimen access; Sebastian Egberts (UT Austin Vertebrate Paleontology Laboratory) for fossil preparation; M. Colbert and J. Maisano (UTCT) for specimen scanning; N. Adam Smith and an anonymous reviewer for their helpful comments, as well as H. Bilger, N. Crouch, S. Davis, L. Legendre, G. Musser, and undergraduate collaborators in J.A.C.'s lab (UT Austin) for helpful comments and discussion. We thank National Science Foundation grant EAR 0938199 (to J.A.C.) for funding. M.A.N. thanks the AMNH Division of Paleontology and the Macaulay Family endowment.

REFERENCES

- Alvarenga, H.M.F.D. 1988. Ave fossil (Gruiformes: Rallidae) dos folhelhos da bacia de taubate, Estado de Sao Paulo, Brasil. Anais da Academia Brasileira de Ciências 60 (3): 321–332.
- Averianov, A. 1991. Bony-toothed birds (Aves: Pelecaniformes: Odontopterygia) from the Late Paleocene and Eocene of the western margin of ancient Asia. Proceedings of the Zoological Institute, USSR Academy of Sciences 239: 3–12. [in Russian]
- Averianov, A., O.R. Potapova, and L.A. Nessov. 1990. On the first native finds of the bones of ancient birds. Proceedings of the Zoological Institute, USSR Academy of Sciences 210: 3–9. [in Russian]
- Bajpai, S., et al. 2008. The oldest Asian record of Anthropoidea. Proceedings of the National Academy of Sciences of the United States of America 105 (32): 11093–11098.
- Baumel, J.J., and L.M. Witmer. 1993. Osteologia. *In* Handbook of avian anatomy: nomina anatomica avium: 45–132. Cambridge: Museum of Comparative Zoology.
- Bendukidze, O.G. 1971. Novyj prestaviteľ semeistva Geranoididae (Aves, Gruiformes) iz eotsenovykh otlozhenij zaisan. Soobshchenija Akademii nauk Gruzinskoĭ SSR/Bulletin of the Academy of Sciences of the Georgian SSR 63: 749–751.
- Bertelli, S., B.E.K. Lindow, G.J. Dyke, and G. Mayr. 2013. Another charadriiform-like bird from the Lower Eocene of Denmark. Paleontological Journal 47 (11): 1282–1301.
- Bourdon, E., C. Mourer-Chauviré, M. Amaghzaz, and B. Bouya. 2008. New specimens of *Lithoptila abdounensis* (Aves, Prophaethontidae) from the Lower Paleocene of Morocco. Journal of Vertebrate Paleontology 28 (3): 751–761.
- Bout, R.G., and G.A. Zweers. 2001. The role of cranial kinesis in birds. Comparative Biochemistry and Physiology A: Molecular and Integrative Physiology 131 (1): 197–205.
- Bowen, G.J. et al. 2002. Mammalian dispersal at the Paleocene/Eocene boundary. Science 295: 2062-2065.
- Buffetaut, E. 2013. The giant bird *Gastornis* in Asia: a revision of *Zhongyuanus xichuanensis* Hou, 1980, from the Early Eocene of China. Paleontological Journal 47 (11): 1302–1307.
- Cheneval, J., and F. Escuillie. 1992. New data concerning *Palaelodus ambiguus* (Aves: Phoenicopteriformes: Palaelodidae): ecological and evolutionary interpretations. Natural History Museum of Los Angeles County Science Series 36: 209–224.
- Clarke, J.A. 2002. The morphology and systematic position of Ichthyornis Marsh and the phylogenetic relationships of basal Ornithurae. Yale University.
- Clarke, J.A., and M.A. Norell. 2002. The morphology and phylogenetic position of *Apsaravis ukhaana* from the Late Cretaceous of Mongolia. American Museum Novitates 3387: 1–46.
- Clarke, J.A., and M.A. Norell. 2004. New avialan remains and a review of the known avifauna from the Late Cretaceous Nemegt Formation of Mongolia. American Museum Novitates 3447: 1–12.
- Cracraft, J. 1973. Systematics and evolution of the Gruiformes (class Aves). 3. Phylogeny of the suborder Grues. Bulletin of the American Museum of Natural History 151 (1): 1–127.

- Cracraft, J., and J.A. Clarke. 2001. The basal clades of modern birds. *In* J. Gauthier and L.F. Gall (editors), New perspectives on the origin and early evolution of birds: proceedings of the international symposium in honor of John H. Ostrom: 143–156. New Haven: Peabody Museum of Natural History, Yale University.
- Dashzeveg, D. 1988. Holarctic correlation of non-marine Palaeocene–Eocene boundary strata using mammals. Journal of the Geological Society 145 (3): 473–478.
- Dashzeveg, D., J.L. Hartenberger, T. Martin, and S. Legendre. 1998. A peculiar minute glires (Mammalia) from the Early Eocene of Mongolia. *In* K.C. Beard and M.R. Dawson (editors), Dawn of the age of mammals in Asia. Bulletin of Carnegie Museum of Natural History 34: 194–209.
- Elzanowski, A., and T.A. Stidham. 2010. Morphology of the quadrate in the Eocene anseriform *Presbyornis* and extant galloanserine birds. Journal of Morphology 271 (3): 305–323.
- Ericson, P.G.P. 1999. New material of *Juncitarsus* (Phoenicopteriformes), with a guide for differentiating that genus from the Presbyornithidae (Anseriformes). Smithsonian Contributions to Paleobiology 89: 245–51.
- Fürbringer, M. 1888. Untersuchungen zur Morphologie und Systematik der Vögel, zugleich ein Beitrag zur Anatomie der Stütz- und Bewegungsorgane. Amsterdam: T. van Holkema.
- Gussekloo, S.W.S. 2005. Cranial kinesis in palaeognathous birds. Journal of Experimental Biology 208 (17): 3409–3419.
- Hackett, S.J., et al. 2008. A phylogenomic study of birds reveals their evolutionary history. Science 320: 1763–1768.
- Hasegawa, Y., et al. 1979. Preliminary notes on the Oligo-Miocene penguin-like birds from Japan (parts I–VII). Kitakyūshū-shi Shizenshi Hakubutsukan kenkyū hōkoku/Bulletin of the Kitakyushu Museum of Natural History 1: 41–60. [in Japanese]
- Hou, L. 1980. New form of the Gastornithidae from the Lower Eocene of the Xichuan, Honan. Vertebrata PalAsiatica 18: 111–15. [in Chinese]
- Hou, L. 1982. New form of the Threskiornithidae from the Upper Eocene of the Minggang, Henan. Vertebrata PalAsiatica 20: 196–202. [in Chinese]
- Hou, L. 1989. A Middle Eocene bird from Sangequan, Xinjiang. Vertebrata PalAsiatica 27: 65–70. [in Chinese]
- Hou, L. 1990. An Eocene bird from Songzi, Hubei province. Vertebrata PalAsiatica 28 (1): 34-42. [in Chinese]
- Hou, L. 1994. A new Paleocene bird from Anhui, China. Vertebrata PalAsiatica 32: 60-65. [in Chinese]
- Hou, L., and P.G.P. Ericson. 2002. A Middle Eocene shorebird from China. Condor 104 (4): 896-899.
- Hwang, S.H., G. Mayr, and M. Bolortsetseg. 2010. The earliest record of a galliform bird in Asia, from the Late Paleocene–Early Eocene of the Gobi Desert, Mongolia. Journal of Vertebrate Paleontology 30 (5): 1642–1644.
- Jarvis, E.D., et al. 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. Science 346: 1320–1331.
- Kondrashov, P.E. 2004. A new hyopsodontid (Mammalia, Condylartha) from the Early Eocene of Mongolia. Bulletin New Mexico Museum of Natural History 26: 165–168.
- Kondrashov, P.E., A.V. Lopatin, and S.G. Lucas. 2004. The oldest known Asian artiodactyl (Mammalia). Bulletin New Mexico Museum of Natural History and Science 26: 205–208.
- Kozlova, E.V. 1960. New fossil birds from Southeastern Gobi. Trudy Problemnykh i Tematicheskikh Soveshchanii/Zoologicheskii Institut (Akademiia Nauk SSSR) 9: 323–329. [in Russian]

- Kurochkin, E.N. 1968. New Oligocene birds from Kazakhstan. Paleontologicheskii Zhurnal 92–101. [in Russian]
- Kurochkin, E.N. 1976. A survey of the Paleogene birds of Asia. *In* S.L. Olson (editor), Collected papers in avian paleontology honoring the 90th birthday of Alexander Wetmore. Smithsonian Contributions to Paleobiology 27: 75–86.
- Kurochkin, E.N. 1981. New representatives and evolution of two archaic Gruiform families in Eurasia. Trudy Sovmestnaja Sovetsko-Mongol-Skaja Paleontologicheskaja Ekspeditsija 15: 59–85. [in Russian]
- Kurochkin, E.N., and G.J. Dyke. 2010. A large collection of *Presbyornis* (Aves, Anseriformes, Presbyornithidae) from the Late Paleocene and Early Eocene of Mongolia. Geological Journal 45: 375–387.
- Kurochkin, E.N., and G.J. Dyke. 2011. The first fossil owls (Aves: Strigiformes) from the Paleogene of Asia and a review of the fossil record of Strigiformes. Paleontological Journal 45 (4): 445–458.
- Kurochkin, E.N., G.J. Dyke, and A.A. Karhu. 2002. A new presbyornithid bird (Aves, Anseriformes) from the Late Cretaceous of southern Mongolia. American Museum Novitates 3386: 1–11.
- Lambrecht, K. 1931. Protoplotus beauforti n.g. n. sp., ein Schlangenhalsvogel aus dem Tertiär von W.-Sumatra. Wetenschappelijke Mededeelingen Dienst van den Mijnbouw in Nederlandisch-Oost Indië 17: 15–24.
- Louchart, A., C. Mourer-Chauviré, P. Vignaud, H.T. MacKaye, and M. Brunet. 2005. A finfoot from the Late Miocene of Toros Menalla (Chad, Africa): palaeobiogeographical and palaeoecological implications. Palaeogeography, Palaeoclimatology, Palaeoecology 222 (1–2): 1–9.
- Mayr, G. 2000. A new basal galliform bird from the Middle Eocene of Messel (Hessen, Germany). Senckenbergiana Lethaea 80 (1): 45–57.
- Mayr, G. 2004. Phylogenetic relationships of the Early Tertiary Messel rails (Aves, Messelornithidae). Senckenbergiana Lethaea 84 (1): 317–322.
- Mayr, G. 2006. New specimens of the Early Eocene stem group galliform *Paraortygoides* (Gallinuloididae), with comments on the evolution of a crop in the stem lineage of Galliformes. Journal of Ornithology 147 (1): 31–37.
- Mayr, G. 2007. The birds from the Paleocene fissure filling of Walbeck (Germany). Journal of Vertebrate Paleontology 27 (2): 394–408.
- Mayr, G. 2009. Paleogene fossil birds. Berlin: Springer.
- Mayr, G. 2014. The Eocene *Juncitarsus* its phylogenetic position and significance for the evolution and higher-level affinities of flamingos and grebes. Comptes Rendus Palevol 13 (1): 9–18.
- Mayr, G. 2015. Cranial and vertebral morphology of the straight-billed Miocene phoenicopteriform bird *Palaelodus* and its evolutionary significance. Zoologischer Anzeiger 254: 18–26.
- Mayr, G. 2016. The world's smallest owl, the earliest unambiguous charadriiform bird, and other avian remains from the Early Eocene Nanjemoy Formation of Virginia (USA). Paläontologische Zeitschrift 90 (4): 747–763.
- Mayr, G. 2017. A small, "wader-like" bird from the Early Eocene of Messel (Germany). Annales de Paléontologie 103 (2): 141–147.
- Mayr, G., and J.A. Clarke. 2003. The deep divergences of neornithine birds: a phylogenetic analysis of morphological characters. Cladistics 19 (6): 527–553.
- Mayr, G., and R.P. Scofield. 2016. New avian remains from the Paleocene of New Zealand: the first Early Cenozoic Phaethontiformes (Tropicbirds) from the southern hemisphere. Journal of Vertebrate Paleontology 36 (1).

- Mayr, G., and R. Smith. 2002. Avian remains from the lowermost Oligocene of Hoogbutsel (Belgium). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre 72: 139–150.
- Mayr, G., and T. Smith. 2013. Galliformes, Upupiformes, Trogoniformes, and other avian remains (?Phaethontiformes and ?Threskiornithidae) from the Rupelian stratotype in Belgium, with comments on the identity of "Anas" benedeni Sharpe, 1899. In U.B. Göhlich and A. Kroh (editors), Paleornithological research 2013. Proceedings of the 8th International Meeting of the Society of Avian Paleontology and Evolution: 23–35. Vienna: Natural History Museum Vienna, 306 pp.
- Mayr, G., and I. Weidig. 2004. The Early Eocene bird *Gallinuloides wyomingensis*—a stem group representative of Galliformes. Acta Palaeontologica Polonica 49 (2): 211–217.
- Mayr, G., and E. Zvonok. 2011. Middle Eocene Pelagornithidae and Gaviiformes (Aves) from the Ukrainian Paratethys. Palaeontology 54 (6): 1347–1359.
- Mayr, G., and E. Zvonok. 2012. A new genus and species of Pelagornithidae with well-preserved pseudodentition and further avian remains from the Middle Eocene of the Ukraine. Journal of Vertebrate Paleontology 32 (4): 914–925.
- Mayr, G., R.S. Rana, A. Sahni, and T. Smith. 2007. Oldest fossil avian remains from the Indian subcontinental plate. Current Science 92 (9): 1266–1269.
- Mayr, G., et al. 2010. *Quercypsitta*-like birds from the Early Eocene of India (Aves, ?Psittaciformes). Journal of Vertebrate Paleontology 30 (2): 467–478.
- Mayr, G., J. Yang, E. De Bast, C. Li, and T. Smith. 2013a. A *Strigogyps*-like bird from the Middle Paleocene of China with an unusual grasping foot. Journal of Vertebrate Paleontology 33 (4): 895–901.
- Mayr, G., et al. 2013b. New specimens of the Early Eocene bird *Vastanavis* and the interrelationships of stem group Psittaciformes. Paleontological Journal 47 (11): 1308–1314.
- Meng, J., and M.C. McKenna. 1998. Faunal turnovers of Palaeogene mammals from the Mongolian plateau. Nature 394: 364–367.
- Milne-Edwards, A. 1863. Mémoire sur la distribution géologique des oiseaux fossiles et description de quelques espèces nouvelles. Annales des Sciences Naturelles. 4 (20): 132–176.
- Mlikovsky, J., and P. Svec. 1986. Review of the Tertiary waterfowl (Aves: Anseridae) of Asia. Věstník Československé Společnosti Zoologické/Acta Societatis Zoologicae Bohemoslovenicae 50: 249–72.
- Mourer-Chauviré, C. 1992. The Galliformes (Aves) from the Phosphorites Du Quercy (France): systematics and biostratigraphy. Natural History Museum of Los Angeles County, Science Series 36: 67–95.
- Mourer-Chauviré, C. 2000. A new species of *Ameripodius* (Aves: Galliformes: Quercymegapodiidae) from the Lower Miocene of France. Palaeontology 43 (3): 481–493.
- Nesbitt, S.J., A.H. Turner, M. Spaulding, J.L. Conrad, and M.A. Norell. 2009. The theropod furcula. Journal of Morphology 270 (7): 856–879.
- Nessov, L.A. 1986. The first record of the Late Cretaceous bird *Ichthyornis* in the Old World and some other bird bones from the Cretaceous and Paleogene of Central Asia. Proceedings of the Zoological Institute, USSR Academy of Sciences 147: 31–38. [in Russian]
- Nessov, L.A. 1988. New Cretaceous and Paleocene birds of Soviet Union and Kazakhstan and their environments. Proceedings of the Zoological Institute, Leningrad 182: 116–123. [in Russian]
- Nessov, L.A. 1992. Mesozoic and Paleogene birds of the USSR and their paleoenvironments. Natural History Museum of Los Angeles County, Science Series 36: 465–478.
- Norell, M.A., and J.A. Clarke. 2001. Fossil that fills a critical gap in avian evolution. Nature 409: 181–184. Olson, S.L. 1974. *Telecrex* restudied: a small Eocene guineafowl. Wilson Bulletin 86 (3): 246–250.
- Olson, S.L., and A. Feduccia. 1980. Relationships and evolution of flamingos (Aves: Phoenicopteridae). Smithsonian Contributions to Zoology 316. Washington D.C.: Smithsonian Institution Press.

- Olson, S.L., and Y. Hasegawa. 1996. A new genus and two new species of gigantic Plotopteridae from Japan (Aves: Pelecaniformes). Journal of Vertebrate Paleontology 16 (4): 742–751.
- Olson, S.L., and D.C. Parris. 1987. The Cretaceous birds of New Jersey. Smithsonian Contributions to Paleobiology 63: 1–22.
- Panteleyev, A.V., and L.A. Nessov. 1987. A small tubinare (Aves: Procellariiformes) from the Eocene of Middle Asia. Trudy Zoologicheskogo Instituta 252: 95–103. [in Russian]
- Peters, D.S. 1987. *Juncitarsus merkeli* n. sp. stützt die Ableitung der Flamingos von Regenpfeifervögeln (Aves: Charadriiformes: Phoenicopteridae). Courier Forschungsinstitut Senckenberg 97: 141–55.
- De Pietri, V.L., and G. Mayr. 2014. Reappraisal of Early Miocene rails (Aves, Rallidae) from central France: diversity and character evolution. Journal of Zoological Systematics and Evolutionary Research 52 (4): 312–322.
- Prum, R.O., et al. 2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. Nature 526: 569–573.
- Sakurai, K., M. Kimura, and T. Katoh. 2008. A new penguin-like bird (Pelecaniformes:Plotopteridae) from the Late Oligocene Tokoro Formation, northeastern Hokkaido, Japan. Oryctos 7: 83–94.
- Samejima, M., and J. Otsuka. 1987. Observations on the quadrate of birds. Journal of Ornithology 35 (4): 129–144. [in Japanese]
- Smith, N.A. 2014. The fossil record and phylogeny of the auklets (Pan-Alcidae, Aethiini). Journal of Systematic Palaeontology 12: 217–236.
- Smith, N.A. 2015. Sixteen vetted fossil calibrations for divergence dating of Charadriiformes (Aves, Neognathae). Palaeontologia Electronica: PE 18.1.4FC. [https://doi.org/10.26879/410]
- Smith, T., K.D. Rose, and P.D. Gingerich. 2006. Rapid Asia-Europe-North America geographic dispersal of earliest Eocene primate *Teilhardina* during the Paleocene-Eocene Thermal Maximum. Proceedings of the National Academy of Sciences 103 (30): 11223–11227.
- Stidham, T.A., and X. Ni. 2014. Large anseriform (Aves: Anatidae: Romainvilliinae?) fossils from the Late Eocene of Xinjiang, China. Vertebrata PalAsiatica 52 (1): 98–111.
- Stidham, T.A., et al. 2005. An ibis-like bird (Aves: cf. Threskiornithidae) from the Late Middle Eocene of Myanmar. Contributions from the Museum of Paleontology, the University of Michigan 31 (7): 179–184.
- van Tets, G.F., P.V. Rich, and H.R. Marino-Hadiwardoyo. 1989. A reappraisal of *Protoplotus beauforti* from the Early Tertiary of Sumatra and the basis of a new Pelecaniform family. Publication of the Geological Research and Development Centre, Indonesia. Paleontology Series 5: 57–75.
- Tugarinov, A. 1940. New data on the Tertiary ornithofauna of the USSR. Doklady Akademii nauk SSSR 26: 314–16.
- Wang, M., G. Mayr, J. Zhang, and Z. Zhou. 2012a. Two new skeletons of the enigmatic, rail-like avian taxon *Songzia* Hou, 1990 (Songziidae) from the Early Eocene of China. Alcheringa 36 (4): 487–499.
- Wang, M., G. Mayr, J. Zhang, and Z. Zhou. 2012b. New bird remains from the Middle Eocene of Guangdong, China. Acta Palaeontologica Polonica 57 (3): 519–526.
- Wetmore, A. 1934. Fossil birds from Mongolia and China. American Museum Novitates 711: 1-16.
- Witmer, L.M. 1990. The craniofacial air sac system of Mesozoic birds (Aves). Zoological Journal of the Linnean Society 100: 327–378.
- Xue, X. 1995. A Paleocene bird discovered in China. Courier Forschungsinstitut Senckenberg 181: 89–93.
- Zelenkov, N.V. 2013. Cenozoic phoenicopteriform birds from Central Asia. Paleontological Journal 47 (11): 1323–1330.

- Zelenkov, N.V. 2018. The earliest Asian duck (Anseriformes: *Romainvillia*) and the origin of Anatidae. Doklady Biological Sciences 483 (1): 225–227.
- Zelenkov, N.V. 2019. A swan-sized anseriform bird from the Late Paleocene of Mongolia. Journal of Vertebrate Paleontology. In press.
- Zelenkov, N.V., and A.V. Panteleyev. In press. A small stem-galliform bird (Aves: Paraortygidae) from the Eocene of Uzbekistan. Comptes Rendus Palevol.
- Zelenkov, N.V., and T.A. Stidham. 2018. Possible filter-feeding in the extinct *Presbyornis* and the evolution of Anseriformes (Aves). Zoologicheskiĭ Zhurnal 97 (8): 943–956.
- Zvonok, E.A., G. Mayr, and L. Gorobets. 2015. New material of the Eocene marine bird *Kievornis* Averianov et al., 1990 and a reassessment of the affinities of this taxon. Vertebrata PalAsiatica 53 (3): 238–244.
- Zvonok, E.A., N.V. Zelenkov, and I.G. Danilov. 2016. A new unusual waterbird (Aves, ?Suliformes) from the Eocene of Kazakhstan. Journal of Vertebrate Paleontology 36 (1): e1035783.

All issues of *Novitates* and *Bulletin* are available on the web (http://digitallibrary. amnh.org/dspace). Order printed copies on the web from: http://shop.amnh.org/a701/shop-by-category/books/scientific-publications.html

or via standard mail from:

American Museum of Natural History—Scientific Publications Central Park West at 79th Street New York, NY 10024

⊗ This paper meets the requirements of ANSI/NISO Z39.48-1992 (permanence of paper).