

AMERICAN MUSEUM *Novitates*

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, N.Y. 10024
Number 3170, 50 pp., 12 figures

May 23, 1996

Catalog of Recent Type Specimens in the Department of Invertebrates, American Museum of Natural History. II. Mollusca Part 1 (Classes Aplacophora, Polyplacophora, Gastropoda [Subclass Opisthobranchia], Bivalvia, and Scaphopoda)

CHRISTOPHER B. BOYKO¹ AND WALTER E. SAGE III²

ABSTRACT

A complete, annotated listing of all type specimens of four classes and one subclass of mollusks in the collection of the Department of Invertebrates at the American Museum of Natural History (as of December 1995) is given. A total of 271 type lots from the five groups is listed. Eight-

een lectotypes are designated herein, and illustrations are provided for one lectotype and 11 other important type specimens, including several holotypes thought to have been lost. An index of genera and species is included for each group.

INTRODUCTION

The Department of Invertebrates at the American Museum of Natural History currently houses a collection of over 350,000

cataloged lots divided among three major categories: fossil (45,000 lots), Recent mollusks (275,000 lots), and other Recent phyla

¹ Department of Zoology, University of Rhode Island, Kingston, Rhode Island 02881-0816.

² Formerly Senior Scientific Assistant, Department of Invertebrates, American Museum of Natural History (deceased).

(30,000 lots). All of the types are housed separately from the main body of the collection. This allows for a greater level of security than can be maintained in a large, general collection area as per ICZN Recommendation 72 (ICZN, 1985: 147).

This catalog is the second in a series of comprehensive listings of all the type material in the department's collection. It is also an updating and expansion of the work of Richards and Old (1969, in part), which contains many inaccuracies and omissions. Included here are all AMNH holotype, lectotype, paratype, paralectotype, and syntype specimens from four classes and one subclass of mollusks: Aplacophora (12 species), Polyplacophora (8 species), Opisthobranchia (22 species), Bivalvia (122 species), and Scaphopoda (36 species). A total of 54 holotype, 19 lectotype, 135 paratype, 45 paralectotype, and 18 syntype AMNH lots is listed in this publication.

This catalog includes an accounting of the type specimens that have been transferred to this institution from other sources. Very often no published record of these transfers is ever made, especially for species coming from private collections, such as those listed below. In the case of several lots, types were published as having been given to the American Museum, but subsequent investigation indicated that the specimens were never deposited. Other members of type series with representatives in AMNH have also been subject to transfer between institutions. The most recent known repositories have been listed, along with the most recent catalog numbers used to identify type lots in other institutions.

The majority of the type specimens included in this catalog come from a few key sources. The two most important pre-1900 collections are those of John C. Jay and William A. Haines. The Jay Collection, purchased for \$10,000 in 1874, was donated to the AMNH and contains type specimens of species described by both Jay and others, such as Isaac Lea, with whom Jay traded specimens. Part of the Haines Collection was donated by Haines himself, the remainder being purchased from his son in 1895. Haines described very few species, but received many type specimens from Isaac Lea and others.

For more information on these and other large collections that have been incorporated into the AMNH mollusk collection, see Richards and Old (1969).

More recent sources of type material included in this catalog are the AMNH Belgian Congo Expedition of 1909–1915, with many new species collected by Herbert Lang and James Chapin and described by Pilsbry and Bequaert (1927), the "Vema" Expedition (1955–1962), with new species described by Luitfried von Salvini-Plawen, Victor Scarabino, and Ernst and Eveline Marcus, and the Gordon Nowell-Usticke Collection, which has been discussed in detail by M. J. Faber (1988).

The AMNH collection has been organized by phylum and "superphylum" (Protozoa). Each division is given a separate series of catalog numbers. As a result of this system of numbering, a chiton may have the same catalog number as a trematode, but the specimens cannot be confused because of the taxonomic distinction between phyla.

Each potential type lot was compared to the original description and figures in order to verify type status, confirm locality and collection data and remove any specimens from the type collection that are not valid types. The publication of this catalog fulfills ICZN Recommendation 72G(4), which states that all museums holding type specimens have the responsibility of making a published catalog of these types available to the scientific community (ICZN, 1985: 147).

This catalog is arranged by class, based primarily on Vaught, 1989. Within each class, the types are listed in alphabetical order by genus and species. An index is provided, which gives a genus- and species-level list within each phylum. This catalog adds to the format of Boyko (1994) by including detailed information on current synonymy and generic placement of all species listed.

One critical area addressed in this catalog is the ambiguous type designations and other type concept problems introduced by Salvini-Plawen, E. Marcus, and Usticke. Although there has been some indication that the species introduced by Salvini-Plawen (1986) should, at least in part, be validated from that date (Bayer, personal commun.), we and others (A. Scheltema, personal commun.; Petit,

personal commun.) believe all species from the 1986 abstract are nomena nuda because there is no way to distinguish them based on the limited or absent characters given. Salvini-Plawen (1992) described all the 1986 species in detail and the names should date from the 1992 paper. Salvini-Plawen (1992) also stated that "all types are deposited in the American Museum." This statement refers to all the holotypes from the 1992 paper, because not all the specimens in each type series were deposited at the AMNH.

Similarly, E. Marcus (1958b) stated that "the types and some of the other material in this article are deposited in the invertebrate section of the American Museum of Natural History." This statement should be read "the [holo]types and some of the other material [including some paratypes] . . ." as, again, not all the specimens in each type series were deposited at the AMNH.

Usticke (1959) unambiguously designated the "pictured" specimen for each new species or subspecies as holotype. However, in his 1969 paper, Usticke did not include any holotype designations and, therefore, all figured specimens from the 1969 paper are syntypes. Usticke (1971) published measurements of all holotypes which, in most cases, correspond to the figured specimens. We refer to this author as "Usticke" rather than "Nowell-Usticke" because the former appellation is how he referred to himself in his own papers.

In the case of 67 species treated herein, no holotype was designated in the original publication. For 19 of these species, one specimen in the AMNH collection is clearly either the figured specimen in the original publication or was labeled by the author of the species as being the holotype. Such specimens not subsequently published as holotype or lectotype, are herein designated lectotype (any postpublication reference to a holotype where none was originally designated is considered the equivalent of a lectotype designation). One of these specimens is figured along with 11 holotypes and syntypes that either were never illustrated (e.g., *Batissa gigantea*) or were cited as lost in other publications (e.g., *Unio callosus*). In addition, one U.S. National Museum specimen is designated lectotype for one species of bivalve (*Iridina coelestis*).

A list is also provided of 45 species that should be represented by type (or possible type) specimens at the AMNH, but for whom types cannot be traced in the collection or the records. Most of the bivalve specimens were either stated or implied to be in collections later transferred to the AMNH, primarily those of Jay and Haines. Since there is no record of these specimens having arrived here, and they have not been located at other major institutions, it is suspected that they are lost. One species on this list (*Chlamys incantata*) was explicitly cited as having been deposited at the AMNH but never arrived. It is of interest to note that only one bivalve holotype from all species described in Pilsbry and Bequaert (1927) is not located in either AMNH, ANSP, or MCZ. This is *Aspatharia chapini*, which has been traced to the Museum of Central Africa at Tervuren, Belgium.

The list of missing opisthobranchs is based on a letter found in the department files from Eveline and Ernst Marcus, (dated 1962, to Libbie Hyman at the AMNH) which includes a list of 38 lots of opisthobranchs sent to AMNH for deposition. While it is possible that these specimens were not types, the Marcus stated that "if the labels are compared with the mentioned publication, one can see whether there are types or not," which would indicate that at least some of these specimens were types. Unfortunately, only seven lots from this list can be found at the AMNH; they are listed in the main opisthobranch section. The remaining lots are presumed lost or destroyed and, since the list refers specific questions on lot size, locality, and collection dates to the individual specimen labels, this information is lost as well.

Each type listing contains four to six categories of data. Within each genus, specific names are followed by authorship data (author, year description was published, and page and figure numbers). The next category indicates the specific kind of type(s) in the collection, as well as AMNH catalog number(s), and the number of specimens in each lot. The locality category contains all pertinent collection data, including county names for U.S. locations where that information can be determined. The collector category indicates the collector(s) of the specimen(s) and the date(s) they were collected, if those data are known.

The remarks category contains information as to provenance of specimens, location of other types in the series (all of the members of each type series are listed as could be located), current synonymy and/or generic placement, etc. The host category is included for species that are parasites or commensals of other species.

| | |
|------|---|
| URI | University of Rhode Island, Department of Zoology, Kingston |
| USNM | United States National Mu- seum (now National Muse- um of Natural History— NMNH), Washington, D.C. |
| WHOI | Woods Hole Oceanographic Institute, Massachusetts |

Institutional Abbreviations

| | |
|--------|---|
| AMNH | American Museum of Natural History, New York |
| ANSP | Academy of Natural Sciences of Philadelphia, Pennsylvania |
| BM(NH) | British Museum (Natural History) (now The Natural History Museum), London |
| CASIZ | California Academy of Sciences, Invertebrate Zoology, Berkeley |
| DMNH | Delaware Museum of Natural History, Wilmington |
| FMNH | Field Museum of Natural History, Chicago |
| LACM | Los Angeles County Museum of Natural History, California |
| MCZ | Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts |
| MHNH-P | Muséum National d'Histoire Naturelle, Paris, France |
| MHNH-M | Museo Nacional de Historia Natural, Montevideo, Uruguay |
| MRAC | Musée Royal de L'Afrique Centrale, Tervuren, Belgium |
| NHMV | National History Museum, Vienna, Austria |
| NSMT | National Science Museum, Tokyo |
| NYZS | New York Zoological Society |
| SBMNH | Santa Barbara Museum of Natural History, California |
| SDNHM | San Diego Natural History Museum, California |
| UMML | University of Miami Marine Laboratory, Florida |
| UMMZ | University of Michigan Museum of Zoology, Ann Arbor |

ACKNOWLEDGMENTS

We would like to thank the following for their help in providing much needed information and assistance: Elizabeth Kools (CASIZ), Luitfried von Salvini-Plawen (NHM Vienna), Ruth Turner and Richard Johnson (MCZ), Gary Rosenberg (ANSP), Firmin Puylaert (Royal Museum of Central Africa), Roland Houart, Louis and Lorraine Schwartz, Lindsey Groves (LACM), John Slapcinsky and Rüdiger Bieler (FMNH), Alan Kabat (USNM) and Paul Scott (SBMNH). Additional thanks to Richard E. Petit, Gene Coan, Alan Harvey (AMNH) and Robert Bullock (URI) for reviewing the manuscript as well as Terrence Gosliner (CASIZ) and Paula Mikkelsen (DMNH) (Opisthobranchia), William G. Lyons (Polyplacophora), Amélie Scheltema (WHOI) (Aplacophora), and William K. Emerson (AMNH) (Scaphopoda) for reviewing the sections in their respective areas of expertise. Thanks also to Andrew Model (AMNH) for the photography and to Alan Harvey and Deborah Levine for assistance with the illustrations.

CLASS APLACOPHORA

| |
|--|
| <i>Falcidens (Lepoderma) acutargatus</i> Salvini- Plawen, 1992: 322, figs. 8, 18, 23c. Holotype: AMNH 265333 . Paratypes: AMNH 265346 (3 specimens). Locality: V-15-22, 09°46.3'N, 79°37.5'W, 974 m, Gulf of Darien, Panama, Atlantic Ocean. Collector: R/V "Vema," Nov. 10, 1958. |
| <i>Falcidens limifossorides</i> Salvini-Plawen, 1992: 321, figs. 6, 16, 23a. Holotype: AMNH 265330 . Paratypes: AMNH 265335 (1 specimen), 265337 (2 specimens), 265338 (1 speci- men), 265339 (3 specimens), 265340 (1 specimen). |

Locality: V-17-5, 38°15'S, 76°00'W, 3739–3824 m (AMNH 265330), V-17-6, 37°57'S, 75°08'W, 4303–4323 m (AMNH 265340), off Chile; V-15-38, 05°00'N, 79°04'W, 3023–3251 m (AMNH 265335), V-15-40, 04°25'N, 78°25'W, 3700–3757 m (AMNH 265337), off Colombia; V-15-42, 07°55'N, 86°00'W, 3087–3091 m (AMNH 265338), V-15-59, 07°05'N, 85°55'W, 2853–2855 m (AMNH 265339), off Costa Rica, Pacific Ocean.

Collector: R/V "Vema," Nov. 15 (AMNH 265335), Nov. 16 (AMNH 265337), Nov. 19 (AMNH 265338), Nov. 30 (AMNH 265339), 1958; March 16 (AMNH 265330), March 21 (AMNH 265340), 1961.

Remarks: Additional paratypes will be deposited in NHMV (Salvini-Plawen, personal commun.). This specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Falcidens (Lepoderma) nontargatus Salvini-Plawen, 1992: 322, figs. 19, 23d.

Holotype: AMNH 265334.

Locality: V-14-32, 34°35'S, 17°31'E, 1861 m, off western cape of Africa, Atlantic Ocean.

Collector: R/V "Vema," April 6, 1958.

Remarks: AMNH 265334 is the holotype by monotypy.

Falcidens procerus Salvini-Plawen, 1992: 321–322, figs. 9, 22, 23e.

Holotype: AMNH 265331.

Paratypes: AMNH 265344 (3 specimens), 265345 (5 specimens).

Locality: V-17-2, 34°29'S, 74°21'W, 4030–4036 m, off Chile, (AMNH 265331, 265344); V-15-74, 07°32'S, 81°26'W, 5759–5760 m, off Peru (AMNH 265345), Pacific Ocean.

Collector: R/V "Vema," Dec. 19, 1958 (AMNH 265345), March 13, 1961 (AMNH 265331, 265344).

Remarks: Additional paratypes will be deposited in NHMV (Salvini-Plawen, personal commun.). This specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Falcidens (Lepoderma) targatus Salvini-Plawen, 1992: 322, figs. 7, 17, 23b.

Holotype: AMNH 265332.

Paratypes: AMNH 265336 (23 specimens).

Locality: V-17-102, 34°25'S, 52°19'W, 73 m, off Uruguay, Atlantic Ocean.

Collector: R/V "Vema," June 27, 1961.

Remarks: Additional paratypes will be deposited in NHMV (Salvini-Plawen, personal commun.). This species is the type of the subgenus *Lepoderma* Salvini-Plawen, 1992, by original designation (Salvini-Plawen, 1992: 322). This specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Limifossor elongatus Salvini-Plawen, 1972: 47–49, figs. 28–31.

Lectotype: AMNH 265351.

Paralectotypes: AMNH 265355 (1 specimen), 265356 (3 specimens).

Locality: V-15-46, 09°22'N, 89°33'W, 3517–3528 m (AMNH 265351), off Costa Rica; V-15-41, 07°02'N, 83°10'W, 2269–2270 m (AMNH 265355), off Panama; V-15-63, 03°15'S, 82°30'W, 2861–2864 m (AMNH 265356), off Ecuador, Pacific Ocean.

Collector: R/V "Vema," Nov. 18 (AMNH 265355), Nov. 20 (AMNH 265351), Dec. 4 (AMNH 265356), 1958.

Remarks: AMNH 265351 was labeled "type" by Salvini-Plawen. Since Salvini-Plawen did not designate a holotype, AMNH 265351 is here designated lectotype. Additional paralectotypes will be deposited in NHMV (Salvini-Plawen, personal commun.). This species is the type of the genus *Psilodens* Salvini-Plawen, 1977, by original designation (Salvini-Plawen, 1977: 415).

Limifossor hexadentatus Salvini-Plawen; 1992: 321, figs. 5, 13.

Holotype: AMNH 265327.

Paratypes: AMNH 265343 (3 specimens).

Locality: V-18-09, 36°17'S, 53°21'W, 547–676 m, Rio de la Plata, Uruguay, Atlantic Ocean.

Collector: R/V "Vema," Feb. 4, 1962.

Remarks: The additional paratype will be deposited in NHMV (Salvini-Plawen, personal commun.). This specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Limifossor holopeltatus Salvini-Plawen, 1992: 321, figs. 4, 15.

Holotype: AMNH 265326.

Paratypes: AMNH 265341 (3 specimens).

Locality: V-17-20, 53°21'S, 70°36'W, 247 m, Strait of Magellan, Chile, Pacific Ocean.

Collector: R/V "Vema," March 29, 1961.

Remarks: Additional paratypes will be deposited in NHMV (Salvini-Plawen, personal commun.). This specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Limifossor profundus Salvini-Plawen, 1992: 318, figs. 3, 14.

Holotype: AMNH 265325.

Paratype: AMNH 265342 (1 specimen).

Locality: V-15-36, 07°30'N, 79°21'W, 1609–1746 m (AMNH 265325); V-15-35, 07°30'N, 79°16'W, 2234 m (AMNH 265342), off Panama, Pacific Ocean.

Collector: R/V "Vema," Nov. 14, 1958.

Remarks: Additional paratypes will be deposited in NHMV (Salvini-Plawen, personal commun.). This specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Prochaetoderma (Lonchoderma) longisquamosum Salvini-Plawen, 1992: 324, fig. 25.

Holotype: AMNH 265328, 265329.

Locality: V-16-15, 45°00'S, 45°46'E, 1622 m, between Prince Edward and Crozet Islands, Indian Ocean.

Collector: R/V "Vema," Jan. 8, 1960.

Remarks: AMNH 265328, 265329 is the holotype by monotypy. The body of the specimen is AMNH 265328 and the radula is AMNH 265329. This species is the type of the subgenus *Lonchoderma* Salvini-Plawen, 1992, by monotypy (Salvini-Plawen, 1992: 324). The specific name is a nomen nudum from Salvini-Plawen, 1986: 76 (Petit, personal commun.; A. Scheltema, personal commun.).

Scutopus chilensis Salvini-Plawen, 1972: 45, figs. 23, 25a, 26, 27.

Lectotype: AMNH 265349.

Paralectotype: AMNH 265350 (1 specimen [part]).

Locality: V-17-27, 53°33.8'S, 70°17.5'W, 263 m, Strait of Magellan, Chile, Pacific Ocean.

Collector: R/V "Vema," March 29, 1961.

Remarks: AMNH 265349 was labeled "type" by Salvini-Plawen. Since Salvini-Plawen did not designate a holotype, AMNH 265349 is here designated lectotype. The paralectotype [part] is a radula slide. Additional paralectotypes as well as the body of AMNH 265350 will be deposited in NHMV (Salvini-Plawen, personal commun.).

Scutopus megaradulatus Salvini-Plawen, 1972: 44, figs. 21, 22, 24, 25b.

Lectotype: AMNH 265347.

Locality: V-15-22, 09°46.3'N, 79°37.5'W, 974 m, Gulf of Darien, Panama, Atlantic Ocean.

Collector: R/V "Vema," Nov. 10, 1958.

Remarks: AMNH 265347 was labeled "type" by Salvini-Plawen. Since Salvini-Plawen did not designate a holotype, AMNH 265347 is here designated lectotype. The paralectotype will be deposited in NHMV (Salvini-Plawen, personal commun.).

CLASS POLYPLACOPHORA

Callistochiton leei Ferreira, 1979: 451–452, figs. 7–12.

Paratype: AMNH 183817 (1 specimen).

Locality: Low tide, less than 1 m depth, about the midpoint on the east side of the island, 29°01'N, 118°13'W, Sealer's Camp, Guadalupe Island, Mexico, Pacific Ocean.

Collector: W. L. Lee and A. J. Ferreira, Jan. 2, 1975.

Remarks: ex Ferreira. The holotype is CASIZ 020232 and additional paratypes are CASIZ 020305, 064068, LACM 1905, USNM 749083, and ANSP 346418 (Ferreira, 1979: 452).

Chaetopleura staphylophera Lyons, 1985: 35–38, figs. 1–14, 28.

Paratype: AMNH 213570 (1 specimen).

Locality: Trawled in 53 m, 30 nautical miles ENE of Melbourne, Florida, 28°10'N, 80°03'W, United States, Atlantic Ocean.

Collector: J. Johnson, Sept. 1981.

Remarks: ex Lyons (Johnson Collection). The holotype is USNM 842109 and additional paratypes are in several institutions (Lyons, 1985: 35–37) as well as ANSP (Lyons, personal commun.).

Chiton viridis var. *rubrocarinata* Usticke, 1969: 7, fig. 1860.

Lectotype: AMNH 198488 (1 specimen).

Paralectotypes: AMNH 270409 (2 specimens).

Locality: La Grange Beach, St. Croix, Virgin Islands, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Usticke Collection 1860 (AMNH 198488) and L. and L. Schwartz (Usticke Collection) (AMNH 270409). AMNH 198488 was the figured specimen (Usticke, 1969) and is here designated lectotype. This subspecific name is a nomen nudum from Usticke, 1959: 90 as "Var. *Chiton viridis rubrocarinatus*." Although the term "var." was employed, all other "varieties" described by Usticke (1969) were called "new s[ub] sp[ecies]" or, in error, "new sp[ecies]." This taxon, described with an infrasubspecific term after 1960, should nevertheless be considered subspecific because it was not published expressly as a "variety" in the traditional sense of the word. This taxon was erroneously cited as "*Chiton rubrocarinata*" in the figure caption. This subspecies is a synonym of *C. viridis* Spengler, 1797 (Usticke, 1971: 31; Faber, 1988: 89).

Ischnochiton dubium Usticke, 1969: 7, fig. 1869.

Lectotype: AMNH 198489 (1 specimen).

Locality: Pointe des flèches, St. Croix, Virgin Islands, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Usticke collection 1869. AMNH 198489 was the figured specimen (Usticke, 1969) and is here designated lectotype. This species is a synonym of *Stenoplax (S.) boogii* (Haddon, 1886) (Kaas and Van Belle, 1987: 144).

Ischnochiton macleani Ferreira, 1978a: 36–39, figs. 1, 2.

Paratype: AMNH 183856 (1 specimen).

Locality: 55 m, on rock and coral, Tagus Cove, 00°16'38"S, 91°22'44"W, Albemarle Island, Galápagos Islands, Ecuador, Pacific Ocean.

Collector: AHF Station 147-34, Jan. 13, 1934.

Remarks: ex LACM-AHF. The holotype is LACM 1853 and additional paratypes are LACM 1854, CASIZ 016260, USNM 771232, ANSP A7213, and SDNHM 69305 (Ferreira, 1978a: 36).

Lepidozona allynsmithi Ferreira, 1974: 173–175, figs. 19–24.

Paratype: AMNH 181727 (3 valves).

Locality: 2–3 m, Caleta el Candelero, on the western side of Isla Espíritu Santo, 24°30'N, 110°23'W, Gulf of California, Mexico, Pacific Ocean.

Collector: A. J. Ferreira, Oct. 18, 1970.

Remarks: ex Ferreira. The holotype is CASIZ 016431 and additional paratypes are CASIZ 016432, 016433, LACM 1727, USNM 709833 and in the University of Costa Rica (Ferreira, 1974: 174–175).

Lepidozona guadalupensis Ferreira, 1978b: 37–38, figs. 13, 14, 32, 33.

Paratype: AMNH 183855 (1 specimen).

Locality: 29°01'N, 118°13'W, Sealer's Camp, Guadalupe Island, Mexico, Pacific Ocean.

Collector: A. J. Ferreira and W. L. Lee, Jan. 1–2, 1975.

Remarks: ex Ferreira. The holotype is CASIZ 016269 and additional paratypes are CASIZ 016270, 016271, LACM 1857, SDNHM 68728, USNM 770960, and ANSP A7226 (Ferreira, 1978b: 37–38).

Plaxiphora excravata Pilsbry, 1893: 327–328.

Syntypes: AMNH 21933 (3 specimens).

Locality: "Australia?"

Collector: Unknown.

Remarks: ex Haines Collection 2710i. Additional syntypes are in the McGill University Collection, Montreal, Canada (Pilsbry, 1893: 328). Two specimens are illustrated herein (fig. 1). Pilsbry, not Carpenter in Pilsbry, was the author of this species. This species is provisionally considered a synonym of *Plaxiphora (P.) albida* (Blainville, 1825) (Kaas and Van Belle, 1994: 294).

CLASS GASTROPODA

SUBCLASS OPISTHOBRANCHIA

Acteon eloiseae Abbott, 1973: 91–92, figs. 1–4.

Paratypes: AMNH 245763 (4 specimens).

Locality: Shallow water near shore, Al Masirah Island, Oman, Indian Ocean.

Collector: Fishermen, 1970.

Remarks: ex Bosch. The holotype is DMNH 27213 and additional paratypes are DMNH 67746 and in the collection of D. T. Bosch (Abbott, 1973: 92). This species is placed

in *Punctacteon* Kuroda and Habe, 1961 (Bosch et al., 1995: 179).

Alderia uda Ev. Marcus and E. Marcus, 1956: 8–14, figs. 8–23.

Holotype: AMNH 3485.

Locality: Near Santos, São Vicente, São Paulo, Brazil, Atlantic Ocean.

Collector: S. A. Gerlach, April 1954.

Remarks: ex Ev. Marcus and E. Marcus. The repository of the paratypes was not cited. This species is a synonym of *A. modesta* (Lovén, 1844) (Ev. Marcus, 1977: 7).

Antiopella mucloc E. Marcus, 1958a: 57.

Lectotype: AMNH 3508.

Locality: Among algae, northern base of the Oceanographic Institute of the University of São Paulo, approximately 180 km ENE Santos, near Ubatuba, Brazil, Atlantic Ocean.

Collector: Unknown, Sept. 4, 1956.

Remarks: ex E. Marcus. This specific name was intended to be first made available from Marcus' (1958b) description published Aug. 13, 1958, and based on a single specimen (AMNH 3508). However, this name was first published with accompanying descriptive notes in Marcus (1958a), received by USNM Aug. 13, 1958. Since the paper received on Aug. 13, 1958, must have been published prior to that date, the name must date from Marcus (1958a). The type series consists of the intended type specimen (not the holotype by monotypy) and the two additional specimens cited by Marcus (1958a). AMNH 3508 was the intended "holotype" and is here designated lectotype. The repository of the paralectotypes is unknown. This species is placed in *Janolus* Bergh, 1884 (Gosliner, 1981: 30).

Atys caribaea var. *corpulenta* Usticke, 1959: 84–85.

Syntype: AMNH 270412 (1 specimen).

Locality: "Altona Bay and Ham Bay," St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: Unknown.

Remarks: ex L. and L. Schwartz (Usticke Collection). This subspecific name was considered a nomen dubium for lack of type specimens (Faber, 1988: 75), but AMNH 270412 was located in 1994 in the portion of the Usticke collection in the possession

of L. and L. Schwartz. Usticke (1959: vi) stated that the figured specimen of each new taxon was the holotype. For this subspecies, however, no figure was given and thus all type specimens are syntypes. This taxon was cited as a probable synonym of *A. caribaea* (Orbigny, 1841) (Usticke MS: 259). However, the name *A. caribaea* as used by Usticke (in the sense of Pilsbry, 1895: 274, pl. 48, fig. 12a) refers to *A. riiseanus* Mörcz, 1875 (Ev. Marcus, 1970: 930).

Atys lineata Usticke, 1959: 85, pl. 4, fig. 15.

Holotype: AMNH 195475.

Paratype: AMNH 270411 (1 specimen).

Locality: Ham Bay, St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Usticke Collection (AMNH 195475) and L. and L. Schwartz (Usticke Collection) (AMNH 270411). AMNH 195475 was the figured specimen (Usticke, 1959). This species is considered a synonym of *A. caribaea* (Orbigny, 1841) by Abbott (1974: 320) but of *A. mandrewii* E. A. Smith, 1872 by Ev. Marcus (1977: 4).

Baeolidia benteva E. Marcus, 1958b: 65–68, figs. 105–111.

Holotype: AMNH 3512.

Locality: From the bottom of a boat, northern base of Oceanographic Institute of the University of São Paulo, approximately 180 km ENE of Santos, near Ubatuba, Brazil, Atlantic Ocean.

Collector: E. Nonato and C. Froelich, July 1956.

Remarks: ex E. Marcus. The repository of the paratype was not cited. This species is placed in *Berghia* Trinchese, 1877 (Ev. Marcus, 1977: 16).

Cadlina evelinae E. Marcus, 1958b: 18–21, figs. 28–33.

Holotype: AMNH 3506 (lost).

Locality: Under stones in the upper littoral, "Ilhabela and Guarujá," near Santos, Brazil, Atlantic Ocean.

Collector: Unknown, "December 1953, July 1954, July 1956."

Remarks: ex Marcus. This specimen was to have been deposited in AMNH (E. Marcus, 1958b: 1). It was lost in the mail and Mar-

cus submitted a voucher specimen (not from the localities cited), AMNH 226497. The repository of the paratypes was not cited. This species is placed in *Tyrinna* Bergh, 1898, by Rudman (1984: 245) but is retained in *Cadlina* by Rios (1994: 211).

Capellinia conicla E. Marcus, 1958b: 41–44, figs. 72–80.

Holotype: AMNH 3509.

Locality: Northern base of the Oceanographic Institute of the University of São Paulo, approximately 180 km ENE Santos, Ubatuba, Brazil, Atlantic Ocean.

Collector: Unknown, Oct. 1956.

Remarks: ex E. Marcus. The repository of the paratypes was not cited. This species is placed in *Eubranchus* Forbes, 1838, as *E. conicus* (Ev. Marcus, 1977: 14).

Catriona perca E. Marcus, 1958b: 45–48, figs. 81–87.

Holotype: AMNH 3513 (lost).

Locality: Lagoons, southern base of the Oceanographic Institute, University of São Paulo, approximately 200 km SW of Santos, Cananéia, Brazil, Atlantic Ocean.

Collector: K. Hosoe, Feb. 1953.

Remarks: ex E. Marcus. This specimen was never received at AMNH. The repository of the paratypes was not cited. This species is placed in *Cuthona* Alder and Hancock, 1855, by Gosliner (1980: 43) but is retained in *Catriona* by Rios (1994: 218).

Doto fragilis umia Ev. Marcus and E. Marcus, 1969: 27–29, figs. 38, 39.

Holotype: AMNH 146379.1, 146379.2.

Paratype: AMNH 226489 (1 specimen).

Locality: V-17-RD29, 60°27'N, 48°31'W, 326–366 m, SW Greenland, Atlantic Ocean.

Collector: R/V "Vema," Sept. 4, 1961.

Remarks: The body of the holotype is AMNH 146379.1 and the radula is AMNH 146379.2. This subspecies is a synonym of *D. chica* E. Marcus, 1960 (Ev. Marcus, 1977: 13).

Elysia vreelandae Ev. Marcus and E. Marcus, 1970: 194–195, figs. 36–40.

Holotype: AMNH 140586 [part].

Paratype: AMNH 140586A (1 specimen).

Locality: On *Codium*, San Agustin, Sonora, Mexico, Pacific Ocean.

Collector: P. Vreeland, Nov. 12, 1966.

Remarks: ex Ev. Marcus. The repository of additional paratypes as well as the remainder of the holotype was not cited.

Facelina coenda E. Marcus, 1958b: 52–55, figs. 88–96.

Holotype: AMNH 3510.

Paratype: AMNH 3511 (1 specimen).

Locality: Among algae in the upper littoral, northern base of the Oceanographic Institute of the University of São Paulo, approximately 180 km ENE Santos, Ubatuba, Brazil, Atlantic Ocean.

Collector: Unknown, Sept. 1956.

Remarks: ex E. Marcus. The repository of additional paratypes was not cited.

Hypselodoris regina Ev. Marcus and E. Marcus, 1970: 199, figs. 54–59.

Holotype: AMNH 140582.1–140582.3.

Locality: 2 m, on *Seriatopora angulata*, Tanikely, Madagascar, Indian Ocean.

Collector: E. Kirsteuer, Dec. 7, 1959.

Remarks: ex Ev. Marcus. This specimen is the holotype by monotypy. The body of the holotype is AMNH 140582.1 and two slides of internal structures are AMNH 140582.2, 140582.3.

Noumeaella isa Ev. Marcus and E. Marcus, 1970: 212, 215, figs. 86–89.

Holotype: AMNH 140584.1–140584.3.

Locality: 2 m, on *Seriatopora angulata*, Tanikely, Madagascar, Indian Ocean.

Collector: E. Kirsteuer, Dec. 6, 1959.

Remarks: ex Ev. Marcus. AMNH 140584.1–140584.3, mounted on three slides, is the holotype by monotypy. The AMNH number for this specimen was incorrectly given as 140854 in the original publication.

Philine thurmanni chilla Ev. Marcus and E. Marcus, 1969: 17–19, fig. 29.

Holotype: AMNH 146378.

Paratypes: AMNH 226486 (1 specimen), 226487 (1 specimen [part]), 226488 (1 specimen [part]).

Locality: V-17-3, 36°15'S, 76°51'W, 4030–4032 m (AMNH 146378, 226486); V-17-15, 47°02'S, 75°36'W, 626–642 m (AMNH 226487, 226488), Chile, Pacific Ocean.

Collector: R/V "Vema," March 14, 1961 (AMNH 146378, 226486) and March 24, 1961 (AMNH 226487, 226488).

Remarks: The repository of additional paratypes as well as the remainder of AMNH 226487 and 226488 was not cited.

Philine thurmanni thurmanni Ev. Marcus and E. Marcus, 1969: 14–17, figs. 23–28.

Holotype: AMNH 146377.

Paratypes: AMNH 226482 (1 specimen), 226483 (39 specimens), 226484 (1 specimen [part]), 226485 (1 specimen [part]).

Locality: V-17-97, 44°29'S, 60°59'W, 101 m (AMNH 146377, 226482); V-17-75, 41°41'S, 59°19'W, 82 m (AMNH 226483–226485), Argentina, Atlantic Ocean.

Collector: R/V "Vema," June 13 (AMNH 146377, 226482) and May 23 (AMNH 226483–226485), 1961.

Remarks: The repository of additional paratypes as well as the remainder of AMNH 226484 and 226485 was not cited.

Platydoris carolynae Mulliner and Sphon, 1974: 209–212, figs. 2–5.

Paratype: AMNH 173729 (1 specimen).

Locality: Station 27, 10 m, 00°25'S, 90°42'W, Jervis Island, Galápagos Islands, Ecuador, Pacific Ocean.

Collector: Ameripagos Expedition, March 24, 1971.

Remarks: ex Ameripagos Expedition. The holotype is CASIZ 019176 and additional paratypes are in the Charles Darwin Research Station, Galápagos Islands, LACM 1619, SDNBM 62826, 62827, USNM 735349 and DMNH 64524 (Mulliner and Sphon, 1974: 210).

Retusa sosa Ev. Marcus and E. Marcus, 1969: 6–8, fig. 7.

Holotype: AMNH 146376.1, 146376.2.

Locality: V-17-74, 41°27'S, 59°33'W, 71 m, Argentina, Atlantic Ocean.

Collector: R/V "Vema," May 23, 1961.

Remarks: The holotype is dissected and mounted on two slides. The repository of the paratype was not cited.

Rostanga byga E. Marcus, 1958b: 22–24, figs. 34–36.

Holotype: AMNH 3507.

Locality: Under a stone in the tidal zone, approximately 100 km ENE of Santos, Ilha Bela, Isla São Sebastião, Brazil, Atlantic Ocean.

Collector: Unknown, June 1956.

Remarks: ex E. Marcus. AMNH 3507 is the holotype by monotypy.

Smaragdinella kirsteueri Ev. Marcus and E. Marcus, 1970: 188–189, figs. 23–27.

Holotype: AMNH 140583.1–140583.3.

Paratype: AMNH 140583A (1 specimen).

Locality: 2 m, on *Seriatopora angulata* and *Millepora tenella*, Tanikely, Madagascar, Indian Ocean.

Collector: E. Kirsteuer, Dec. 7–8, 1959.

Remarks: ex Ev. Marcus. The body of the holotype is AMNH 140583.1 and the remainder is mounted on two slides (AMNH 140583.2, 140583.3).

Stiliger (Stiliger) erbsus Ev. Marcus and E. Marcus, 1970: 192–194, figs. 32–34.

Holotype: AMNH 140585.1, 140585.2.

Locality: 2.5 m, on *Acropora corymbosa*, Tanikely, Madagascar, Indian Ocean.

Collector: E. Kirsteuer, July 14, 1959.

Remarks: ex Ev. Marcus. AMNH 140585.1, 140585.2, mounted on two slides, is the holotype by monotypy.

Stiliger (Stiliger) talis Ev. Marcus and E. Marcus, 1956: 1–3, figs. 1–7.

Holotype: AMNH 3484.

Locality: Near Santos, São Vicente, São Paulo, Brazil, Atlantic Ocean.

Collector: S. A. Gerlach, April 1954.

Remarks: ex Ev. Marcus. The repository of the paratypes was not cited.

Potential opisthobranch type specimens that should be in the AMNH Collection but were not present or cataloged as of June 1995 (see Introduction for details).

Berthella tupala E. Marcus, 1957: 416–418, figs. 58–69 (syntype(s)?).

Cadlina rumia E. Marcus, 1955: 119–121, figs. 78–89 (syntype(s)?).

Catriona tina E. Marcus, 1957: 459–461, figs. 187–198 (syntype(s)?).

Chelidonura evelinae E. Marcus, 1955: 95–100, figs. 8–19 (syntype(s)?).

Discodoris pusae E. Marcus, 1955: 147–149, figs. 151–165 (syntype(s)?).

Doris bovena E. Marcus, 1955: 131–133, figs. 109–118 (syntype(s)?).

- Doto pita* E. Marcus, 1955: 169–170, figs. 223–225 (syntype(s)?).
- Doto uva* E. Marcus, 1955: 167–168, figs. 218–222 (holotype by monotypy?).
- Elysia canguzua* E. Marcus, 1955: 111–112, figs. 45–48, 60–65 (syntype(s)?).
- Elysia cauze* E. Marcus, 1957: 405–408, figs. 35–44 (syntype(s)?).
- Elysia serca* E. Marcus, 1955: 113–115, figs. 49–52, 59 (syntype(s)?).
- Favorinus auritulus* E. Marcus, 1955: 181–183, figs. 259–269 (syntype(s)?).
- Ganitus evelinae* E. Marcus, 1953: 173–175, figs. 24–40 (syntype(s)?).
- Glossodoris neona* E. Marcus, 1955: 124–126, figs. 90–101 (syntype(s)?).
- Hallaxa apefae* E. Marcus, 1957: 421–422, figs. 73–80 (holotype by monotypy?).
- Hancockia ryrca* E. Marcus, 1957: 451–454, figs. 164–174 (holotype by monotypy?).
- Janolus comis* E. Marcus, 1955: 170–172, figs. 226–236 (syntype(s)?).
- Okenia evelinae* E. Marcus, 1957: 438–440, figs. 128–137 (syntype(s)?).
- Phyllaplysia engeli* E. Marcus, 1955: 105–106, figs. 29–37 (syntype(s)?).
- Piseinotecus divae* E. Marcus, 1955: 177–178, figs. 244–248 (syntype(s)?).
- Pluscula cuica* E. Marcus, 1953: 177–181, figs. 41–49 (syntype(s)?).
- Polycera odhneri* E. Marcus, 1955: 158–160, figs. 193–202 (syntype(s)?).
- Polycerella conyna* E. Marcus, 1957: 429–432, figs. 104–110 (syntype(s)?).
- Siraius ilo* E. Marcus, 1955: 135–137, figs. 119–125 (holotype by monotypy?).
- Taringa telopia*, E. Marcus, 1955: 152–155, figs. 166–179 (syntype(s)?).
- Thordisa diuda* E. Marcus, 1955: 140–142, figs. 133–140 (holotype by monotypy?).
- Trapania maringa* E. Marcus, 1957: 442–443, figs. 138–145 (syntype(s)?).

CLASS BIVALVIA

- Aequipecten (Corymbichlamys) iredalei* Powell, 1958: 68–69, pl. 11, figs. 1, 2, text fig. A2.
- Paratypes: AMNH 105519 (5 valves).
- Locality: Station 675, 29°13.5'S, 177°57'W, 58–60 m, Raoul Island, Kermadec Islands, Pacific Ocean.
- Collector: R/V "Galathea," March 3, 1952.
- Remarks: ex Powell. The holotype is in the Zoological Museum, Copenhagen, Denmark (Powell, 1958: 66) and additional paratypes are FMNH 118997. This species is placed in the genus *Corymbichlamys* Iredale, 1939 (Rombouts, 1991: 36).
- Anodonta montezuma* Lea, 1841: 31.
- Holotype: AMNH 30871.
- Locality: Central America.
- Collector: Unknown.
- Remarks: ex Jay Collection 2057. AMNH 30871 is the holotype by monotypy and was figured by Lea (1842: 240–241, pl. 23, fig. 55). Although previously cited as lost by Johnson (1974: 92), this specimen was located at AMNH in 1994 and is fig. 2 herein. This taxon is a subspecies of *Anodontites* (A.) *trigonus* (Spix, 1827) (Haas, 1969: 561).
- Anodonta pavonia* Lea, 1838: 78–79, pl. 21, fig. 65.
- Paralectotypes: AMNH 30400 (6 specimens).
- Locality: "Little Beaver" [River, Lawrence Co.], Ohio, United States, North America.
- Collector: J. P. Kirtland.
- Remarks: ex Jay Collection 2064. Johnson (1974: 108) stated that the figured specimen (USNM 86514) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are USNM 86514a. This species is a synonym of *Strophitus undulatus* (Say, 1817) (Johnson, 1970: 367).
- Anodonta subcylindracea* Lea, 1838: 106, pl. 24, fig. 117.
- Syntypes: AMNH 31131 (1 valve) (lost), 31131a (1 specimen).
- Locality: Oak Orchard Creek, Orleans Co., New York, United States, North America.
- Collector: Unknown.

Remarks: ex Jay Collection 2073 (AMNH 31131) and ANSP 41045 (AMNH 31131a). The figured specimen could not be located in USNM or AMNH (Johnson, 1974: 139) and was not found at AMNH or USNM during the 1994–1995 search. AMNH 31131a (fig. 3 herein), is inscribed in Lea's hand with the species name and type locality and is close to, but does not appear to match, the figured specimen. We therefore consider AMNH 31131a a syntype. Neither specimen was cited by Johnson (1974: 139). This species is a synonym of *Anodontoides ferussacianus* (Lea, 1834) (Clarke, 1973: 92).

Anodontites irisans W. B. Marshall, 1926: 10–11, pl. 2, figs. 3, 5; pl. 3, fig. 7.

Paratypes: AMNH 35038 (2 specimens).

Locality: Venezuela, South America.

Collector: Unknown.

Remarks: These specimens were identified as paratypes by I. S. Oldroyd, who supplied Marshall with the material upon which the description was based. The holotype is USNM 359920 (Marshall, 1926: 11) and additional paratypes are BM(NH) 1926.2.3.29, 1926.2.3.30 (Johnson, 1971: 76). This species is a synonym of *A. (A.) patagonicus leotaudi* (Guppy, 1864) (Haas, 1969: 567).

Arca (Anadara) mazatlanica Hertlein and Strong, 1943: 156–157, pl. 1, figs. 1, 4.

Paratypes: AMNH 73931 (2 specimens).

Locality: Station 153-D-2, 23°6'N, 106°47'W, 19 mi west of Mazatlan, Mexico, Pacific Ocean.

Collector: Templeton Crocker Expedition, April 27, 1936.

Remarks: ex NYZS. The holotype is CASIZ 064296 and additional paratypes are CASIZ 064297 (Hertlein and Strong, 1943: 156). This species is placed in *Anadara* Gray, 1847, sensu stricto (Keen, 1971: 44).

Argina brevifrons var. *bucaruana* Sheldon and Maury in Maury, 1922: 200–201, pl. 2, fig. 16; pl. 3, fig. 10.

Syntypes: AMNH 27184 (3 valves).

Locality: Port of Tonosi, Bucaru, Los Santos Province, Panama, Pacific Ocean.

Collector: A. Olsson, 1918.

Remarks: ex Olsson via Maury. These specimens are labeled paratypes, but no holo-

type was designated in the original publication. The repository of additional syntypes was not cited. This taxon is a synonym of *Lunarca brevifrons* (Sowerby, 1833) (Keen, 1971: 50).

Asaphis nana Powell, 1958: 75, pl. 9, fig. 5, text fig. A6.

Paratype: AMNH 105514 (1 specimen).

Locality: Station 674, 29°15'S, 177°57'W, 75–85 m, Raoul Island, Kermadec Islands, Pacific Ocean.

Collector: R/V "Galathea," March 3, 1952.

Remarks: ex Powell. The holotype is in the Zoological Museum, Copenhagen, Denmark (Powell, 1958: 66) and additional paratypes are in the South Australian Museum (Willan, 1993: 90) and FMNH 119000. This species is provisionally placed in *Agnomyax* Stewart, 1930 (Willan, 1993: 90–91).

Aspatharia flava Pilsbry and Bequaert, 1927: 418–419, pl. 43, figs. 3, 3a; pl. 44, figs. 2, 2a.

Holotype: AMNH 72063.

Paratypes: AMNH 72064, 72190 (1 specimen each), 72191 (2 specimens).

Locality: 02°25'N, 27°30'E, Medje (AMNH 72063, 72064); 03°40'N, 29°40'E, Faradjé (AMNH 72190); 03°50'N, 30°10'E, Aba (AMNH 72191), Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: AMNH 72063 was pl. 43, figs. 3, 3a and AMNH 72064 was pl. 44, figs. 2, 2a (Pilsbry and Bequaert, 1927). Additional paratypes are MCZ 142360 (Johnson, 1956: 114) and ANSP 134205 (Johnson and Baker, 1973: 155), 132300, 132301, MRAC 57.395–57.399. This species is a synonym of *A. (A.) r. rugifera* (Dunker, 1858) (Haas, 1969: 588–589).

Basterotia pustula Usticke, 1971: 29–30, pl. 5, fig. 1618.

Syntypes: AMNH 198484 (2 valves).

Locality: St. Johns, Antigua and Barbuda, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1618. Usticke (1971: 2) stated that figured holotypes were segregated in his collection. Since neither of the figured unpaired valves in AMNH 198484 matches the given "ho-

lototype" dimensions, both valves must be syntypes.

Batissa gigantea Prime, 1860: 112.

Holotype: AMNH 31427.

Locality: Unknown.

Collector: United States Exploring Expedition to Japan under Commodore Perry.

Remarks: ex Jay Collection 10945+. AMNH 31427 is the holotype by monotypy. Although previously cited as missing by Johnson (1959: 453), this specimen was located at AMNH in 1994 (fig. 4 herein).

Bornia bibsae Usticke, 1969: 6, pl. 1, fig. 1616.

Lectotype: AMNH 198483 (1 valve).

Locality: South of Frederiksted, St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1616. AMNH 198483 was the figured specimen (Usticke, 1969) and is here designated lectotype.

Brachidontes exustus var. *rosaceus* Usticke, 1969: 5, pl. 1, fig. 1453.

Lectotype: AMNH 198479 (1 valve).

Locality: North of Frederiksted, St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1453. AMNH 198479 was the figured specimen (Usticke, 1969) and is here designated lectotype. Although the term "var." was employed, all "varieties" (including this one) described by Usticke (1969) were called "new s[ub] sp[ecies]" or, in error, "new sp[ecies]." This taxon, described with an infrasubspecific term after 1960, should nevertheless be considered subspecific as it was not published expressly as a "variety" in the traditional sense of the word. This taxon was erroneously cited as "*Brachidontes rosaceus*" in the figure caption. This subspecies is a synonym of *B. lavalleanus* Orbigny, 1842 (Usticke, 1971: 29) [= *B. exustus* (Linnaeus, 1758) (Rios, 1994: 236)].

Brachidontes pseudorecurvus Usticke, 1969: 5, pl. 1, fig. 1455.

Lectotype: AMNH 198480 (1 valve).

Locality: St. Thomas, U.S. Virgin Islands, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1455. AMNH 198480 was the figured specimen (Usticke,

1969) and is here designated lectotype. This species is a synonym of *Ischadium cubitus* (Say, 1821) (Usticke, 1971: 29).

Caelatura mesafricana Pilsbry and Bequaert, 1927: 402–403, pl. 32, figs. 2, 2a.

Holotype: AMNH 72093.

Paratype: AMNH 72202 (1 specimen).

Locality: 04°10'N, 29°40'E, Garamba River (AMNH 72093), 02°25'N, 27°30'E, Medje (AMNH 72202), Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, Feb. 1913 (AMNH 72093) and H. Lang, 1915 (AMNH 72202).

Remarks: Additional paratypes are ANSP 133850 (Johnson and Baker, 1973: 162). AMNH 72093 was the figured specimen and AMNH 72202 was the "shorter specimen" cited by Pilsbry and Bequeart (1927: 403). This species is the type of the genus *Mesafra* Haas, 1936, by original designation (Haas, 1936: 84).

Caelatura (Laevirostris?) rotula Pilsbry and Bequaert, 1927: 405–406, pl. 33, figs. 5, 5a.

Paratypes: AMNH 106840 (3 specimens).

Locality: 04°25'S, 15°20'E, Leopoldville, Congo River, Belgian Congo [Zaire], Africa.

Collector: J. Bequaert.

Remarks: ex MCZ. The holotype is ANSP 133847 (Johnson and Baker, 1973: 168) and additional paratypes are MCZ 161863 (Johnson, 1956: 133). *Coelatura* Conrad, 1853, was unjustifiably emended to *Caelatura* by Simpson (1900), making *Caelatura* a junior objective synonym (Rosenberg et al., 1990: 29). This species is a synonym of *Coelatura (Laevirostris) bourguignoni* (Rochebrune, 1886) (Haas, 1936: 66).

Caelatura stanleyvillensis Pilsbry and Bequaert, 1927: 403, pl. 32, figs. 3, 3a, 4; pl. 33, fig. 3.

Holotype: AMNH 72094.

Paratypes: AMNH 72095 (1 specimen), 72200 (20 specimens).

Locality: 00°30'N, 25°15'E, Stanleyville, Congo River, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: Additional paratypes are MCZ 142354 (Johnson, 1956: 136), ANSP

133852, FMNH 77363, and MRAC 56.360–56.362. AMNH 72094 was pl. 32, figs. 3, 3a and AMNH 72095 was pl. 32, fig. 4 (Pilsbry and Bequaert, 1927). This species is a synonym of *C. (Zairia) elegans* (Rochebrune, 1886) (Haas, 1936: 67). See remarks under the previous species for justification in correcting the spelling of the genus to *Coelatura*.

Calyptogena (Ectenagena) magnifica Boss and Turner, 1980: 165–177, figs. 1–9, 10 f, g, 11, 12 d–f, 13.

Paratype: AMNH 206087 (1 specimen).

Locality: Dive 984, 00°48.24'N, 86°13.47'W, 2450 m, Galápagos Rift, Pacific Ocean.

Collector: DSRV Alvin.

Remarks: ex MCZ 288829. The holotype is MCZ 288500 and additional paratypes are in MCZ as well as several other museum collections (Boss and Turner, 1980: 165).

Cardium (Laevicardium) clarionense Hertlein and Strong, 1947: 144–145, pl. 1, figs. 5–7, 14.

Paratype: AMNH 160330 (1 valve).

Locality: Station 136-D-15, 23°28'30"N, 109°25'W, 40 fathoms, in mud and crushed shell, Arena Bank, Gulf of California, Mexico, Pacific Ocean.

Collector: Templeton Crocker Expedition, April 20, 1936.

Remarks: ex NYZS. The holotype is CASIZ 065551 and additional paratypes are CASIZ 065552, 064508 (Hertlein and Strong, 1947: 145). This species is placed in *Laevicardium* Swainson, 1840, sensu stricto (Bernard, 1983: 38).

Cardium parvulum Li, 1930: 259, pl. 3, fig. 22.

Holotype: AMNH 268088 (1 valve).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22082. AMNH 268088 is the holotype by monotypy. This species is a synonym of *Tachycardium (Mexicardia) procerum* (Sowerby, 1833) (Bernard, 1983: 38).

Chione (Chione) guatulcoensis Hertlein and Strong, 1948: 182, pl. 1, figs. 2, 4, 6; pl. 2, figs. 1, 8, 12, 13.

Paratype: AMNH 160347 (1 specimen).

Locality: Station 195-D-9, 15°44'28", 96°07'51"W, 7 fathoms, in sand and crushed shell, Port Guatulco, Mexico, Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition, Dec. 5, 1937.

Remarks: ex NYZS. The holotype is CASIZ 065557 and additional paratypes are CASIZ 065558, 064492 (Hertlein and Strong, 1948: 182).

Chione minor Usticke, 1969: 6, pl. 1, fig. 1650.

Lectotype: AMNH 198485 (1 valve).

Locality: Mangrove swamps, "Antigua and Barbuda," Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1650. AMNH 198485 was the figured specimen (Usticke, 1969) and is here designated lectotype.

Chlamydella favus lemchei Powell, 1958: 70–71, pl. 9, figs. 7, 8.

Paratypes: AMNH 105518 (6 valves).

Locality: Station 674, 29°15'S, 177°57'W, 75–85 m, Raoul Island, Kermadec Islands, Pacific Ocean.

Collector: R/V "Galathea," March 3, 1952.

Remarks: ex Powell. The holotype is in the Zoological Museum, Copenhagen, Denmark (Powell, 1958: 71) and additional paratypes are FMNH 118999.

Condylonucula maya Moore, 1977: 124, figs. 4, 5.

Paratypes: AMNH 183858 (5 valves).

Locality: Chancanab Lagoon, 2 m, Cozumel, Quintana Roo, Mexico, Atlantic Ocean.

Collector: D. R. Moore, Nov. 29, 1971.

Remarks: ex Moore. The holotype is USNM 758533 and additional paratypes are USNM 758536, MCZ 288443, ANSP 344389, DMNH 120579, FMNH 198080, UMML 28-2810, and in the collection of the author (Moore, 1977: 124).

Corbula altirostris Li, 1930: 263–264, pl. 5, fig. 37.

Holotype: AMNH 268092 (1 valve).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22097. AMNH 268092 is the holotype by mono-

typy. This species is a synonym of *Mulinia pallida* (Broderip and Sowerby, 1829) (Keen, 1971: 207).

Corbula glypta Li, 1930: 264, pl. 5, figs. 38, 38a.

Holotype: AMNH 268094.

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22098. AMNH 268094 is the holotype by monotypy. This species is a synonym of *C. (Tenucorbula) tenuis* Sowerby, 1833 (Bernard, 1983: 58).

Crassatellites rufus Li, 1930: 257, pl. 3, fig. 16.

Holotype: AMNH 268086 (1 valve).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22076. AMNH 268086 is the holotype by monotypy. This species is a synonym of *Eucrassatella gibbosa* Sowerby, 1832 (Keen, 1971: 104).

Crassinella lata Usticke, 1969: 5, pl. 1, fig. 1545.

Lectotype: AMNH 198481 (1 valve).

Locality: Ham Bay, St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1545. AMNH 198481 was the figured specimen (Usticke, 1969) and is here designated lectotype.

Crassinella skoglundae Coan, 1979: 6–7, figs. 12–15.

Paratype: AMNH 200376 (1 specimen).

Locality: Bahia Banderas, 20°45'N, 105°24'W, 7.5–15 m, La Cruz, Nayarit, Mexico, Pacific Ocean.

Collector: C. and P. Skoglund, July 31, 1973.

Remarks: ex Skoglund. The holotype is CASIZ 085659 and additional paratypes are in CASIZ, LACM 1904, ANSP, MCZ 288512, and SDNHM (Coan, 1979: 7).

Cuspidaria (Tropidomya) nana Oldroyd, 1918: 28.

Paratype: AMNH 58306 (1 specimen).

Locality: In clay, Monterey Bay, California, United States, Pacific Ocean.

Collector: Unknown.

Remarks: ex Oldroyd. The holotype was in the Oldroyd Collection, Stanford University (Oldroyd, 1918: 28), now CASIZ 060981. The holotype was figured by Oldroyd (1924: 99, pl. 13, figs. 8, 9). This species is a synonym of *Sphenia luticola* (Valenciennes, 1846) (Bernard, 1983: 58).

Cyrena jayensis Lea, 1834: 108, pl. 17, fig. 52.

Syntype: AMNH 31337 (1 specimen).

Locality: Sumatra, [Indonesia].

Collector: Unknown.

Remarks: ex Jay Collection 760. Johnson (1959: 457) cited MCZ 176905 as a paratype. Since Lea did not designate a holotype and no lectotype has since been designated, all specimens in the type series are syntypes. AMNH 31337 is fig. 5 herein.

Cyrenoida rosea brevidentata Pilsbry and Bequaert, 1927: 376–377, text fig. 85.

Holotype: AMNH 72097.

Locality: 05°55'S, 12°35'E, Kunga, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: The paratype is ANSP 133310.

Donax dorotheae Morrison, 1971: 554–556, pl. 1, fig. Do.

Paratypes: AMNH 162251 (134 specimens), 181631 (33 specimens), 189618 (34 specimens), 213566 (30 specimens), 223498 (55 specimens), 225197 (580 specimens), 225198 (1820 specimens), 276090 (1370 specimens).

Locality: Grand Isle, Louisiana, United States, Atlantic Ocean.

Collector: J. P. E. Morrison, Jan. 3, 1970.

Remarks: ex USNM, MCZ and several private collections. The holotype is USNM 679773 and additional paratypes are in USNM and ANSP (Morrison, 1971: 555), SBMNH 32286, 35047 (Scott et al., 1990: 15), MCZ 302542, LACM 1604, and FMNH [uncataloged]. This species is a synonym of *D. texasanus* Philippi, 1847 (Adamkewicz and Harasewych, 1994: 24).

Donax striatus var. *intermedius* Usticke, 1969: 7, pl. 1, fig. 1769.

Lectotype: AMNH 198486 (1 valve).

Locality: Trinidad, Trinidad and Tobago, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1769. AMNH 198486 was the figured specimen (Usticke, 1969) and is here designated lectotype. Although the term "var." was employed, all "varieties" (including this one) described by Usticke (1969) were called "new s[ub] sp[ecies]" or, in error, "new sp[ecies]." This taxon, described with an infrasubspecific term after 1960, should nevertheless be considered subspecific as it was not published expressly as a "variety" in the traditional sense of the word. This taxon was erroneously cited as "*Donax intermedius*" in the figure caption.

Egeria nux Pilsbry and Bequaert, 1927: 367–368, pl. 29, figs. 4–7, text fig. 82.

Holotype: AMNH 72092 (lost).

Paratypes: AMNH 72180 (40 valves), 72181 (8 valves).

Locality: 0°S, 12°40'E, Malela, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: The holotype (Pilsbry and Bequaert, 1927: pl. 29, fig. 4) cannot be located in the AMNH collection. Additional paratypes are MCZ 181377, ANSP 133889, and MRAC 55.317–55.326. *Egeria* Roissy, 1805, is a synonym of *Galatea* Bruguiere, 1797 (Vaught, 1989: 133); therefore this species should be placed in *Galatea*.

Egeria tenuicula langi Pilsbry and Bequaert, 1927: 366–367, pl. 29, figs. 1–3.

Holotype: AMNH 72091.

Paratypes: AMNH 72185 (89 valves).

Locality: 0°S, 12°40'E, Malela, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: AMNH 72091 was pl. 29, fig. 1 (Pilsbry and Bequaert, 1927). Additional paratypes are MCZ 181379, ANSP 133886, 133888, and MRAC 55.331–55.350. See remarks under *Egeria nux* for justification in placing this species in *Galatea*.

Elliptio mcmichaeli Clench and Turner, 1956: 170–171, pl. 7, figs. 1, 2.

Paratypes: AMNH 271104 (6 specimens).

Locality: Choctawhatchee River, 8 mi west of Miller Cross Roads, Holmes Co., Florida, United States, North America.

Collector: W. J. Clench, R. D. Turner, and D. F. McMichael, Sept. 11, 1954

Remarks: ex MCZ. The holotype is MCZ 191922 and additional paratypes are in several museum collections (Clench and Turner, 1956: 170). This species is placed in the subgenus *Elliptio* Rafinesque, 1820 (Haas, 1969: 232). The spelling of "*E. mcmichaeli*" by Haas (1969: 232) was an unjustified emendation (Bieler, 1988: 423).

Ensis myrae Berry, 1953: 398–401, pl. 29, figs. 5, 6; text figs. 3, 4.

Paratype: AMNH 221880 (1 specimen).

Locality: Long Beach, California, United States, Pacific Ocean.

Collector: Unknown.

Remarks: ex Chace Collection via DuShane. The holotype is CASIZ 064460 and additional paratypes are in several museums and private collections (Berry, 1953: 399–400).

Ensis tropicalis Hertlein and Strong, 1955: 203, pl. 3, figs. 34, 35.

Holotype: AMNH 73421 (1 valve).

Paratypes: AMNH 73422 (4 valves), 73422A (30+ valves).

Locality: Station 7, South Passage, dredged from sandy bottom in 15 fathoms, Pearl Islands, Panama, Pacific Ocean.

Collector: "Askoy" Expedition, Feb. 13, 1941.

Remarks: Additional paratypes are CASIZ 032846 (Hertlein and Strong, 1955: 203) and FMNH 77534.

Eupera doellojuradoi Klappenbach, 1962: 102–106, figs. 1–3.

Paratypes: AMNH 138279 (4 specimens).

Locality: Rio de la Plata, Puerto Platero, Depto. Colonia, Uruguay, South America.

Collector: A. Figueiras, May 14, 1961.

Remarks: ex Collection Malacology, MNHN-M 1215. The holotype is MNHN-M 1214 and additional paratypes are MNHN-M 878, 989, 1215 (Klappenbach, 1962: 104–105) and MCZ 256925.

Eupera primei Klappenbach, 1967: 111–114, figs. 1–3.

Paratypes: AMNH 89158 (69 specimens).

Locality: Near Ucalayi River, Pequino, Peru, South America.

Collector: H. Bassler, Nov. 1923.

Remarks: ex Collection Malacology,

MNHN-M 1309. The holotype is MNHN-M 1220 and additional paratypes are MNHN-M 1309 (Klappenbach, 1967: 111–112).

Glauconome cumingi Prime, 1862b: 384–385, pl. 14, fig. 4.

Syntypes: AMNH 31550 (2 specimens).

Locality: Malacca, [Malaysia], Pacific Ocean.

Collector: Unknown.

Remarks: ex Jay Collection 10949 and 10950+. AMNH 31551, from the Constable Collection, was incorrectly cited by Johnson (1959: 459) as part of the type series; Constable had no type material for this species. Additional syntypes could not be located in either MCZ or BM(NH) (Johnson, 1959: 449).

Glauconome jayana Prime, 1861b: 354.

Lectotype: AMNH 31537.

Locality: Australia.

Collector: Unknown.

Remarks: ex Jay Collection 10950. No types could be located in MCZ and a lectotype was selected from the Jay Collection and figured by Johnson (1959: 457, pl. 5, fig. 5.)

Glycymeris gordoni form *tobagoensis* Usticke, 1959: 2–3.

Syntype: AMNH 271074 (1 valve).

Locality: South of Fredericksted, St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Usticke Collection via L. and L. Schwartz. Although a potentially infrasubspecific term ("form") was used in this pre-1960 description (see ICZN, 1985: 87—article 45g), this taxon should be considered subspecific because Usticke (1959) used the terms "form," "variety," and "subspecies" interchangeably and even mixed terms within a single species description. Usticke (1959: vi) designated "as holotypes the new species and sub-species that are pictured." This subspecies was not figured in the original publication and, therefore, no holotype was designated. AMNH 271074 is fig. 6 herein. This taxon was also cited as "*Glycymens* (sic) *gordoni* form *tobagoensis* Usticke (N. Sp.)" (Usticke, 1959: 2).

Iphigenia congo Pilsbry and Bequaert, 1927: 373–374, text fig. 83.

Paratypes: AMNH 72188 (34 valves).

Locality: 06°S, 12°40'E, Malela, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: The holotype is ANSP 133884a and additional paratypes are ANSP 133884, 134042, MCZ 181390, and MRAC 55.353–55.360.

Iridina coelestis Lea, 1838: 82–83, pl. 22, fig. 70.

Paralectotype: AMNH 31292 (1 specimen).

Locality: Africa.

Collector: Unknown.

Remarks: ex Jay Collection 1991. Johnson (1974: 32) indicated that all syntypes were lost. The specimen figured by Lea (1838) is USNM 86785, located during the 1994–1995 search, and is here designated lectotype. The Jay specimen was located at AMNH in 1994 and is figured for comparison (fig. 7 herein). An additional paralectotype is possibly USNM 86785a (there may be only two specimens in the type series or more, depending on how Lea's wording is interpreted). This species is a synonym of *Mutela* (*M.*) *dubia* (Gmelin, 1791) (Haas, 1969: 579).

Isocardia tetragona A. Adams and Reeve, 1850: 76–77, pl. 22, fig. 1.

Paralectotype: AMNH 51095 (1 specimen).

Locality: Japan, Pacific Ocean.

Collector: H. M. S. "Samarang," 1843–1846.

Remarks: ex Constable Collection. The label is marked "Adam's type specimen." Kosuge (1994: 42) stated that the holotype was BM(NH) 1874.12.11.270. Since A. Adams and Reeve did not designate a holotype, Kosuge's action constituted a lectotype designation. This specific name is preoccupied by *I. tetragona* Koch and Dunker, 1837, and was replaced with *Meocardia samarangiae* Bernard, Cai and Morton, 1993 (Bernard et al., 1993: 67).

Legrandina aucklandica Powell, 1933: 233, fig. 1.

Paratypes: AMNH 136332 (4 specimens).

Locality: Beach drift, Faith Harbour, Auckland Islands, New Zealand, Pacific Ocean.

Collector: J. Bollons.

Remarks: ex Mrs. Winton Barker. The holotype is in the M. K. Mestayer Collection and additional paratypes are in the M. K.

Mestayer Collection (Powell, 1933: 233) and LACM 1356 (Sphon, 1971: 2). This species is placed in *Perrierina* Bernard, 1897, as *P. (Legrandina) aucklandica* (Powell, 1979: 403).

Limopsis (Limopsis) diegensis Dall, 1908: 395, pl. 15, figs. 13, 15.

Syntypes: AMNH 226332 (25 specimens).

Locality: Station 2923, 32°40'30"N, 117°31'30"W, 822 fathoms, off San Diego, California, United States, Pacific Ocean.

Collector: "Albatross," Jan. 19, 1889.

Remarks: ex USNM. Additional syntypes are USNM 122585 (Dall, 1908: 395), LACM 1166 (Sphon, 1971: 2), USNM 110699, 110700, 111422, SBMNH 34060 (Scott et al., 1990: 10), MCZ 27844, 187523, and FMNH 41068, 90695.

Macoma (Psammacoma) panamensis spectri Hertlein and Strong, 1949a: 91-92, pl. 1, figs. 9, 10, 16.

Paratypes: AMNH 160341 (2 valves), 270833 (7 valves).

Locality: Station 143-D-3, 26°57'N, 111°56'W, 35 fathoms, in mud and crushed shell, Santa Inez Bay, Gulf of California, Mexico, Pacific Ocean (AMNH 160341) and no precise locality, probably Gulf of California, Mexico, Pacific Ocean (AMNH 270833).

Collector: Templeton Crocker Expedition, April 13, 1936 (AMNH 160341).

Remarks: ex NYZS. The holotype is CASIZ 031198 and additional paratypes are CASIZ 031199 (Hertlein and Strong, 1949a: 91). This subspecies is a synonym of *M. siliqua* (C. B. Adams, 1852) (Bernard, 1983: 45).

Mactra (Micromactra) fonsecana Hertlein and Strong, 1950: 232, pl. 2, figs. 16, 19, 20.

Paratypes: AMNH 160102, 160103 (1 valve each), 160104 (2 valves), 160342 (1 specimen).

Locality: Station 199-D-1, Meanguera Island, Gulf of Fonseca, El Salvador (AMNH 160102); beach, Corinto, Nicaragua (AMNH 160103); Station 197-D-2, 14 fathoms, mud, 7 mi west of Champerico, Guatemala (AMNH 160104); Potosi and Monypenny Point, Gulf of Fonseca, Nicaragua (AMNH 160342), Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition.

Remarks: ex NYZS. The holotype is CASIZ 061534 and additional paratypes are CASIZ 061535, 064511 (Hertlein and Strong, 1950: 232).

Mondocondylaea crebristriata Anthony, 1865: 205-206, pl. 18, fig. 1.

Paralectotypes: AMNH 271105 (3 specimens).

Locality: Pegu, Burma [Myanmar], Asia.

Collector: Unknown.

Remarks: ex MCZ. Johnson (1956: 111) stated that the holotype was MCZ 161872. Since Anthony did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are MCZ 161872 (Johnson, 1956: 111), 178935, ANSP 41761, 125804 (Johnson and Baker, 1973: 152), and USNM 86327. This species is a synonym of *Pseudodon i. inoscularis* (Gould, 1844) (Brandt, 1974: 267).

Monocondylaea mardinensis Lea, 1865: 286.

Paralectotype: AMNH 106843 (1 specimen).

Locality: Stream into Tigris River, near Mardin, Iraq, Asia.

Collector: Unknown.

Remarks: ex C. M. Wheatley via MCZ. Johnson and Baker (1973: 161) stated that the specimen (USNM 86354) figured by Lea (1869: 252-253, pl. 30, fig. 67) was the holotype. Since Lea did not designate a holotype, Johnson and Bakers' action constituted a lectotype designation. Additional paralectotypes are ANSP 41034 (Johnson and Baker, 1973: 161), MCZ 152886, 178756, 240335, 293262. This species is a synonym of *Leguminaia wheatleyi* (Lea, 1862) (Haas, 1969: 143).

Montacuta cruzensis Usticke, 1969: 6, pl. 1, fig. 1609.

Lectotype: AMNH 198482 (1 valve).

Locality: St. Croix, U.S. Virgin Islands, Atlantic Ocean.

Collector: G. Usticke.

Remarks: ex Usticke Collection 1609. AMNH 198482 was renumbered as 1611 by Usticke (1971: 29). AMNH 198482 was the figured specimen (Usticke, 1969) and is here designated lectotype.

Mutela garambae Pilsbry and Bequaert, 1927: 431-432, pl. 40, figs. 1, 1a, 2.

Holotype: AMNH 72067.

Locality: 04°10'N, 29°40'E, in sand and mud of the river bed, Middle Garamba River, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, Feb. 1913.

Remarks: AMNH 72067 was pl. 40, figs. 1, 1a (Pilsbry and Bequaert, 1927). Paratypes are ANSP 134051 (Johnson and Baker, 1973: 156) and MRAC 58.443. This species is a synonym of *M. (M.) dubia* (Gmelin, 1791) (Haas, 1969: 579–580).

Mutela garambae praetenuis Pilsbry and Bequaert, 1927: 432, pl. 40, fig. 3.

Holotype: AMNH 72066.

Locality: 03°50'N, 30°10'E, Near Aba, Aba River, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Chapin, 1909–1915.

Remarks: AMNH 72066 was the figured specimen (Pilsbry and Bequaert, 1927). A paratype is ANSP 132309 (Johnson and Baker, 1973: 167). This subspecies is a synonym of *M. (M.) dubia* (Gmelin, 1791) (Haas, 1969: 579–580).

Mutela langi Pilsbry and Bequaert, 1927: 432–433, pl. 39, figs. 3, 3a, 3b.

Paratypes: AMNH 72198 (62 valves).

Locality: 06°S, 12°50'E, in bed of an arm of the Congo River, Zambi, Belgian Congo [Zaire], Africa.

Collector: H. Lang and J. Bequaert, 1915.

Remarks: The holotype is ANSP 133860a (Johnson and Baker, 1973: 160) and additional paratypes are MCZ 41068 (Johnson, 1956: 121), ANSP 132311, 133860, UMMZ 239354 (Johnson, 1979: 32), FMNH 77355, and MRAC 58.471–58.496. This species is a synonym of *M. (M.) dubia* (Gmelin, 1791) (Haas, 1969: 579–580).

Mya japonica Jay, 1856: 292, pl. 1, figs. 7, 10.

Syntype: AMNH 34162 (1 specimen).

Locality: Volcano Bay, Yedo Island, Japan, Pacific Ocean.

Collector: U.S. Naval Expedition to Japan, 1852–1854.

Remarks: ex Jay Collection 10995. This specimen was figured by Habe (1977, pl. 1, fig. 10; pl. 2, figs. 1–3). This species is placed in the subgenus *Arenomya* Winckworth, 1930 (Higo and Goto, 1993: 664).

Mycetopus emarginatus Lea, 1860: 308.

Paralectotypes: AMNH 30459 (7 specimens).

Locality: Siam [Thailand], Asia.

Collector: Unknown.

Remarks: ex Haines Collection. Johnson (1974: 52) stated that the specimen (USNM 86787) figured by Lea (1863: 398–399, pl. 50, fig. 305) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is the type of the genus *Solenaria* Conrad, 1869, by monotypy as *S. emarginata* (Conrad, 1869: 249), but may be a species of the American genus *Mycetopoda* Orbigny, 1835 (Brandt, 1974: 334).

Nucula calcicola Moore, 1977: 120–122, figs. 1–3.

Paratypes: AMNH 183857 (10 valves).

Locality: Chancanab Lagoon, 2 m, Cozumel, Quintana Roo, Mexico, Atlantic Ocean.

Collector: D. R. Moore, Nov. 29, 1971.

Remarks: ex Moore. The holotype is USNM 758535 and additional paratypes are USNM 758537, DMNH 120581, ANSP 344387, MCZ 288445, BM(NH), in Leiden, MNHN-P, and UMML 28-2812 (Moore, 1977: 120–121).

Nucula eightsii Jay, 1839: 113, pl. 1, figs. 12, 13.

Syntype: AMNH 56119 (1 specimen).

Locality: New South Shetland Island, Atlantic Ocean.

Collector: J. G. Eights.

Remarks: ex Jay Collection 1447. Jay, not Couthouy in Jay, is the author of this species (Johnson, 1946: 39). Dell cited this specimen as "Jay's figured type." This statement does not imply that AMNH 56119 is the lectotype and Dell (1990) cited this specimen as the "figured syntype." Neither Dell (1963) nor Richards and Old (1969), who based their citation of AMNH 56119 as the "holotype" on Dell (1963), were therefore making a lectotype designation. This species is placed in *Yoldia* Möller, 1842, as *Y. (Aequiyoldia) eightsi* (Dell, 1990: 10).

Nucula (Nucula) interflucta Marinovich, 1973: 7–8, fig. 1.

Paratypes: AMNH 172412 (21 specimens).

Locality: 20°13'S, 70°10'W, Near end of Ave.

- Baquedana, Iquique, Tarapaca Province, Chile, Pacific Ocean.
- Collector: L. Marincovich, June–Aug. 1964.
- Remarks: ex LACM 64-16. The holotype is LACM 1581 and additional paratypes are LACM 1582, USNM 709015, CASIZ 085610, SDNHM, and ANSP (Marincovich, 1973: 7).
- Nucula (Nuculopsis) schencki* Hertlein and Strong, 1940: 384, pl. 1, figs. 8–10.
- Paratype: AMNH 160806 (1 specimen).
- Locality: Station 195-D-9, 15°44'28"N, 96°07'51"W, 7 fathoms, in sand and crushed shell, Port Guatulco, Mexico, Pacific Ocean.
- Collector: Eastern Pacific "Zaca" Expedition, Dec. 5, 1937.
- Remarks: ex NYZS. The holotype is CASIZ 065653 and additional paratypes are CASIZ 065654, 065655. This species is placed in the subgenus *Nucula* Lamarck, 1799 (Bernard, 1983: 9).
- Nuculana (Costelloleda) marella* Hertlein, Hanna, and Strong in Hertlein and Strong, 1940: 399–400, pl. 2, figs. 12, 13.
- Paratypes: AMNH 73925 (4 valves), 160807 (2 valves).
- Locality: Station 221-D-1-5, 07°54'15"-07°52'30"N, 82°02'45"-82°01'W, 35–40 fathoms, in sandy mud, Gulf of Chiriquí, Panama, Pacific Ocean.
- Collector: Eastern Pacific "Zaca" Expedition, March 13, 1938.
- Remarks: ex NYZS. The holotype is CASIZ 065656. This species is placed in the subgenus *Nuculana* Link, 1807 (Allen and Hannah, 1986: 232–233).
- Parabornia squillina* Boss, 1965: 2–6, figs. 1–3.
- Holotype: AMNH 112526.
- Paratypes: AMNH 112527 (8 specimens).
- Host: *Lysiosquilla scabricauda* (Malacostraca).
- Locality: Attached to the ventral surface of the abdomen and thorax of host, Cristobal, Canal Zone, Panama, Atlantic Ocean.
- Collector: Unknown, Aug. 23, 1934.
- Remarks: AMNH 112526 was fig. 1 (Boss, 1965). Additional paratypes are USNM 653701 (Boss, 1965: 4). This species is the type of the genus *Parabornia* Boss, 1965, by monotypy (Boss, 1965: 2).
- Pecten fililectus* Li, 1930: 255, pl. 2, fig. 10.
- Holotype: AMNH 268083 (1 valve).
- Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.
- Collector: D. F. MacDonald, 1907.
- Remarks: ex Columbia University 22070. AMNH 268083 is the holotype by monotypy. *P. filitextus* Pilsbry (1931: 429) was an unjustified emendation and therefore a junior synonym of *P. fililectus*. This species is a synonym of *Argopecten circularis* (Sowerby, 1835) (Keen, 1971: 87).
- Pecten islandicus pugetensis* Oldroyd, 1920: 136, pl. 4, figs. 5, 6.
- Paratype: AMNH 30549 (1 specimen).
- Locality: Off San Juan Islands, Puget Sound, Washington, United States, Pacific Ocean.
- Collector: Unknown.
- Remarks: ex Oldroyd. The holotype was in the Oldroyd Collection at Stanford University (Oldroyd, 1920: 136), now CASIZ 064278. Additional paratypes are CASIZ 066375 and SBMNH 34893 (Scott et al., 1990: 12). This subspecies is a synonym of *Chlamys hastata* (Sowerby, 1843) (Bernard, 1983: 24).
- Pecten latiauritus* [sic] *indentus* Li, 1930: 256, pl. 2, fig. 13.
- Lectotype: AMNH 268085 (1 valve).
- Paralectotype: AMNH 268085a (1 valve).
- Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.
- Collector: D. F. MacDonald, 1907.
- Remarks: ex Columbia University 22073. AMNH 268085 was the figured specimen (Li, 1930) and is here designated lectotype. This subspecies is a synonym of *Leptopecten (Pacipecten) tumbezensis* (Orbigny, 1846) (Bernard, 1983: 25).
- Pecten latiauritus* [sic] *splendens* Li, 1930: 256, pl. 2, fig. 12.
- Holotype: AMNH 268084 (1 valve).
- Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.
- Collector: D. F. MacDonald, 1907.
- Remarks: ex Columbia University 22072. AMNH 268084 is the holotype by monotypy. This subspecies is a synonym of *Lep-*

topecten (Pacipecten) tumbezensis (Orbigny, 1846) (Bernard, 1983: 25).

Pecten (Leptopecten) velero bolleyi Hertlein and Strong, 1946: 60–61, pl. 1, fig. 6.

Paratypes: AMNH 160346 (3 valves).

Locality: Station 203-D-1, 10°56'05"N, 85°49'25"W, 15 fathoms, Port Parker, Costa Rica, Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition, Jan. 20, 1938.

Remarks: ex NYZS. The holotype is CASIZ 064334 and additional paratypes are CASIZ 064335 (Hertlein and Strong, 1946: 61). This subspecies is accorded species status in *Leptopecten* Verrill, 1897 sensu stricto (Bernard, 1983: 25).

Penitella hopkinsi Kennedy and Armentrout, 1989: 320–324, figs. 1–13.

Paratype: AMNH 232083 (1 specimen).

Locality: Southeast margin of modern emergent marine abrasion platform on southwest facing end of Wingham Island, 59°59'28.2"N, 144°22'36"W, Gulf of Alaska, Alaska, United States, Pacific Ocean.

Collector: J. M. Armentrout, July 16, 1975.

Remarks: ex Kennedy. The holotype is LACM 2387 and additional paratypes are in several museum collections and in the collection of the junior author (Kennedy and Armentrout, 1989: 321, 323).

Penitella richardsoni Kennedy, 1989: 316–318, figs. 7–15.

Paratypes: AMNH 104306 (2 specimens).

Locality: At minus 2.3 tide, in soft clay rocks, 5 mi east of Port Orchard, Waterman, Washington, United States, Pacific Ocean.

Collector: E. Wingard, May 30, 1961.

Remarks: ex Kennedy. The holotype is LACM 2378 and additional paratypes are in several museum collections (Kennedy, 1989: 318).

Phacoides (Lucinisca) liana Pilsbry, 1931: 435, pl. 41, fig. 3.

Holotype: AMNH 268087 (1 valve) (lost).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22080. This specimen could not be located at AMNH, ANSP, MCZ or USNM. This spe-

cies is a synonym of *Lucina centrifuga* (Dall, 1901) (Keen, 1971: 121).

Pisidium fallax Sterki, 1896: 20.

Syntypes: AMNH 64869 (9 valves).

Locality: Tuscarawas River, on sand and gravel, New Philadelphia, Ohio, United States, North America.

Collector: V. Sterki, Oct.–Nov. 1891.

Remarks: ex Sterki. Additional syntypes are MCZ 88389 (Johnson, 1959: 451). The species was figured by Herrington (1962, pl. 4, fig. 5; pl. 7, fig. 11).

Pitar (Hyphantosoma) aletes Hertlein and Strong, 1948: 172–173, pl. 1, figs. 9, 11–13.

Paratype: AMNH 160329 (1 valve).

Locality: Gulf of California, Mexico, Pacific Ocean.

Collector: Templeton Crocker Expedition, 1936.

Remarks: ex NYZS. The holotype is CASIZ 065554 (Hertlein and Strong, 1948: 172).

Pitar (Lamelliconcha) frizzelli Hertlein and Strong, 1948: 176, pl. 1, figs. 1, 5, 7; pl. 2, fig. 11.

Paratypes: AMNH 160333 (2 valves).

Locality: Station 136-D-26, 23°27'N, 109°24'W, 45 fathoms, in sand and crushed shell, Arena Bank, Gulf of California, Mexico, Pacific Ocean.

Collector: Templeton Crocker Expedition, May 1, 1936.

Remarks: ex NYZS. The holotype is CASIZ 065555 and additional paratypes are CASIZ 065556, 064491.

Raeta maxima Li, 1930: 263, pl. 5, fig. 35.

Lectotype: AMNH 268093 (1 valve).

Paralectotype: AMNH 268093a (1 valve).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22095. AMNH 268093 was the figured specimen (Li, 1930) and is here designated lectotype. This species is a synonym of *Harvella elegans* (Sowerby, 1825) (Keen, 1971: 206).

Salaputium iredalei Powell, 1958: 72–73, pl. 9, fig. 6.

Paratypes: AMNH 105515 (10 valves).

Locality: Station 674, 29°15'S, 177°57'W, 75–85 m, Raoul Island, Kermadec Islands, Pacific Ocean.

Collector: R/V "Galathea," March 3, 1952.

Remarks: ex Powell. The holotype is in the Zoological Museum, Copenhagen, Denmark (Powell, 1958: 66) and additional paratypes are FMNH 118975.

Semele aurora Tursch and Pierret, 1964: 35–36, figs. 1–3.

Paratype: AMNH 111915 (1 specimen).

Locality: Off Rio de Janeiro, 30 fathoms, in sand, Brazil, Atlantic Ocean.

Collector: B. Tursch.

Remarks: ex Tursch. The holotype is CASIZ 064686 and an additional paratype is Museu Nacional, Rio de Janeiro 3116 (Tursch and Pierret, 1964: 35–36). This species is a synonym of *Semele modesta* (Reeve, 1853) (Rios, 1994: 275).

Semele craneana Hertlein and Strong, 1949b: 241–242, pl. 1, figs. 19, 22.

Paratype: AMNH 160337 (1 valve).

Locality: Station 136-D-26, 23°27'N, 109°24'W, 45 fathoms, in sand and crushed shell, Arena Bank, Gulf of California, Mexico, Pacific Ocean.

Collector: Templeton Crocker Expedition, May 1, 1936.

Remarks: ex NYZS. The holotype is CASIZ 064521 and additional paratypes are CASIZ 064534, 064497 (Hertlein and Strong, 1949b: 241). This species is placed in the subgenus *Amphidesma* Lamarck, 1818 (Coan, 1988: 16).

Semele jamesi Coan, 1988: 33–35, figs. 62, 63.

Paratypes: AMNH 232169 (3 valves).

Locality: 05°33'N, 87°02'30"W, 46–69 m, in coral rubble, Chatham Bay, Cocos Island, Costa Rica, Pacific Ocean.

Collector: D. Shasky, M. Montoya and K. Kaiser, May 27, 1985.

Remarks: ex SBMNH 34942. The holotype is SBMNH 34941 and additional paratypes are in several museum collections (Coan, 1988: 33).

Septifer zeteki Hertlein and Strong, 1946: 71–72, pl. 1, figs. 1, 2.

Paratype: AMNH 160073 (1 specimen).

Locality: Taboga Island, 25 fathoms, Panama, Pacific Ocean.

Collector: J. Zetek, 1937–1938.

Remarks: ex NYZS. The holotype is CASIZ 064328 and additional paratypes are CASIZ 064329–064331 (Hertlein and Strong, 1946: 71).

Solen crockeri Hertlein and Strong, 1950: 225–226, pl. 1, figs. 3, 5, 7.

Paratype: AMNH 160343 (1 specimen).

Locality: Station 199-D-2, 13°02'30"N, 87°29'30"W, 5 fathoms, Monypenny Point, Gulf of Fonseca, Nicaragua, Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition, Dec. 24, 1937.

Remarks: ex NYZS. The holotype is CASIZ 039970 and additional paratypes are CASIZ 064509, 061533 (Hertlein and Strong, 1950: 225). This species is placed in the subgenus *Solen* Linnaeus, 1758 (Bernard, 1983: 41).

Solen viridis var. *aldrigei* Usticke, 1969: 7, pl. 1, fig. 1780.

Lectotype: AMNH 198487 (1 valve).

Locality: Ramsey Bay, Antigua and Barbuda, Atlantic Ocean.

Collector: R. Aldridge.

Remarks: ex Usticke Collection 1780. AMNH 198487 was the figured specimen (Usticke, 1969) and is here designated lectotype. Although the term "var." was employed, all "varieties" (including this one) described by Usticke (1969) were called "new s[ub] sp[ecies]" or, in error, "new sp[ecies]." This taxon, described with an infrasubspecific term after 1960, should nevertheless be considered subspecific as it was not published expressly as a "variety" in the traditional sense of the word. This taxon was erroneously cited as "*Solen aldrigei*" in the figure caption.

Somalipecten cranmerorum Waller, 1986: 42–45, figs. 1–13.

Paratypes: AMNH 222557 (1 specimen), 222558 (6 specimens), 241626 (1 specimen).

Locality: Trawled from 150–300 m, off Somalia, Indian Ocean.

Collector: Taiwanese fisherman, 1985.

Remarks: ex USNM (AMNH 222557), D. Dan (AMNH 222558) and M. C. Hayes Collection (AMNH 241626). The holotype

is USNM 859034 and additional paratypes are USNM 859035, 859036, and in several museum and private collections (Waller, 1986: 45). This species is the type of the genus *Somalipecten* Waller, 1986, by original designation (Waller, 1986: 41).

Sphaerium hendersoni Sterki, 1906: 69–70.
Syntypes: AMNH 172860 (3 specimens).
Locality: Waterholes, Crow Creek, 25 mi northeast of Greeley, Weld Co., Colorado, United States, North America.

Collector: J. Henderson.

Remarks: ex University of Colorado via W. Weeks. Additional syntypes are MCZ 10405 (Johnson, 1959: 455), SBMNH 35050 (Scott et al., 1990: 15), MCZ 36569, and FMNH 59070, 59527, 69443. This species is a synonym of *S. striatinum* (Lamarck, 1818) (Herrington, 1962: 27).

Tellina (Moerella) arenica Hertlein and Strong, 1949a: 68–69, pl. 1, figs. 5, 11.

Paratypes: AMNH 160339 (1 specimen), 160344 (1 valve).

Locality: Between Puntarenas and Bat Bat Island, Puntarenas, Costa Rica (AMNH 160339) and Station 136-D-20, 23°30'N, 109°26'W, 43 fathoms, Arena Bank, Gulf of California, Mexico (AMNH 160344), Pacific Ocean.

Collector: Templeton Crocker Expedition, 1932 (AMNH 160339) and April 30, 1936 (AMNH 160344).

Remarks: ex NYZS. The holotype is CASIZ 031181 and additional paratypes are CASIZ 031182 (Hertlein and Strong, 1949a: 68). This species is a synonym of *T. (Angulus) carpenteri* Dall, 1900 (Bernard, 1983: 42).

Tellina (Eurytellina) eburnea askoyana Hertlein and Strong, 1955: 197, pl. 3, figs. 3, 13–15, 20, 21, 23.

Paratype: AMNH 73432 (1 specimen).

Locality: Station 19, sample 35, 14–33 m, gray sandy mud bottom, Pinas Bay, Panama, Pacific Ocean.

Collector: "Askoy" Expedition, Feb. 24, 1941.

Remarks: The holotype is CASIZ 031183 and additional paratypes are CASIZ 032842, 032843 (Hertlein and Strong, 1955: 197). AMNH 73432 was pl. 3, fig. 3 (Hertlein and Strong, 1955). This subspecies is a syn-

onym of *T. (E.) eburnea* Hanley, 1844 (Bernard, 1983: 43).

Tellina (Scissula) nicoyana Hertlein and Strong, 1949a: 85–86, pl. 1, figs. 23–26.

Paratypes: AMNH 160092 (6 valves), 160336 (1 specimen).

Locality: Station 213-D-11, 09°44'52"N, 84°51'25"W to 09°42'N, 84°56'W, 35 fathoms, in mud, Ballena Bay, Gulf of Nicoya, Costa Rica, Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition, Feb. 25, 1938.

Remarks: ex NYZS. The holotype is CASIZ 031190 and additional paratypes are CASIZ 031191, 064494 (Hertlein and Strong, 1949a: 85). This species is the type of the subgenus *Hertellina* Olsson, 1961 by monotypy (Olsson, 1961: 409).

Tellina panamanensis Li, 1930: 262, pl. 5, fig. 32.

Lectotype: AMNH 268089 (1 valve).

Paralectotype: AMNH 268090 (1 valve).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22092. AMNH 268089 (Li's figured specimen) was designated lectotype by Pilsbry (1931: 436, pl. 41, figs. 5, 6). This species is a synonym of *T. (Eurytellina) eburnea* Hanley, 1844 (Keen, 1971: 215).

Tellina tenuilineatus Li, 1930: 262, pl. 5, fig. 33.

Holotype: AMNH 268091 (1 valve).

Locality: Mouth of Rio Grande, in mud, 10–40 ft, 1 mi from mainland near La Boca, Panama Bay, Panama, Pacific Ocean.

Collector: D. F. MacDonald, 1907.

Remarks: ex Columbia University 22093. AMNH 268091 is the holotype by monotypy. This species is a synonym of *Temoconcha cognata* (C. B. Adams, 1852) (Keen, 1971: 234).

Tellina (Tellinella) zacae Hertlein and Strong, 1949a: 65–66, pl. 1, figs. 12, 13, 17.

Paratype: AMNH 160334 (1 specimen).

Locality: Station 136-D-32, 23°24'30"N, 109°24'24"W to 23°28'N, 109°23'30"W, 42 fathoms, Arena Bank, Gulf of California, Mexico, Pacific Ocean.

Collector: Templeton Crocker Expedition, May 2, 1936.

Remarks: ex NYZS. The holotype is CASIZ 031200 and additional paratypes are CASIZ 064495 (Hertlein and Strong, 1949a: 65).

Transennella gerrardi Abbott, 1958: 130–131, pl. 4, figs. a–c, text fig. 7.

Paratypes: AMNH 108391 (5 valves).

Locality: In sand and broken coral, 5 fathoms, Georgetown Anchorage, Grand Cayman Island, Cayman Islands, Atlantic Ocean.

Collector: A. J. Ostheimer III, Oct. 28, 1954.

Remarks: ex ANSP. The holotype is ANSP 199507 and additional paratypes are ANSP 199506, 199508–199510, USNM 619551 (Abbott, 1958: 131), MCZ 211610 and MNHN-M (Olazarri et al., 1972: 4).

Unio africanus Lea, 1856: 94.

Paratypes: AMNH 30820 (3 specimens). Locality: Cape of Good Hope, South Africa, Africa.

Collector: E. Verreaux.

Remarks: ex Haines Collection 1481. Johnson (1974: 11) stated that the specimen (USNM 85233) figured by Lea (1858: 300–301, pl. 27, fig. 15) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paratypes are USNM 85233a. This species is a synonym of *Cafferia caffra* (Krauss, 1848) (Haas, 1969: 155).

Unio boydianus Lea, 1840: 286–287.

Paratype: AMNH 61912 (1 specimen).

Locality: Oak Orchard Creek, Orleans Co., New York, United States, North America.

Collector: Dr. Boyd.

Remarks: ex Jay Collection 1515. Johnson (1974: 22) stated that the specimen (USNM 86126) figured by Lea (1842: 216, pl. 16, fig. 32) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is a synonym of *Lampsilis (L.) radiata* (Gmelin, 1791) (Burch, 1975: 160).

Unio burtoni Woodward, 1859: 349, pl. 47, fig. 1.

Paratype: AMNH 56105 (1 specimen). Locality: Lake Tanganyika, Kenya, Africa.

Collector: Captain Speke.

Remarks: This specimen bears an original (unknown) collection number 11955. Johnson (1971: 80) stated that the figured specimen (BM(NH) 1859.12.23.9) was the holotype. Since Woodward did not designate a holotype, Johnson's action constituted a lectotype designation. An additional paratype is BM(NH) 1965220 (Johnson, 1971: 80). This species is placed in *Coelatura* Conrad, 1853, and is the type of the subgenus *Grandidieria* Bourguignat, 1885, by original designation (Bourguignat, 1885: 6).

Unio callosus Lea, 1841: 31.

Holotype: AMNH 30561.

Locality: Ohio Canal, 12 mi below Columbus, Franklin Co., Ohio, North America.

Collector: Unknown.

Remarks: ex Jay Collection 1524. AMNH 30561 is the holotype by monotypy and was figured by Lea (1842: 239–240, pl. 23, fig. 54). Johnson (1974: 26) stated that this specimen was lost, but it was located at AMNH in 1994 and is fig. 8 herein. This species is a possible senior synonym of *Nephronaias (Sphenonaias) mexicana* (Philippi, 1847) (Haas, 1969: 206).

Unio cucumoides Lea, 1840: 285.

Holotype: AMNH 30897.

Locality: Hunter's River, New South Wales, Australia.

Collector: Unknown.

Remarks: ex Jay Collection 1608. AMNH 30897 is the holotype by monotypy and was figured by Lea (1842: 192–193, pl. 7, fig. 2) and is fig. 9 herein. This species is a synonym of *Cucumerunio novaehollandiae* (Gray, 1834) (Smith, 1992: 17).

Unio eximius Lea, 1856: 93.

Paratypes: AMNH 30785 (4 specimens, 1 valve).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1642. Johnson (1974: 54) stated that the specimen (USNM 83900) figured by Lea (1858: 294–295, pl. 25, fig. 8) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paratypes are

USNM 83900a and FMNH 77358. This species is placed in *Physunio* Simpson, 1900, sensu stricto (Haas, 1969: 89).

Unio gravidus Lea, 1856: 93.

Paralectotypes: AMNH 30772 (9 specimens, 1 valve), 31281 (2 specimens, 1 valve).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1672. Johnson (1974: 67) stated that the specimen (USNM 83935) figured by Lea (1858: 292–293, pl. 24, fig. 5) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are USNM 83935a, MCZ 221746, and ANSP 41748 (Johnson and Baker, 1973: 157). This species is the type of the genus *Physunio* Simpson, 1900, by original designation (Simpson, 1900: 830), but is a junior synonym of *P. (P.) superbus* (Lea, 1843) (Haas, 1969: 90).

Unio hainesianus Lea, 1856: 92.

Lectotype: AMNH 31070.

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection. Lea (1858: 289–290, pl. 21, fig. 1) stated "there are two specimens only of this fine species among the shells from Mr. Haines." This contradicts Simpson's (1900: 582) claim that "the type consists of a single valve of a young individual." AMNH 31070 is the figured specimen that Johnson (1974: 67) stated could not be found in USNM or AMNH; it was located at AMNH in 1994. It is here designated lectotype and is fig. 10 herein. USNM 83933, cited as a syntype by Johnson (1974: 67), is therefore the paralectotype. This species is the type of the genus *Chamberlainia* Simpson, 1900, by original designation as *C. hainesiana* (Simpson, 1900: 582).

Unio housei Lea, 1856: 92.

Paralectotypes: AMNH 29223 (3 specimens), 31039 (5 specimens).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1684. Johnson (1974: 71) stated that the specimen (USNM 83951) figured by Lea (1858: 291,

pl. 23, fig. 3) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. An additional paralectotype is MCZ 221846. This species is a synonym of *Hyriopsis (Limnoscapha) myersiana* (Lea, 1856) (Brandt, 1974: 275).

Unio humilis Lea, 1856: 93.

Paralectotypes: AMNH 30825 (1 specimen, 4 valves).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1891. Johnson (1974: 72) stated that the specimen (USNM 84153) figured by Lea (1858: 296–297, pl. 26, fig. 10) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are USNM 84153a. This species is placed in *Indonaia* Prashad, 1918 (Brandt, 1974: 286).

Unio inornatus Lea, 1856: 93.

Paralectotypes: AMNH 30813 (6 specimens), 30837 (2 specimens).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1688. Johnson (1974: 74) stated that the specimen (USNM 85196) figured by Lea (1858: 293–294, pl. 24, fig. 6) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is a synonym of *Uniantra contradens tumidula* (Lea, 1856) (Brandt, 1974: 290).

Unio jayensis Lea, 1838: 28–29, pl. 9, fig. 23.

Paralectotype: AMNH 30563 (1 valve).

Locality: Florida, United States, North America.

Collector: Unknown.

Remarks: ex Jay Collection 1693. Johnson (1974: 77) stated that the valve figured by Lea (USNM 86031) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. USNM 86031a (1 specimen) cannot be a paralectotype since Lea (1838) stated that he had only "single opposed valves of two individuals of nearly the same size." This species is placed in *Elliptio* Raf-

inesque, 1820, sensu stricto (Haas, 1969: 220–221).

Unio lutens Lea, 1856: 93.

Holotype: AMNH 29109.

Locality: Newville, Burma [Myanmar], Asia.
Collector: Mrs. Vinton.

Remarks: ex Haines Collection 1727. AMNH 29109 is the holotype by monotypy and was figured by Lea (1858: 291–292, pl. 24, fig. 4). Lea (1858: 291) corrected this specific name to *U. luteus*, as he considered the original spelling to be incorrect. This is an unjustified emendation because there is no evidence in the original publication that *U. lutens* was a typographical error. Johnson (1974: 87) stated that this specimen was lost, but it was located at AMNH in 1994 and is fig. 11 herein. This species is a synonym of *Parreysia corrugata* (Müller, 1774) (Haas, 1969: 118).

Unio lutulentus Gould, 1850: 295.

Paralectotypes: AMNH 30917 (3 specimens).

Locality: New Zealand.

Collector: United States Exploring Expedition, 1838–1842.

Remarks: ex Crooke Collection 17490. These specimens are types (W. J. Clench, on specimen label). The species was figured by Gould (1852: 428, pl. 37, figs. 542, 542a–c). The lectotype is USNM 5927 (McMichael, 1958: 428) and additional paralectotypes are USNM 17399 and MCZ 88092, 155765 (McMichael and Hiscock, 1958: 462). This species is a synonym of *Hyridella (Echyridella) menziesi* (Gray, 1843) (McMichael and Hiscock, 1958: 460).

Unio myersianus Lea, 1856: 92.

Paralectotypes: AMNH 29224 (2 specimens), 31007 (1 specimen), 31038 (6 specimens).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1745. Johnson (1974: 95) stated that the specimen (USNM 83950) figured by Lea (1858: 290, pl. 22, fig. 2) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is placed in *Hyriopsis* Conrad, 1853, as *H. (Limnoscapha) myersiana* (Brandt, 1974: 275).

Unio novi-eboraci Lea, 1838: 104–105, pl. 24, fig. 114.

Paralectotypes: AMNH 29006 (2 specimens, 1 valve).

Locality: Oak Orchard Creek, Orleans Co., New York, United States, North America.
Collector: Unknown.

Remarks: ex Jay Collection 1762. Johnson (1974: 100) stated that the figured specimen (USNM 86042) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is a synonym of *Villosa iris* (Lea, 1829) (Oesch, 1984: 201).

Unio nucleus Lea, 1856: 94.

Paralectotype: AMNH 30840 (1 specimen).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1762. Johnson (1974: 100) stated that the specimen (USNM 533741) figured by Lea (1858: 306–307, pl. 28, fig. 21) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is placed in *Scabies* Haas, 1911 (Haas, 1911: 105).

Unio occatus Lea, 1860: 307–308.

Paralectotype: AMNH 31251 (1 specimen).

Locality: Bengal, India, Asia.

Collector: Unknown.

Remarks: ex Crooke Collection 17510. AMNH 31251 is a type (W. J. Clench, on specimen label). Johnson (1974: 103) stated that the specimen (USNM 84077) figured by Lea (1863: 398, pl. 50, fig. 304) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. An additional paralectotype is USNM 84077a. This species is placed in *Indonaia* Prashad, 1918, as *I. occata* (Haas, 1969: 114).

Unio phaselus Lea, 1856: 94.

Paralectotypes: AMNH 30827 (4 specimens, 1 valve).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1807. Johnson (1974: 112) stated that the specimen (USNM 84143) figured by Lea (1858: 297–298, pl. 26, fig. 11) was the holotype. Since Lea did not designate a holotype, John-

son's action constituted a lectotype designation. This species is placed in *Scabies* Haas, 1911 (Brandt, 1974: 283).

Unio rusticus Lea, 1856: 93.

Paralectotypes: AMNH 30810 (12 specimens).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1864. Johnson (1974: 127) stated that the specimen (USNM 84082) figured by Lea (1858: 294, pl. 25, fig. 7) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are ANSP 41681 (1 specimen) (Johnson and Baker, 1973: 169), MCZ 221747 (1 valve), and USNM 84082a (2 specimens); a total of 15 1/2 paralectotypes. Lea's (1858) statement that "about a dozen of this species" were examined by him is vague enough to permit the recognition of this number of specimens in the type series. This taxon is a subspecies of *Uniandra contradens* (Lea, 1838) as *U. c. rustica* (Brandt, 1974: 291).

Unio sagittarius Lea, 1856: 93.

Paralectotype: AMNH 30809 (1 specimen).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection. Johnson (1974: 127) stated that the specimen (USNM 85934) figured by Lea (1858: 298, pl. 26, fig. 12) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. An additional paralectotype is USNM 85934a. This species is a synonym of *Ensidens ingallsianus* (Lea, 1852) (Brandt, 1974: 288).

Unio scobinatus Lea, 1856: 93.

Paralectotypes: AMNH 30832 (14 specimens, 1 valve).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1867. Johnson (1974: 130) stated that the specimen (USNM 84145) figured by Lea (1858: 299–300, pl. 26, fig. 13) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are MCZ 175555 (Johnson, 1956: 134), 297361, and USNM 84145a, 84146. This species is the

type of the genus *Scabies* Haas, 1911 (Haas, 1911: 105) by subsequent designation as *S. scobinata* (Haas, 1969: 63), but is a junior synonym of *S. crispata* (Gould, 1843) (Brandt, 1974: 281).

Unio shuttleworthii Lea, 1856: 94.

Paralectotype: AMNH 30905 (1 specimen).

Locality: Australia.

Collector: R. J. Shuttleworth.

Remarks: ex Haines Collection 1870.

McMichael and Hiscock (1958: 415) stated that the holotype is USNM 84416, figured by Lea (1858: 304–305, pl. 28, fig. 19). Since Lea did not designate a holotype, McMichael and Hiscock's action constituted a lectotype designation. *Alathyria p. pertexta* Iredale, 1934, is the replacement name for *U. shuttleworthii* Lea, 1856 (non Kuster, 1855) (Smith, 1992: 16).

Unio sloatianus Lea, 1840: 287.

Holotype: AMNH 56104.

Locality: Chattahoochee River, Georgia, United States, North America.

Collector: L. W. Sloat.

Remarks: ex Jay Collection 1872. AMNH 56104 is the holotype by monotypy and was figured by Lea (1842: 271, pl. 16, fig. 33) and is fig. 12 herein. This species is the type of the genus *Elliptoideus* Frierson, 1927, by monotypy (Frierson, 1927: 33).

Unio superbus Lea, 1843: 1.

Paralectotype: AMNH 30776 (1 specimen).

Locality: New Holland [Australia].

Collector: Unknown.

Remarks: ex Jay Collection 1895. This species was described in a printed sheet dated Aug. 19, 1843 (Scudder, 1885: 42). AMNH 30776 is a type (W. J. Clench, on specimen label). Johnson (1974: 142) stated that the specimen (USNM 83934) figured by Lea (1846: 281, pl. 42, fig. 11) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This species is the type of the genus *Physunio* Simpson, 1900 (Simpson, 1900: 830), as the senior synonym of *U. gravidus* Lea, 1856 (Haas, 1969: 89–90).

Unio tumidulus Lea, 1856: 93.

Paralectotype: AMNH 30815 (1 valve).

Locality: Siam [Thailand], Asia.

Collector: S. R. House.

Remarks: ex Haines Collection 1912. Johnson (1974: 149) stated that the specimen (USNM 84612) figured by Lea (1858: 295–296, pl. 25, fig. 9) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. This taxon is a subspecies of *Unianдра contradens* (Lea, 1838) as *U. c. tumidula* (Brandt, 1974: 290).

Unio verreauianus Lea, 1856: 94.

Paralectotypes: AMNH 30819 (4 specimens).

Locality: Cape of Good Hope, South Africa, Africa.

Collector: E. Verreau.

Remarks: ex Haines Collection 1918. Johnson (1974: 153) stated that the specimen (USNM 84418) figured by Lea (1858: 301–302, pl. 27, fig. 16) was the holotype. Since Lea did not designate a holotype, Johnson's action constituted a lectotype designation. Additional paralectotypes are USNM 84418a and MCZ 240231 (ex AMNH). This species is a synonym of *Caferia caffra* (Krauss, 1848) (Haas, 1969: 155).

Venerupis peruviana Jay, 1839: 113, pl. 1, figs. 14, 15.

Syntypes: AMNH 56118 (3 specimens).

Locality: Coast of Peru, Pacific Ocean.

Collector: Unknown.

Remarks: ex Jay Collection 486. Richards and Old (1969) stated that "one ... is marked 'fig'd type' and is therefore taken to be the holotype." This was a flawed rationale, as no specimen can be a holotype unless so stated in the original publication, and the action of Richards and Old (1969) should not be considered a lectotype designation. In fact, no single specimen matches Jay's figures well enough to be definitely called the figured specimen. This species is a synonym of *Petricola (Rupellaria) denticulata* Sowerby, 1834 (Keen, 1971: 199).

Volsella (Volsella) salvadora Hertlein and Strong, 1946: 73, pl. 1, figs. 7, 11.

Paratype: AMNH 160345 (1 valve).

Locality: Station 199-D-6, 13°02'30"N, 87°29'30"W, 4 fathoms, off Monypenny Point, Gulf of Fonseca, Nicaragua, Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition, Dec. 24, 1937.

Remarks: ex NYZS. The holotype is CASIZ 064332 (Hertlein and Strong, 1946: 73). This species is placed in *Lioberus* Dall, 1898, as *L. salvadoricus* (Bernard, 1983: 19).

Type bivalve specimens that should be in the AMNH Collection but were not present or cataloged as of June 1995 (see Introduction for details).

Anodonta senegalensis Lea, 1859: 154 (paralectotype).

Cardium setosum Redfield, 1848: 168, pl. 11, figs. 1a–c (syntype—AMNH 29954?; specimen needs to be seen for verification of status).

Chlamys (Chlamys) incantata Hertlein, 1972: 2–4, figs. 1–5 (paratype).

Corbicula erosa Prime, 1861a: 126 (paralectotype).

Corbicula gracilis Prime, 1862b: 389–390, pl. 14, fig. 7 (paralectotype).

Corbicula lydigiana Prime, 1861b: 355–356 (paralectotype).

Corbicula notata Prime, 1861a: 127 (paralectotype).

Corbicula violacea Prime, 1861a: 128 (paralectotype).

Cyrena fortis Prime, 1861b: 355 (paralectotype).

Glauconome sumatrensis Prime, 1862b: 384, pl. 14, fig. 5 (paralectotype).

Glycymeris gordoni Usticke, 1959: 2–3, pl. 1, fig. 1 (holotype).

Mactra gigas Usticke, 1969: 7 (syntype).

Pecten yessoensis Jay, 1856: 293–294, pl. 3, figs. 3, 4; pl. 4, figs. 1, 2 (syntype).

Phacoides cancellata Usticke, 1969: 6, pl. 1, fig. 1578 (syntype).

Psammobia olivacea Jay, 1856: 292, pl. 1, figs. 8, 9 (syntype).

Unio evitatus Lea, 1866: 133 (paralectotype).

Unio rajahensis Lea, 1841: 30 (paralectotype).

Venus (Gemma) manhattensis Prime, 1862a: 482–483, text fig. (syntype).

CLASS SCAPHOPODA

Annulipulsellum euzkadii Scarabino, 1986b: 4–5, figs. 8–20.

Paratype: AMNH 202773 (1 specimen).

Locality: Cruise 25, Station 307, 3835–3862 m, 12°34.4'–12°40.8'N, 58°59.3'–59°09.2'W, Atlantic Ocean.

Collector: R/V "Knorr," March 3, 1972.

Remarks: ex Scarabino. The holotype is MCZ 293952 and additional paratypes are MCZ 293953–293955 (also MCZ 293956, not cited in publication), MNHN-M 14770–14772 and MNHN-P (Scarabino, 1986b: 4).

Cadulus acus Dall, 1889: 432, pl. 27, fig. 11. Paralectotypes: AMNH 148337 (2 specimens), 148342 (8 specimens).

Locality: 30 fathoms, Samana Bay, Santo Domingo, Atlantic Ocean.

Collector: Capt. Couthouy, 1850.

Remarks: ex USNM 95379 (now USNM 887454). Henderson (1920: 140) stated that USNM 95379 was the holotype. Since Dall did not designate a holotype, Henderson's action constituted a lectotype designation. Additional paralectotypes are USNM 95379 (Henderson, 1920: 140) (now USNM 887454) and MCZ 7751 (Turner, 1955: 317). MCZ 7751 is not the lectotype, despite being cited as the holotype by Turner (1955: 317). This species is placed in *Gadila* Gray, 1847 (Scarabino *in* Rios, 1994: 309).

Cadulus albicomatus Dall, 1890: 295, pl. 9, fig. 8.

Syntype: AMNH 146292 (1 specimen).

Locality: Station 2792, 00°37'S, 81°00'W, 401 fathoms, off Manta, Ecuador, Pacific Ocean.

Collector: "Albatross," March 2, 1888.

Remarks: ex USNM 602251. Additional syntypes are USNM 87539, 602251, and SBMNH 35018 (Scott et al., 1990: 18). This species is the type of the subgenus *Striocadulus* Emerson, 1962, by original designation (Emerson, 1962: 478).

Cadulus (Platyschides) arctus Henderson, 1920: 124, pl. 19, fig. 19.

Paratype: AMNH 148348 (1 specimen).

Locality: Station 2404, 28°44'N, 85°16'W, 60 fathoms, off Cape San Blas, Florida, United States, Atlantic Ocean.

Collector: "Albatross," March 15, 1885.

Remarks: ex USNM 151830. The holotype is USNM 323955 and additional paratypes are USNM 151830 (Henderson, 1920: 124).

Cadulus (Gadila) atlanticus Henderson, 1920: 137–139, pl. 20, fig. 2.

Paratypes: AMNH 148338 (2 specimens).

Locality: Station 2115, 35°49'30"N, 74°34'45"W, 843 fathoms, off Cape Hatteras, North Carolina, United States, Atlantic Ocean.

Collector: "Albatross," Nov. 11, 1883.

Remarks: ex USNM. The holotype is USNM 78238 and additional paratypes are in USNM (Henderson, 1920: 139) and MCZ 186815 (ex Station 2682).

Cadulus (Gadila) californicus Pilsbry and Sharp, 1898: 180–181, pl. 34, figs. 5–8.

Paralectotypes: AMNH 148391 (4 specimens).

Locality: Station 2923, 32°40'30"N, 117°31'30"W, 822 fathoms, off San Diego, California, United States, Pacific Ocean.

Collector: "Albatross," Jan. 19, 1889.

Remarks: ex USNM 107698a. The lectotype (USNM 107698) was designated by Shimmek (1989: 235). Additional paralectotypes are USNM 107698a (Pilsbry and Sharp, 1898: 181), 203136, and SBMNH 35038 (Scott et al., 1990: 18). This species is placed in *Polyschides* Pilsbry and Sharp, 1898 (Turgeon et al., 1988: 51).

Cadulus (Polyschides) dalli Pilsbry and Sharp, 1898: 155–156, pl. 30, figs. 19–23.

Syntypes: AMNH 148395 (3 specimens).

Locality: Station 2783, 51°02'30"S, 74°08'30"W, 122 fathoms, West Coast of Patagonia, Chile, Pacific Ocean.

Collector: "Albatross," Feb. 6, 1888.

Remarks: ex USNM 122746 (now USNM 602253). Additional syntypes are USNM 122736, 122746 (not 123746 as cited in Pilsbry and Sharp, 1898: 156), 122746 (now USNM 602253), and SBMNH 35039 (Scott et al., 1990: 18).

Cadulus (Gadila) fusiformis Pilsbry and Sharp, 1898: 193–194, pl. 35, fig. 14.

Paratype: AMNH 148398 (1 specimen).

Locality: 4 fathoms, San Pedro, California, United States, Pacific Ocean.

Collector: J. G. Cooper.

Remarks: ex USNM 602247. The holotype is USNM 133809 (Pilsbry and Sharp, 1898: 194). Additional paratypes are USNM 7957 and 602247. This species is a synonym of *C. aberrans* Whiteaves, 1887 (Shimek, 1989: 235).

Cadulus (Cadulus) halius Henderson, 1920: 149, pl. 20, fig. 12.

Paratype: AMNH 148332 (1 specimen).

Locality: Station 115, 120 fathoms, off English Harbor, Antigua, Atlantic Ocean.

Collector: University of Iowa Expedition of 1918.

Remarks: ex USNM 314936. The holotype was in the State University of Iowa (now USNM 596599) and additional paratypes are USNM 314936 (Henderson, 1920: 149).

Cadulus (Platyschides) macleani Emerson, 1978: 122, figs. 1-3.

Paratype: AMNH 213440 (1 specimen).

Locality: Station 66-22, 24°55'N, 109°46'W, 18-55 m on sand and shell bottom, Mueritos Bay, Baja California Sur, Mexico, Pacific Ocean.

Collector: McLean, Oringer and Marincovich, April 8, 1966.

Remarks: ex LACM 1887. The holotype is LACM 1886 and additional paratypes are LACM 1887 (Emerson, 1978: 122).

Cadulus (Gadila) mayori Henderson, 1920: 133-134, pl. 19, fig. 3.

Paratypes: AMNH 148339 (7 specimens).

Locality: Station 33, 16 fathoms, Tortugas, Florida, United States, Atlantic Ocean.

Collector: "Eolis," 1911.

Remarks: ex USNM 314784. The holotype is USNM 314783 and additional paratypes are USNM 157809, 314784-314816, 323953 (Henderson, 1920: 133-134), MCZ 48411 (Turner, 1955: 319), and ANSP 44914 from "Eolis" station 43 (ex USNM 314786). This species is placed in *Gadila* Gray, 1834 (Turgeon et al., 1988: 51).

Cadulus (Platyschides) miamensis Henderson, 1920: 129-130, pl. 19, fig. 18.

Paratypes: AMNH 148351 (2 specimens).

Locality: Station 340, 209 fathoms, off Fowey

Light, Florida, United States, Atlantic Ocean.

Collector: "Eolis," 1917.

Remarks: ex USNM 314772a. The holotype is USNM 314772 and additional paratypes are USNM 108176a, 314769-314781, 330522 (Henderson, 1920: 130), MCZ 48413 (Turner, 1955: 319), and ANSP 44912 from "Eolis" station 348 (ex USNM 314774). This species is placed in *Polyschides* Pilsbry and Sharp, 1898 (Turgeon et al., 1988: 51).

Cadulus (Polyschides) nedallisoni Emerson, 1978: 120-122, figs. 4-6.

Holotype: AMNH 160349.

Paratypes: AMNH 183875 (39 specimens), 183875a (1 specimen).

Locality: Station 200-D-19, 12°28'03"N, 87°12'39"W, 22-24 m, Corinto, Nicaragua, Pacific Ocean.

Collector: Eastern Pacific "Zaca" Expedition, Jan. 5, 1938.

Remarks: ex NYZS. Additional paratypes are LACM 2117 (Groves, personal commun.). The holotype was figs. 4, 5 and AMNH 183875a was fig. 6 (Emerson, 1978).

Cadulus (Platyschides) nitidus Henderson, 1920: 129, pl. 19, fig. 9.

Paratypes: AMNH 148353 (3 specimens).

Locality: Station 6062, 25-30 fathoms, Mayaguez Harbor, Puerto Rico, Atlantic Ocean.

Collector: "Fish Hawk," Jan. 20, 1899.

Remarks: ex USNM 161582. The holotype was USNM 161582a (now USNM 887474) and additional paratypes are USNM 161582 (Henderson, 1920: 129).

Cadulus (Gadila) panamensis Sharp and Pilsbry in Pilsbry and Sharp, 1898: 191-192, pl. 36, figs. 23-25.

Paralectotypes: AMNH 148402 (6 specimens).

Locality: Station 2805, 07°56'S, 79°41'30"W, 51.5 fathoms, Panama Bay, Panama, Pacific Ocean.

Collector: "Albatross," March 30, 1888.

Remarks: ex USNM 122795 (now USNM 602248). The lectotype (USNM 122795) was designated by Emerson (1971: 78), who considered *C. perpusillus* (Sowerby, 1832) the senior synonym. Additional paralectotypes were USNM 122795 (now USNM

602248) and SBMNH 35014 (Scott et al., 1990: 18).

Cadulus (Cadulus) platensis Henderson, 1920: 147–148, pl. 20, fig. 14.

Paratype: AMNH 148340 (1 specimen).

Locality: Station 2668, 30°58'30"N, 79°38'30"W, 294 fathoms, off Fernandina, Florida, United States, Atlantic Ocean.

Collector: "Albatross," May 5, 1886.

Remarks: ex USNM 108172. The holotype was USNM 330845a (now USNM 887476) and additional paratypes are USNM 95447, 108172, 108279, 330845 (Henderson, 1920: 147–148).

Cadulus (Platyschides) providensis Henderson, 1920: 122, pl. 18, fig. 14.

Paratype: AMNH 148352 (1 specimen).

Locality: Station 2668, 30°58'30"N, 79°38'30"W, 294 fathoms, off Fernandina, Florida, United States, Atlantic Ocean.

Collector: "Albatross," May 5, 1886.

Remarks: ex USNM 108176. The holotype is USNM 94055 and additional paratypes are USNM 94055a, 108176 (Henderson, 1920: 122).

Cadulus (Gadila) regularis Henderson, 1920: 137, pl. 19, fig. 14.

Paratypes: AMNH 148341 (21 specimens).

Locality: Station 2668, 30°58'30"N, 79°38'30"W, 294 fathoms, off Fernandina, Florida, United States, Atlantic Ocean.

Collector: "Albatross," May 5, 1886.

Remarks: ex USNM 108175. The holotype was USNM 330676a (now USNM 887475) and additional paratypes are USNM 108175, 108278, 330675 (Henderson, 1920: 137).

Cadulus (Platyschides) simpsoni Henderson, 1920: 127–128, pl. 19, fig. 17.

Paratype: AMNH 148357 (1 specimen).

Locality: Station 6062, 25–30 fathoms, Mayaguez, Puerto Rico, Atlantic Ocean.

Collector: "Fish Hawk," Jan. 20, 1899.

Remarks: ex USNM 161580a. The holotype is USNM 161580 and additional paratypes are USNM 161580a, 314677, 314932 (Henderson, 1920: 127).

Cadulus (Gadila) striatus Dall in Pilsbry and Sharp, 1898: 179–180, pl. 35, figs. 9–13.

Syntypes: AMNH 148405 (3 specimens).

Locality: Station 3354, 07°09'45"N, 80°50'W, 322 fathoms, Gulf of Panama, Panama, Pacific Ocean.

Collector: "Albatross," Feb. 23, 1891.

Remarks: ex USNM 122992 (now USNM 602252). Additional syntypes are USNM 122992 (Dall in Pilsbry and Sharp, 1898: 180) and were 122992 (now USNM 602252); also SBMNH 35023 (Scott et al., 1990: 18).

Cadulus (Polyschides) tetrodon Pilsbry and Sharp, 1898: 151–152, pl. 29, figs. 14–18.

Paralectotype: AMNH 148360 (1 specimen).

Locality: 5 mi off Cape Florida, 8 fathoms, Florida, United States, Atlantic Ocean.

Collector: W. H. Rush.

Remarks: ex USNM 84953. Henderson (1920: 101) stated that ANSP 71070 was the "type." Since Pilsbry and Sharp did not designate a holotype, Henderson's action constituted a lectotype designation. Additional paralectotypes are ANSP 71070 (Pilsbry and Sharp, 1898: 152) and USNM 84953 (Henderson, 1920: 101). This species is a synonym of *Polyschides tetraschistus* (Watson, 1879) (Scarabino in Rios, 1994: 310).

Cadulus (Cadulus) transitorius transitorius Henderson, 1920: 143–144, pl. 19, fig. 6.

Paratypes: AMNH 148334 (8 specimens).

Locality: Station 2668, 30°58'30"N, 79°38'30"W, 294 fathoms, off Fernandina, Florida, United States, Atlantic Ocean.

Collector: "Albatross," May 5, 1886.

Remarks: ex USNM 108170. The holotype is USNM 330589 and additional paratypes are USNM 108170, 314821–314823, 323779, 323813, and 330676 (not 330678 as cited in Henderson, 1920: 144).

Cadulus (Gadila) verrilli Henderson, 1920: 136–137, pl. 19, fig. 13.

Paratype: AMNH 148343 (1 specimen).

Locality: Station 871, 115 fathoms, off Martha's Vineyard, Massachusetts, United States, Atlantic Ocean.

Collector: "Fish Hawk," Sept. 4, 1880.

Remarks: ex USNM 38693. The holotype is USNM 38693 and additional paratypes are USNM 38693, 38694, 61144 (Henderson, 1920: 137), and MCZ 186819 (ex Station 871).

Costentalina vema Scarabino, 1986b: 2–3, pl. 1, figs. 1–7.

Holotype: AMNH 202762.

Paratype: AMNH 202763 (1 specimen).

Locality: V-17-81, 44°33'S, 44°19'W, 5332 m (AMNH 202762); V-17-85, 47°35'S, 43°21'W, 5495 m (AMNH 202763), Argentine Basin, Argentina, Atlantic Ocean. Collector: R/V "Vema," May 27, 1961 (AMNH 202762) and June 6, 1961 (AMNH 202763).

Remarks: AMNH 202762 was the figured specimen (Scarabino, 1986b).

Dentalium (Antalis) agile subagile Henderson, 1920: 38–39, pl. 4, fig. 8.

Paratype: AMNH 148306 (1 specimen)

Locality: Station 2513, 43°34'N, 63°56'30"W, 134 fathoms, between Halifax and Le Havre Banks, Nova Scotia, Canada, Atlantic Ocean.

Collector: "Albatross," July 11, 1885.

Remarks: ex USNM 50191. The holotype was USNM 50191a (now USNM 887470) and additional paratypes are USNM 50188, 50189, 50191–50193, 52302, 314325 (Henderson, 1920: 39), and ANSP 119158 from "Eolis" station 2513 (ex Henderson). This taxon is considered a full species (Abbott, 1974: 385).

Dentalium (Antalis) bartletti Henderson, 1920: 55–56, pl. 8, figs. 2, 7.

Paratypes: AMNH 148300, 148301 (1 specimen each).

Locality: Station 230, 13°13'20"N, 61°18'45"W, 464 fathoms, off St. Vincent Island, Atlantic Ocean.

Collector: "Blake," Feb. 20, 1879.

Remarks: ex USNM 95371. The holotype is USNM 224986 and additional paratypes are USNM 93121, 93840, 94081, 95368 (not 94368), 95369–95371, 314449, 314450, 323776, 330524 (Henderson, 1920: 55–56).

Dentalium (Antalis) ceratum flavum Henderson, 1920: 51–52, pl. 7, fig. 1.

Paratype: AMNH 148298 (1 specimen).

Locality: Station 325, 95 fathoms, off Sand Key, Florida, United States, Atlantic Ocean.

Collector: "Eolis," 1917.

Remarks: ex USNM 314400. The holotype is USNM 314391 and additional paratypes

are USNM 314391a–314405, 323016 (Henderson, 1920: 52) and ANSP 132113 from "Eolis" station 333 (ex Henderson). This subspecies is a synonym of *D. (A.) ceratum* Dall, 1881 (Abbott, 1974: 385).

Dentalium (Episiphon) johnsoni Emerson, 1952: 5–6, pl. 1, fig. 2.

Paratypes: AMNH 148289 (2 specimens).

Locality: Station 25, 18°32'15"N, 66°22'10"W, 240–300 fathoms, off Puerto Rico, Atlantic Ocean.

Collector: First Johnson-Smithsonian Expedition, Feb. 7, 1933.

Remarks: ex USNM 429714. The holotype is USNM 603543 and additional paratypes are USNM 429714 (Emerson, 1952: 5). This species is a synonym of *Episiphon di-dymum* (Watson, 1879) (Scarabino in Rios, 1994: 308).

Dentalium (Dentalium) rebeccaense Henderson, 1920: 31–32, pl. 3, fig. 2.

Paratype: AMNH 148314 (1 specimen).

Locality: Station 33, 16 fathoms, between Tortugas and Rebecca Shoals, Florida, United States, Atlantic Ocean.

Collector: "Eolis," 1911.

Remarks: ex USNM 314293a. The holotype is USNM 314293 and additional paratypes are USNM 314293a, 314294 (Henderson, 1920: 32). This species is a synonym of *Dentalium americanum* Chenu, 1843 (Scarabino in Rios, 1994: 305).

Dentalium solidum Verrill, 1884: 215–216.

Paralectotypes: AMNH 148370 (3 specimens).

Locality: Station 2084, 40°15'50"N, 67°05'15"W, 1290 fathoms, off Georges Bank, New England, United States, Atlantic Ocean.

Collector: "Albatross," Sept. 5, 1883.

Remarks: ex USNM 34911. The lectotype (USNM 34687) was selected and figured by Henderson (1920: 62–63, pl. 9, fig. 3). Additional paralectotypes are USNM 34687a, 34688, 34904, 34911, 35635, 35636, 35645 (Verrill, 1884: 216). None of the other specimens cited by Henderson (1920: 63) under *D. (Fissidentalium) meridionale verrilli* (new name for *D. solidum* Verrill, 1884, not *D. solidum* Hutton, 1873) are from Verrill's original type series. This species is a synonym of *Fissidentalium*

candidum (Jeffreys, 1877) (Scarabino *in* Rios, 1994: 307).

Dentalium (Episiphon) sowerbyi pelliceri Henderson, 1920: 80–81, pl. 13, figs. 7–9.

Paratypes: AMNH **148322** (35 specimens), **148323** (6 specimens).

Locality: Station 203, 3–12 fathoms, Cabanas Harbor (AMNH 148322) and Station 208, 1–12 fathoms, Bahia Honda (AMNH 148323), Cuba, Atlantic Ocean.

Collector: “Barrera”.

Remarks: ex USNM 314565a (AMNH 148323) and USNM 314567 (AMNH 148322). The holotype is USNM 314565 and additional paratypes are USNM 314565a–14573 (Henderson, 1920: 81) and ANSP 119157 from “Eolis” station 203 (ex Henderson). This subspecies is a synonym of *Episiphon sowerbyi* (Guilding, 1834) (Scarabino *in* Rios, 1994: 308).

Pulsellum salishorum E. Marshall, 1980: 149–152, figs. 1–6.

Paratypes: AMNH **198610** (6 specimens).

Locality: 27 m, sticky mud, East Sound, Orcas Island, San Juan Islands, Washington, United States, Pacific Ocean.

Collector: E. Marshall, Aug. 16, 1977.

Remarks: ex E. Marshall. The holotype is USNM 782263 and additional paratypes are USNM 782264, LACM 1929, ANSP 352474, National Museum of Canada 86068 and Washington State Museum 35663 (E. Marshall, 1980: 150).

Siphonodentalium isaotakii Habe, 1953: 297, 299.

Paratypes: AMNH **148421** (6 specimens).

Locality: Tokyo Bay, Honshu, Japan, Pacific Ocean.

Collector: Unknown.

Remarks: ex Habe. The holotype is NSMT 54817 and an additional paratype is NSMT 54818 (Inaba and Oyama, 1977: 52). The

species was figured by Habe (1963: text fig. 52).

Siphonodentalium japonicum Habe, 1960: 294.

Paratypes: AMNH **148416** (3 specimens).

Locality: Tomioka, Amakusa, Kumamoto Prefecture, Japan, Pacific Ocean.

Collector: Unknown.

Remarks: ex Habe. The holotype is NSMT 54819 and an additional paratype is NSMT 54820 (Inaba and Oyama, 1977: 56). The species was figured by Habe (1963: text figs. 35, 36).

Siphonodentalium okudai Habe, 1953: 297, 299, figs. 745, 759, 760.

Paratype: AMNH **148420** (1 specimen).

Locality: Akkeshi Bay, Hokkaido, Japan, Pacific Ocean.

Collector: Unknown.

Remarks: ex Habe. The holotype is NSMT 54867 and an additional paratype is NSMT 54868 (Inaba and Oyama, 1977: 90).

Wemersoniella duartei Scarabino, 1986a: 5–6, 8, 12–19, figs. 6–11, 17–21.


Holotype: AMNH **202770**.

Paratypes: AMNH **202771** (approx. 20 specimens), **202772** (approx. 30 specimens).


Locality: V-17-80, 43°58'S, 52°09'W, 5781 m (AMNH 202770, 202771); V-17-81, 44°33'S, 49°19'W, 5332 m (AMNH 202772), “Cuenca Argentina,” Atlantic Ocean.

Collector: R/V “Vema,” May 26, 1961 (AMNH 202770, 202771) and May 27, 1961 (AMNH 202772).

Remarks: AMNH 202770 was figs. 6–8 (Scarabino, 1986a). Some paratypes are fragmented. Additional paratypes were cited as MCZ 293962, MNHN-M 14760, 14761, and MNHN-P (Scarabino, 1986a: 5), although in the case of the MCZ lot only a placeholder (“7 lots to come”) is present in the MCZ collection.



1



2

3

Figs. 1-3. 1. *Plaxiphora excavata* Pilsbry, 1893 (syntypes—AMNH 21933). 2. *Anodonta montezuma* Lea, 1841 (holotype—AMNH 30871). 3. *Anodonta subcylindracea* Lea, 1838 (syntype—AMNH 31131a). All figures $\times 1$.







Fig. 4. *Batissa gigantea* Prime, 1860 (holotype—AMNH 31427) $\times \frac{1}{2}$.



Figs. 5-7. 5. *Cyrena jayensis* Lea, 1834 (syntype—AMNH 31337) $\times 1$. 6. *Glycymeris gordoni tobagoensis* Usticke, 1959 (syntype—AMNH 271074) $\times 1\frac{1}{2}$. 7. *Iridina coelestis* Lea, 1838 (paralectotype—AMNH 31292) $\times 1$.



Figs. 8, 9. 8. *Unio callosus* Lea, 1841 (holotype—AMNH 30561) $\times 1$. 9. *Unio cucumoides* Lea, 1840 (holotype—AMNH 30897) $\times \frac{3}{4}$.



Figs. 10, 11. **10.** *Unio hainesianus* Lea, 1856 (lectotype, herein—AMNH 31070) $\times \frac{1}{2}$. **11.** *Unio lutens* Lea, 1856 (holotype—AMNH 29109) $\times 1$.




Fig. 12. *Unio sloatianus* Lea, 1840 (holotype—AMNH 56104) $\times 1$.

REFERENCES

- Abbott, R. T.
- 1958. The marine mollusks of Grand Cayman Island, British West Indies. Monogr. Acad. Nat. Sci. Philadelphia. 11: 138 pp., 5 pls., 7 text figs.
 - 1973. *Acteon eloiseae*, a new opisthobranch from Arabia. *Nautilus* 87(4): 91–92, 5 figs.
 - 1974. American seashells, 2nd ed. New York: Van Nostrand. 663 pp., 24 pls., 6405 text figs.
- Adamkewicz, L., and M. G. Harasewych
- 1994. Molecular markers and the biogeography of *Donax*. Am. Malacol. Union Program and Abstr. 60th annual meeting: 24.
- Adams, A., and L. Reeve
- 1848–50. Mollusca. In A. Adams, The zoology of the voyage of H. M. S. Samarang; under the command of Captain Sir Edward Belcher, C. B., F. R. A. S., F. G. S. during the years 1843–1846. London: Reeve, Benham and Reeve. 87 pp., 24 pls.
- Allen, J. A., and F. J. Hannah
- 1986. A reclassification of the Recent genera of the subclass Protobranchia (Mollusca: Bivalvia). *J. Conchol.* 32(4): 225–249, 50 figs.
- Anthony, J. G.
- 1865. Descriptions of two new species of *Mondocondylaea*. *Am. J. Conchol.* 1(3): 205–206, pl. 18, figs. 1, 3.
- Bernard, F. R.
- 1983. Catalogue of the living Bivalvia of the eastern Pacific Ocean: Bering Strait to Cape Horn. *Can. Spec. Publ. Fish. Aquat. Sci.* 61: 102 pp.
- Bernard, F. R., Y. Y. Cai, and B. Morton
- 1993. Catalogue of the living marine bivalve molluscs of China. Hong Kong: Hong Kong Univ. Press. 146 pp., 6 figs.
- Berry, S. S.
- 1953. West American razor-clams of the genus *Ensis*. *Trans. San Diego Soc. Nat. Hist.* 11(15): 393–404, pl. 29, figs. 5–6; 4 text figs.
- Bieler, R.
- 1988. A note on unjustified emendations. *Veliger* 30(4): 423–424.
- Bosch, D. T., S. P. Dance, R. G. Moolenbeek, and P. G. Oliver
- 1995. Seashells of Eastern Arabia. Dubai, United Arab Emirates: Motivate Publishing. 296 pp.
- Boss, K. J.
- 1965. A new mollusk (Bivalvia, Erycinidae) commensal on the stomatopod crustacean *Lysiosquilla*. *Am. Mus. Novitates* 2215: 11 pp., 3 figs.
- Boss, K. J., and R. D. Turner
- 1980. The giant white clam from the Galapagos Rift, *Calyptogena magnifica* species novum. *Malacologia* 20(1): 161–194, 13 figs.
- Bourguignat, J. R.
- 1885. Monographie d'un nouveau genre d'acéphale du Lac Tanganyika. *Bull. Soc. Malacol. France*, Paris, 2: 1–12, 1 pl.
- Boyko, C. B.
- 1994. Catalog of Recent type specimens in the Department of Invertebrates, American Museum of Natural History. I. Micro-Invertebrates (Phyla Sarcomastigophora, Gnathostomula, Gastrotricha, Rotifera, and Tardigrada). *Am. Mus. Novitates* 3106: 44 pp.
- Brandt, R.
- 1974. The non-marine aquatic Mollusca of Thailand. *Arch. Molluskenkd.* 105(1–4): 423 pp., 30 pls.
- Burch, J. B.
- 1975. Freshwater unionacean clams (Mollusca: Pelecypoda) of North America, 2nd ed. Hamburg, MI: Malacological Publications. 204 pp., 252 figs.
- Clarke, A. H.
- 1973. The freshwater molluscs of the Canadian Interior Basin. *Malacologia* 13(1–2): 1–509, 28 pls., 9 text figs.
- Clench, W. J., and R. D. Turner
- 1956. Freshwater mollusks of Alabama, Georgia, and Florida from the Escambia to the Suwannee River. *Bull. Florida State Mus.* 1(3): 97–239, 9 pls.
- Coan, E. V.
- 1979. Recent eastern Pacific species of the crassatellid bivalve genus *Crassinella*. *Veliger* 22(1): 1–11, 22 figs.
 - 1988. Recent eastern Pacific species of the bivalve genus *Semele*. *Ibid.* 31(1–2): 1–42, 64 figs.
- Conrad, T. A.
- 1869. Notes on Recent and fossil shells, with descriptions of new genera. *Am. J. Conchol.* 4(4): 246–249.
- Dall, W. H.
- 1889. Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico (1877–78) and in the Caribbean Sea (1879–80)...XXIX. Report on the Mollusca. Part II. Gastropoda and Scaphopoda.

- Bull. Mus. Comp. Zool. 18: 1–492, pls. 10–40.
1890. Scientific results of explorations by the U.S. Fish Commission Steamer Albatross. VII. Preliminary report on the collection of Mollusca and Brachiopoda obtained in 1887–88. Proc. U.S. Natl. Mus. 12(773): 219–362, pls. 5–14.
1908. Reports on the dredging operations off the west coast of Central America to the Galapagos, to the west coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz. . XIV. The Mollusca and the Brachiopoda. Bull. Mus. Comp. Zool. 43(6): 205–487, 19 pls.
- Dell, R. K.
1963. The identity of *Yoldia (Aequiyoldia) eightsi* (Couthouy, in Jay, 1839). Proc. Malacol. Soc. London 35(6): 247–249, 1 fig.
1990. Antarctic Mollusca: with special reference to the fauna of the Ross Sea. R. Soc. New Zealand Bull. 27: 311 pp., 482 figs.
- Emerson, W. K.
1952. The scaphopod mollusks collected by the first Johnson-Smithsonian Deep-Sea Expedition. Smithson. Misc. Collect. 117(6): 1–14, 1 pl.
1962. A classification of the scaphopod mollusks. J. Paleontol. 36(3): 461–482, pls. 76–80.
1971. *Cadulus (Gadila) perpusillus* (Sowerby, 1832), an earlier name for *C. (G.) panamensis* Sharp and Pilsbry, 1898. Nautilus 84(3): 77–81, 4 figs.
1978. Two new eastern Pacific species of *Cadulus*, with remarks on the classification of the scaphopod mollusks. Ibid. 92(3): 117–123, 9 figs.
- Faber, M. J.
1988. Studies on West Indian marine mollusks 13. The malacological taxa of Gordon W. Nowell-Usticke. De Kreukel 24(4–5): 67–102.
- Ferreira, A. J.
1974. The genus *Lepidozona* in the Panamic Province, with the description of two new species (Mollusca: Polyplacophora). Veliger 17(2): 162–180, 24 figs.
- 1978a. A new species of chiton (Neoloricata: Ischnochitonidae) from the Galapagos Islands. Bull. South. California Acad. Sci. 77(1): 36–39, 2 figs.
- 1978b. The genus *Lepidozona* (Mollusca: Polyplacophora) in the temperate eastern Pacific, Baja California to Alaska, with the description of a new species. Veliger 21(1): 19–44, 33 figs.
1979. The genus *Callistochiton* Dall, 1879 (Mollusca: Polyplacophora) in the eastern Pacific, with the description of a new species. Ibid. 21(4): 444–466, 27 figs.
- Frierson, L. S.
1927. A classified and annotated check list of the North American naiades. Waco, TX: Baylor Univ. Press. 111 pp.
- Gosliner, T. M.
1980. The systematics of the Aeolidacea (Nudibranchia: Mollusca) of the Hawaiian Islands, with descriptions of two new species. Pac. Sci. 33(1): 37–77, 20 figs.
1981. The South African Janolidae (Mollusca, Nudibranchia) with the description of a new genus and two new species. Ann. S. African. Mus. 86(1): 42 pp., 28 figs.
- Gould, A. A.
1850. Shells from the United States Exploring Expedition. Proc. Boston Soc. Nat. Hist. 3: 292–296.
1852. Mollusca & shells. In United States Exploring Expedition. During the years 1838, 1839, 1840, 1841, 1842. Under the command of Charles Wilkes, U.S.N. 12: 497 pp., 621 figs.
- Haas, F.
- 1910–1920. Die unioniden. Syst. Conchol. Cab. 9 (2, II): 1–344, 73 pls.
1936. Binnen-mollusken aus inner-Africa, hauptsachlich gesammelt von Dr. F. Haas wahrend der Schomburgk Expedition in den jahren 1931/32. Abh. Senckenburgischen Naturforsch. Ges. 431: 156 pp, 8 pls.
1969. Superfamilia Unionacea. Das Tierreich 88: 663 pp., 5 figs.
- Habe, T.
1953. Genera of Japanese shells. Pelecypoda and Scaphopoda 4: 281–326, figs. 731–770.
1960. New species of molluscs from the Amakusa Marine Biological Laboratory, Reihioku-cho, Amakusa, Kumamoto Pref., Japan. Publ. Seto Mar. Biol. Lab. 8(2): 289–298, 5 figs.
1963. A classification of the scaphopod mollusks found in Japan and its adjacent areas. Bull. Nat. Sci. Mus. (Tokyo) 6(3): 252–281, pls. 37–38, 56 text figs.
1977. Type specimens of four Japanese shells described by J. C. Jay. Venus Japanese J. Malacol. 36(3): 153–156, 2 pls.
- Henderson, J. B.
1920. A monograph of the East American scaphopod mollusks. U.S. Natl. Mus. Bull. 111: 177 pp., 20 pls.
- Herrington, H. B.
1962. A revision of the Sphaeriidae of North

- America (Mollusca: Pelecypoda). *Misc. Publ. Mus. Zool. Univ. Michigan* 118: 74 pp., 7 pls., 2 text figs.
- Hertlein, L. G.
 1972. Description of a new species of *Chlamys* (Mollusca: Pelecypoda) from the Galapagos Islands. *Proc. California Acad. Sci.*, 4th ser. 39(1): 1–6, 5 figs.
- Hertlein, L. G., and A. M. Strong
 1940. Eastern Pacific expeditions of the New York Zoological Society. XXII. Mollusks from the West coast of Mexico and Central America. Part I. *Zoologica (New York)* 25, part 4(25): 369–430, 2 pls.
1943. Eastern Pacific expeditions of the New York Zoological Society. XXXII. Mollusks from the West coast of Mexico and Central America. Part II. *Ibid.* 28, part 3(19): 149–168, 1 pl.
1946. Eastern Pacific expeditions of the New York Zoological Society. XXXIV. Mollusks from the West Coast of Mexico and Central America. Part III. *Ibid.* 31, part 2(5): 53–76, 1 pl.
1947. Eastern Pacific expeditions of the New York Zoological Society. XXXVI. Mollusks from the West coast of Mexico and Central America. Part V. *Ibid.* 31, part 4(10): 129–150, 1 pl.
1948. Eastern Pacific expeditions of the New York Zoological Society. XXXIX. Mollusks from the West coast of Mexico and Central America. Part VI. *Ibid.* 33, part 4(13): 163–198, 2 pls.
- 1949a. Eastern Pacific expeditions of the New York Zoological Society. XL. Mollusks from the West coast of Mexico and Central America. Part VII. *Ibid.* 34, part 2(9): 63–97, 1 pl.
- 1949b. Eastern Pacific expeditions of the New York Zoological Society. XLI. Mollusks from the West coast of Mexico and Central America. Part VIII. *Ibid.* 34, part 4(19): 239–258, 1 pl.
1950. Eastern Pacific expeditions of the New York Zoological Society. XLII. Mollusks from the West coast of Mexico and Central America. Part IX. *Ibid.* 35, part 4(19): 217–252, 2 pls.
1955. Marine mollusks collected by the "Askooy" Expedition to Panama, Colombia and Ecuador in 1941. *Bull. Am. Mus. Nat. Hist.* 107(2): 159–318, 3 pls.
- Higo, S., and Y. Goto
 1993. A systematic list of molluscan shells from the Japanese Islands and the adjacent area. Tokyo. 863 pp.
- [ICZN] International Commission on Zoological Nomenclature
 1985. International Code of Zoological Nomenclature (3rd ed.). 338 pp. International Trust for Zoological Nomenclature.
- Inaba, T., and K. Oyama
 1977. Catalogue of molluscan taxa described by Tadashige Habe during 1939–1975, with illustrations of hitherto unfigured species (for commemoration of his sixtieth birthday). Tokyo: 185 pp., 7 pls.
- Jay, J. C.
 1839. A catalogue of the shells, arranged according to the Lamarckian system; together with descriptions of new or rare species contained in the collection of John C. Jay, M. D. 3rd ed. New York: Wiley and Putnam. 126 pp., 10 pls.
1856. Report on the shells collected by the Japan Expedition, under the command of Commodore M. C. Perry, U.S.N., together with a list of Japan shells. In *Narrative of the Expedition of an American squadron to the China seas and Japan, performed in the years 1852, 1853, and 1854, under the command of Commodore M. C. Perry, United States Navy, by order of the government of the United States*, vol. 2. Washington, D.C. pp. 291–297, 5 pls.
- Johnson, R. I.
 1946. Joseph Pitty Couthouy. A bibliography and catalogue of his species. *Occas. Pap. Mollusks Mus. Comp. Zool. Harv. Univ.* 1(5): 33–40, pl. 8.
1956. The types of naiades (Mollusca: Unionidae) in the Museum of Comparative Zoology. *Bull. Mus. Comp. Zool.* 115(4): 101–142, 2 pls.
1959. The types of Corbiculidae and Sphaeriidae (Mollusca: Pelecypoda) in the Museum of Comparative Zoology, and a bio-bibliographic sketch of Temple Prime, an early specialist of the group. *Ibid.* 120(4): 431–479, 8 pls.
1970. The systematics and zoogeography of the Unionidae (Mollusca: Bivalvia) of the southern Atlantic Slope region. *Ibid.* 140(6): 263–450, 22 pls., 5 text figs.
1971. The types and figured specimens of Unionacea (Mollusca: Bivalvia) in the British Museum (Natural History). *Bull. Br. Mus. (Nat. Hist.) Zool.* 20(3): 75–108, 2 pls.
1974. Lea's unionid types or Recent and fossil taxa of Unionacea and Mutelacea introduced by Isaac Lea, including the location of all the extant types. *Spec. Occas. Publ. Dep. Mollusks Mus. Comp. Zool.* 2: 159 pp.

1979. The types of Unionacea (Mollusca: Bivalvia) in the Museum of Zoology, The University of Michigan. *Malacol. Rev.* 12: 29–36.
- Johnson, R. I., and H. B. Baker
1973. The types of Unionacea (Mollusca: Bivalvia) in the Academy of Natural Sciences of Philadelphia. *Proc. Acad. Nat. Sci. Philadelphia* 125(9): 145–186, 10 pls.
- Kaas, P., and R. A. Van Belle
1987. Monograph of living chitons (Mollusca: Polyplacophora) vol. 3. Suborder Ischnochitonina. *Ischnochitonidae: Chaetopleurinae, & Ischnochitoninae (pars). Additions to vols. 1 & 2.* Leiden: E. J. Brill. 302 pp., 117 figs.
1994. Monograph of living chitons (Mollusca: Polyplacophora) vol. 5. Suborder Ischnochitonina. *Ischnochitonidae: Ischnochitoninae (concluded); Callistoplacinae; Mopaliidae. Additions to volumes 1–4.* Leiden: E. J. Brill. 402 pp., 141 figs.
- Keen, M.
1971. Sea shells of tropical West America. Marine mollusks from Baja California to Peru, 2nd ed. Stanford, CA: Stanford Univ. Press. 1064 pp., 22 pls.
- Kennedy, G. L.
1989. Status of *Penitella gabbii* (Tryon, 1863) in the eastern and western Pacific, and description of the previously misidentified eastern Pacific species (Bivalvia: Pholadidae). *Veliger* 32(3): 313–319, 15 figs.
- Kennedy, G. L., and J. M. Armentrout
1989. A new species of chimney-building *Penitella* from the Gulf of Alaska (Bivalvia: Pholadidae). *Veliger* 32(3): 320–325, 13 figs.
- Klappenbach, M. A.
1962. Una nueva especie de "Eupera" (Moll. Pelecypoda) del Uruguay. *Rev. Mus. Argentino Cienc. Nat. "Bernardino Rivadavia"* 8(8): 101–106, 3 figs.
1967. "Eupera primei" sp. n. de la region del Rio Ucayali, Peru (Mollusca, Pelecypoda). *Atas do Simposio sobre a Biota Amazonica* 3 (Limnologia): 109–115, 3 figs.
- Kosuge, S.
1994. Revision of the genus *Meocardia* (Bivalvia Glossidae). *Bull. Inst. Malacol. Tokyo* 3(3): 36–49, pls. 13–18.
- Lea, I.
1834. Observations on the naiades; and descriptions of new species of that, and other families. *Trans. Am. Philos. Soc. n. ser.* 5(2): 23–119, 19 pls.
1838. Description of new freshwater and land shells. *Ibid.* 6(1): 1–154, 24 pls.
1840. Descriptions of new fresh water and land shells. *Proc. Am. Philos. Soc.* 1(13): 284–289.
1841. On fresh water and land shells. *Ibid.* 2(17): 30–35.
1842. Description of fresh water and land shells. *Trans. Am. Philos. Soc., n. ser.* 8(12): 163–250, pls. 5–27.
1843. Description of twelve new species of uniones. Privately Printed. 1 sheet.
1846. Description of new fresh water and land shells. *Trans. Am. Philos. Soc.* 9(10): 275–282, pls. 39–42.
1856. Descriptions of twenty-five new species of exotic uniones. *Proc. Acad. Nat. Sci. Philadelphia* 8(2): 92–95.
1858. Descriptions of exotic genera and species of the family Unionidae. *J. Acad. Nat. Sci. Philadelphia, ser. 2, 3(4)*: 289–321, pls. 21–33.
1859. Descriptions of twenty-one new species of exotic Unionidae. *Proc. Acad. Nat. Sci. Philadelphia* 11: 151–154.
1860. Descriptions of three new species of exotic Unionidae. *Ibid.* 12: 307–308.
1863. New exotic Unionidae. *J. Acad. Nat. Sci. Philadelphia, ser. 2, 5(4)*: 377–400, pls. 41–50.
1865. Description of six new species of western Asiatic Unionidae. *Proc. Acad. Nat. Sci. Philadelphia* 16: 285–286.
1866. Description of five new species of the genus *Unio*. *Ibid.* 18: 133.
1869. New Unionidae, Melanidae, etc., chiefly of the United States. *J. Acad. Nat. Sci. Philadelphia, ser. 2, 6(3)*: 249–302, pls. 29–45.
- Li, C. C.
1930. The Miocene and Recent Mollusca of Panama Bay. *Bull. Geol. Soc. China* 9(3): 249–296, 8 pls.
- Lyons, W. G.
1985. *Chaetopleura staphylophera* (Polyplacophora: Chaetopleuridae), a new species from the southeastern United States and Bahamas. *Nautilus* 99(2–3): 35–44, 30 figs.
- Marcus, E.
1953. Three Brazilian sand-Opisthobranchia. *Bol. Fac. Ciênc. Letr. Univ. São Paulo Zoologia* 18: 165–203, 49 figs.
1955. Opisthobranchia from Brazil. *Ibid.* 20: 89–200, 285 figs.
1957. On Opisthobranchia from Brazil (2). *J.*

- Linn. Soc. London 43(292): 390–486, 246 figs.
- 1958a (1956). Notes on Opisthobranchia. Bol. Inst. Oceanogr. 7(1–2): 31–78, 8 pls.
- 1958b. On western Atlantic opisthobranchiate gastropods. Am. Mus. Novitates 1906: 82 pp., 111 figs.
- Marcus, Ev.
- 1970. Opisthobranchs from northern Brazil. Bull. Mar. Sci. 20(4): 922–951, 49 figs.
 - 1977. An annotated checklist of the western Atlantic warm water opisthobranchs. J. Molluscan Stud., suppl. 4: 22 pp.
- Marcus, Ev., and E. Marcus
- 1956. On two sacoglossan slugs from Brazil. Am. Mus. Novitates 1796: 21 pp., 23 figs.
 - 1969. Opisthobranchian and lamellarian gastropods collected by the "Vema." Ibid. 2368: 33 pp., 39 figs.
 - 1970. Some gastropods from Madagascar and West Mexico. Malacologia 10(1): 181–223, 93 figs.
- Marincovich, L., Jr.
- 1973. Intertidal mollusks of Iquique, Chile. Los Angeles Cty. Mus. Nat. Hist. Sci. Bull. 16: 49 pp., 102 figs.
- Marshall, E.
- 1980. *Pulsellum salishorum* spec. nov., a new scaphopod from the Pacific Northwest. Veliger 23(2): 149–152, 6 figs.
- Marshall, W. B.
- 1926. New land and fresh-water mollusks from Central and South America. Proc. U.S. Natl. Mus. 69(12): 1–12, 3 pls.
- Maury, C. J.
- 1922. The Recent arcas of the Panamic Province. Palaeontol. Am. 1(4): 163–208, pls. 29–31.
- McMichael, D. F.
- 1958. The nature and origin of the New Zealand freshwater mussel fauna. Trans. R. Soc. New Zealand 85(3): 427–432.
- McMichael, D. F., and I. D. Hiscock
- 1958. A monograph of the freshwater mussels (Mollusca: Pelecypoda) of the Australian region. Aust. J. Mar. Freshwater Res. 9(3): 372–508, 19 pls., 18 text figs.
- Moore, D. R.
- 1977. Small species of Nuculidae (Bivalvia) from the tropical western Atlantic. Nautilus 91(4): 119–128, 7 figs.
- Morrison, J. P. E.
- 1971. Western Atlantic *Donax*. Proc. Biol. Soc. Washington 83(48): 545–568, 2 pls.
- Mulliner, D. K., and G. G. Sphon
- 1974. A new *Platydoris* (Gastropoda: Nudibranchia) from the Galápagos Islands.
- Trans. San Diego Soc. Nat. Hist. 17(15): 209–216, 5 figs.
- Oesch, R. D.
- 1984. Missouri naiades: a guide to the mussels of Missouri. Jefferson City, MO: Missouri Department of Conservation. 270 pp., illus.
- Olazarri, J., A. Mones, and E. Duarte
- 1972. Lista de los ejemplares-tipo depositados en el Museo Nacional de Historia Natural de Montevideo, Uruguay. Comun. Zool. Mus. Hist. Nat. Montevideo 10(133): 8 pp.
- Oldroyd, I. S.
- 1918. A new species of *Cuspidaria* from Monterey. Nautilus 32(1): 28.
 - 1920. New species of West Coast shells. Ibid. 33(4): 135–136, pl. 4, 6 text figs.
 - 1924. The marine shells of the West Coast of North America. Stanford Univ. Publ. Geol. Sci. 1(1): 248 pp., 57 pls.
- Olsson, A. A.
- 1961. Mollusks of the tropical eastern Pacific particularly from the southern half of the Panamic-Pacific faunal province (Panama to Peru). Ithaca, NY: Paleontological Research Institution. 574 pp., 86 pls.
- Pilsbry, H. A.
- 1892–93. Manual of conchology. ser. 1, vol. 14. Polyplacophora, (Chitons.) Lepidopleuridae, Ischnochitonidae, Chitonidae, Mopaliidae. Philadelphia, PA: Conchological Section, Academy Natural Sciences. 350 pp., 68 pls.
 - 1893–95. Manual of conchology. ser. 1, vol. 15. Polyplacophora, (Chitons.) Acanthochitonidae, Cryptoplacidae and Appendix. Tectibrachiata. Philadelphia, PA: Conchological Section, Academy Natural Sciences. 436 pp., 61 pls.
 - 1931. The Miocene and recent Mollusca of Panama Bay. Proc. Acad. Nat. Sci. Philadelphia 83: 427–440, pl. 41, 5 text figs.
- Pilsbry, H. A., and J. Bequaert
- 1927. The aquatic mollusks of the Belgian Congo, with a geographical and ecological account of Congo malacology. Bull. Am. Mus. Nat. Hist. 53(2): 69–602, pls. 10–77, 94 text figs.
- Pilsbry, H. A., and B. Sharp
- 1897–98. Manual of conchology. ser. 1, vol. 17. Scaphopoda and Aplacophora. Index to genera and subgenera, volumes II to XVII. Philadelphia, PA: Conchological Section, Academy Natural Sciences. 348 pp., 48 pls.
- Powell, A. W. B.

1933. New species of marine Mollusca from the subantarctic islands of New Zealand. Proc. Malacol. Soc. London 20(5): 232–236, pl. 20, 4 text figs.
1958. Mollusca of the Kermadec Islands. Part I. Rec. Auckland Inst. Mus. 5(1–2): 65–85, pls. 9–11, text fig. A.
1979. New Zealand Mollusca : marine, land and freshwater shells. Auckland, New Zealand: William Collins. 500 pp., 78 pls., 120 text figs.
- Prime, T.
1860. Description of two new species of the genus *Batissa*, with notes on that genus. Ann. Lyc. Nat. Hist. New York 7: 112–115.
- 1861a. Descriptions of new species of *Cyrena*, *Corbicula* and *Sphaerium*. Proc. Acad. Nat. Sci. Philadelphia. 13: 125–128.
- 1861b. Diagnoses d'espèces nouvelles. J. Conchyliol. 9(4): 354–356.
- 1862a. Description of a new species of Mollusca of the genus *Venus*. Ann. Lyc. Nat. Hist. New York 7: 482–483, text fig.
- 1862b. Description d'espèces nouvelles des genres *Glaucome*, *Cyrena*, *Batissa* et *Corbicula*. J. Conchyliol. 10(4): 383–390, pl. 13, figs. 1–2; pl. 14.
- Redfield, J. H.
1848. Description of some new species of shells. Ann. Lyc. Nat. Hist. New York 4: 163–168, pls. 10, 11.
- Richards, M. C., and W. E. Old, Jr.
1969. A catalogue of molluscan type specimens in the Department of Living Invertebrates, the American Museum of Natural History, New York, New York, U.S.A. New York: Dep. of Living Invertebrates, Am. Mus. Nat. Hist. 147 pp.
- Rios, E. C.
1994. Seashells of Brazil. 2nd ed. Mus. Oceanografico da FURG. 368 pp., 113 pls.
- Rombouts, A.
1991. Guidebook to pecten shells—Recent Pectinidae and Propeamussiidae of the world. The Netherlands: Universal Book Services. 157 pp., 29 pls.
- Rosenberg, G., A. E. Bogan, and E. E. Spamer
1990. *Coelatura* Conrad, 1853, *Caelatura* Conrad, 1865 and *Coelatura* Pfeiffer, 1877 (Mollusca): a tale of two diphthongs. *Nautilus* 104(1): 29–32.
- Rudman, W. B.
1984. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: a review of the genera. Zool. J. Linn. Soc. 81: 115–273, 102 figs.
- Salvini-Plawen, L. von
1972. Die Caudofoveata des Mittelmeeres und das genus *Scutopus* (Mollusca, Aculifera). Fifth European Mar. Biol. Symp.: 27–51, 31 figs.
1977. Caudofoveata (Mollusca) des Forschungsprojektes Polymede. Bull. Mus. Nat. Hist. Nat., Paris, 3e ser. 447 (Zoologie 310): 413–421, 6 figs.
1986. On certain Caudofoveata from the 'Vema' Expedition. Abstracts, Ninth Int. Malacol. Congr., Edinburgh, 1986: 76.
1992. On certain Caudofoveata from the VEMA-Expedition. Proc. Ninth Int. Malacol. Congr., Edinburgh, 1986: 317–333, 25 figs.
- Scarabino, V.
- 1986a. Nuevos taxa abisales de la clase Scaphopoda (Mollusca). Comun. Zool. Mus. Hist. Nat. Montevideo 11(155): 20 pp., 24 figs.
- 1986b. Systematics of Scaphopoda (Mollusca), I. Three new bathyal and abyssal taxa of the order Gadilida from the South and North Atlantic Ocean. Ibid. 11(161): 16 pp., 20 figs.
- Scott, P. H., F. G. Hochberg, and B. Roth
1990. Catalog of Recent and fossil molluscan types in the Santa Barbara Museum of Natural History. I. Caudofoveata, Polyplacophora, Bivalvia, Scaphopoda, and Cephalopoda. Veliger 33 (suppl. 1): 27 pp., 3 figs.
- Scudder, N. P.
1885. The published writings of Isaac Lea, LL. D. Bull. U.S. Natl. Mus. 23: lix + 278 pp.
- Shimek, R. L.
1989. Shell morphometrics and systematics: a revision of the slender, shallow-water *Cadulus* of the northeastern Pacific (Scaphopoda: Gadilida). Veliger 32(3): 233–246, 5 figs.
- Simpson, C. T.
1900. Synopsis of the naiades, or pearly freshwater mussels. Proc. U.S. Natl. Mus. 22(1205): 501–1044.
- Smith, B. J.
1992. Non-marine Mollusca. In Houston, W. W. K. (ed.), Zoological catalogue of Australia. Canberra: AGPS vol. 8: 405 pp.
- Sphon, G. G.
1971. Type specimens of Recent mollusks in the Los Angeles County Museum of Natural History. Los Angeles Cty Mus. Contrib. Sci. 213: 37 pp., 21 figs.
- Sterki, V.

1896. Descriptions of new pisidia. *Nautilus* 10(2): 20–21.
1906. *Sphaerium hendersoni* n. sp. *Ibid.* 20(6): 69–70.
- Turgeon, D. D., A. E. Bogan, E. V. Coan, W. K. Emerson, W. G. Lyons, W. L. Pratt, C. F. E. Roper, A. Scheltema, F. G. Thompson, and J. D. Williams
1988. Common and scientific names of aquatic invertebrates from the United States and Canada: mollusks. *Am. Fish. Soc. Spec. Publ.* 16: 277 pp., 12 pls.
- Turner, R. D.
1955. Scaphopods of the *Atlantis* dredgings in the western Atlantic with a catalogue of the scaphopod types in the Museum of Comparative Zoology. *Pap. Mar. Biol. and Oceanogr., Deep-Sea Res.* 3 (suppl.): 309–320.
- Tursch, B., and J. Pierret
1964. New species of mollusks from the coast of Brazil. *Veliger* 7(1): 35–37, 5 figs.
- Usticke, G. W. N.
1959. A check list of the marine shells of St. Croix, U.S. Virgin Islands, with random annotations. Burlington, VT: Lane Press. 90 pp., 4 pls.
1969. A supplementary listing of new shells (illustrated) to be added to the check list of the marine shells of St. Croix. Privately published. 32 pp., 6 pls.
1971. A supplementary listing of new shells (illustrated), revised edition, to be added to the check list of the marine shells of St. Croix. Privately published. 32 pp., 6 pls.
- MS. [A check list of the marine shells of St. Croix and the Lesser Antilles]. MS on file, American Museum Natural History, New York. 276 pp. (Gastropoda; pp. 202–222 missing) + 105 pp. (Bivalvia, Polyplacophora, Scaphopoda and Cephalopoda).
- Vaught, K. C.
1989. A classification of the living Mollusca. Melbourne, FL: American Malacologists, Inc. 195 pp.
- Verrill, A. E.
1884. Second catalogue of Mollusca, recently added to the fauna of the New England coast and the adjacent parts of the Atlantic, consisting mostly of deep sea species, with notes on others previously recorded. *Trans. Conn. Acad.* 6(1): 139–294, pls. 28–32.
- Waller, T. R.
1986. A new genus and species of scallop (Bivalvia: Pectinidae) from off Somalia and the definition of a new tribe Decatopectinini. *Nautilus* 100(2): 39–46, 13 figs.
- Willan, R. C.
1993. Taxonomic revision of the family Psammobiidae (Bivalvia: Tellinoidea) in the Australian and New Zealand region. *Rec. Aust. Mus., suppl.* 18: 132 pp., 416 figs.
- Woodward, S. P.
1859. On some new freshwater shells from Central Africa. *Proc. Zool. Soc. London:* 348–350, pl. 47.

INDEX, BY CLASS AND SUBCLASS

APLACOPHORA

- acutargatus*, *Falcidens* (*Lepoderma*), 4
chilensis, *Scutopus*, 6
elongatus, *Limifosssor*, 5
Falcidens, 4, 5
hexadentatus, *Limifosssor*, 5
holopeltatus, *Limifosssor*, 6
Lepoderma, 4, 5
Limifosssor, 5, 6
limifosssorides, *Falcidens*, 4
Lonchoderma, 6
longisquamosum, *Prochaetoderma* (*Lonchoderma*), 6

- megaradulatus*, *Scutopus*, 6
nontargatus, *Falcidens* (*Lepoderma*), 5
procerus, *Falcidens*, 5
Prochaetoderma, 6
profundus, *Limifosssor*, 6
Psilodens, 5
Scutopus, 6
targatus, *Falcidens* (*Lepoderma*), 5

POLYPLACOPHORA

- albida*, *Plaxiphora* (*Plaxiphora*), 7
allynsmithi, *Lepidozona*, 7

- boogii*, *Stenoplax* (*Stenoplax*), 7
Callistochiton, 6
Chaetopleura, 6
Chiton, 6, 7
dubium, *Ischnochiton*, 7
excavata, *Plaxiphora*, 7
guadalupensis, *Lepidozona*, 7
Ischnochiton, 7
leei, *Callistochiton*, 6
Lepidozona, 7
macleani, *Ischnochiton*, 7
Plaxiphora, 7
rubrocarinata, *Chiton viridis* var., 6
staphylophera, *Chaetopleura*, 6

Stenoplax, 7
viridis, Chiton, 7

GASTROPODA

OPISTHOBRANCHIA

- Acteon*, 7
- Alderia*, 8
- Antiopella*, 8
- apefiae*, *Hallaxa*, 11
- Atys*, 8
- auritulus*, *Favorinus*, 11
- Baeolidia*, 8
- benteva*, *Baeolidia*, 8
- Berghia*, 8
- Berthella*, 10
- bovena*, *Doris*, 10
- byga*, *Rostanga*, 10
- Cadlina*, 8, 10
- canguzua*, *Elysia*, 11
- Capellinia*, 9
- caribaea*, *Atys*, 8
- carolynae*, *Platydoris*, 10
- Catriona*, 9, 10
- cauze*, *Elysia*, 11
- Chelidonura*, 10
- chica*, *Doto*, 9
- chilla*, *Philine thurmanni*, 9
- coenda*, *Facelina*, 9
- comis*, *Janolus*, 11
- conicla*, *Capellinia*, 9
- coniclus*, *Eubranchus*, 9
- conyna*, *Polycerella*, 11
- corpulenta*, *Atys caribaea* var., 8
- cuica*, *Pluscula*, 11
- Cuthona*, 9
- Discodoris*, 10
- diuda*, *Thordisa*, 11
- divae*, *Piseinotucus*, 11
- Doris*, 10
- Doto*, 9, 11
- eloiseae*, *Acteon*, 7
- Elysia*, 9, 11
- engeli*, *Phyllaplysia*, 11
- erssus*, *Stiliger* (*Stiliger*), 10
- Eubranchus*, 9
- evelinae*, *Cadlina*, 8
- evelinae*, *Chelidonura*, 10
- evelinae*, *Ganitus*, 11
- evelinae*, *Okenia*, 11
- Facelina*, 9

- Favorinus*, 11
- Ganitus*, 11
- Glossodoris*, 11
- Hallaxa*, 11
- Hancockia*, 11
- Hypselodoris*, 9
- ilo*, *Siraius*, 11
- isa*, *Noumeaella*, 9
- Janolus*, 8, 11
- kirsteueri*, *Smaragdinella*, 10
- lineata*, *Atys*, 8
- mandrewii*, *Atys*, 8
- maringa*, *Trapania*, 11
- modesta*, *Alderia*, 8
- mucloc*, *Antiopella*, 8
- neona*, *Glossodoris*, 11
- Noumeaella*, 9
- odhneri*, *Polycera*, 11
- Okenia*, 11
- perca*, *Catriona*, 9
- Philine*, 9, 10
- Phyllaplysia*, 11
- Piseinotucus*, 11
- pita*, *Doto*, 11
- Platydoris*, 10
- Pluscula*, 11
- Polycera*, 11
- Polycerella*, 11
- Punctacteon*, 8
- pusae*, *Discodoris*, 10
- regina*, *Hypselodoris*, 9
- Retusa*, 10
- riiseanus*, *Atys*, 8
- Rostanga*, 10
- rumia*, *Cadlina*, 10
- ryrca*, *Hancockia*, 11
- serca*, *Elysia*, 11
- Siraius*, 11
- Smaragdinella*, 10
- sosa*, *Retusa*, 10
- Stiliger*, 10
- talis*, *Stiliger* (*Stiliger*), 10
- Taringa*, 11
- telopia*, *Taringa*, 11
- Thordisa*, 11
- thurmanni*, *Philine thurmanni*, 10
- tina*, *Catriona*, 10
- Trapania*, 11
- tupala*, *Berthella*, 10
- Tyrinna*, 9
- uda*, *Alderia*, 8
- umia*, *Doto fragilis*, 9

uva, *Doto*, 10
vreelandae, *Elysia*, 9

BIVALVIA

- Aequipecten*, 11
- Aequiyoldia*, 19
- africanus*, *Unio*, 24
- Agnomyax*, 12
- Alathyria*, 27
- aldridgei*, *Solen viridis* var., 22
- aletes*, *Pitar* (*Hyphanotosoma*), 21
- altirostris*, *Corbula*, 14
- Amphidesma*, 22
- Anadara*, 12
- Angulus*, 23
- Anodonta*, 11, 28
- Anodontites*, 11, 12
- Anodontoides*, 12
- Arca*, 12
- arenica*, *Tellina* (*Moerella*), 23
- Arenomya*, 19
- Argina*, 12
- Argopecten*, 20
- Asaphis*, 12
- askoyana*, *Tellina* (*Eurytellina*)
eburnea, 23
- Aspatharia*, 3, 12
- auklandica*, *Legrandina*, 17
- auklandica*, *Perrierina* (*Legrandina*), 18
- aurora*, *Semele*, 22
- Basterotia*, 12
- Batissa*, 3, 13
- bibsae*, *Bornia*, 13
- bolleyi*, *Pecten* (*Leptopecten*)
velero, 21
- Bornia*, 13
- bourguignati*, *Coelatura* (*Laevisrostris*), 13
- boydianus*, *Unio*, 24
- Brachidontes*, 13
- brevidentata*, *Cyrenoida rosea*, 15
- brevifrons*, *Lunarca*, 12
- bucaruana*, *Argina brevifrons*, 12
- burtoni*, *Unio*, 24
- Caelatura*, 13
- Cafferia*, 24, 28
- caffra*, *Cafferia*, 24, 28
- calcicola*, *Nucula*, 19
- callosus*, *Unio*, 3, 24
- Calyptogena*, 14

- cancellata*, *Phacoides*, 28
Cardium, 14, 28
carpenteri, *Tellina (Angulus)*, 23
centrifuga, *Lucina*, 21
Chamberlainia, 25
chapini, *Aspatharia*, 3
Chione, 14
Chlamydella, 14
Chlamys, 3, 20, 28
circularis, *Argopecten*, 20
clarionense, *Cardium (Laevicardium)*, 14
Coelatura, 13, 14, 24
coelestis, *Iridina*, 3, 17
cognata, *Temnoconcha*, 23
Condylonucula, 14
congo, *Iphigenia*, 17
contradens, *Uniandra*, 25, 27, 28
Corbicula, 28
Corbula, 14, 15
corrugata, *Parreysia*, 26
Corymbichlamys, 11
Costelloleda, 20
craneana, *Semele*, 22
cranmerorum, *Somalipecten*, 22
Crassatellites, 15
Crassinella, 15
crebristriata, *Mondocondyla*, 18
crispata, *Scabies*, 27
crockeri, *Solen*, 22
cruzensis, *Montacuta*, 18
cubitus, *Ischadium*, 13
Cucumerunio, 24
cucumoides, *Unio*, 24
cumingi, *Glaucome*, 17
Cuspidaria, 15
Cyrena, 15, 28
Cyrenoida, 15
denticulata, *Petricola (Rupellaria)*, 28
diegensis, *Limopsis (Limopsis)*, 18
doellojuradoi, *Eupera*, 16
Donax, 15
dorotheae, *Donax*, 15
dubia, *Mutela (Mutela)*, 17, 19
eburnea, *Tellina (Eurytellina)*, 23
Echyridella, 26
Ectenagena, 14
Egeria, 16
eightsi, *Nucula*, 19
eightsi, *Yoldia (Aequioldia)*, 19
elegans, *Coelatura (Zairia)*, 14
elegans, *Harvella*, 21
Elliptio, 16, 25
Elliptoideus, 27
emarginata, *Solenaia*, 19
emarginatus, *Mycetopus*, 19
Ensidens, 27
Ensis, 16
erosa, *Corbicula*, 28
Eucrassatella, 15
Eupera, 16
Eurytellina, 23
evitatus, *Unio*, 28
eximius, *Unio*, 24
exustus, *Brachidontes*, 13
fallax, *Pisidium*, 21
ferrussacianus, *Anodontoides*, 12
fililectus, *Pecten*, 20
filitextus, *Pecten*, 20
flava, *Aspatharia*, 12
fonsecana, *Mactra (Micromactra)*, 18
fortis, *Cyrena*, 28
frizzelli, *Pitar (Lamelliconcha)*, 21
Galatea, 16
garambae, *Mutela*, 18
Gemma, 29
gerrardi, *Transenella*, 24
gibbosa, *Eucrassatella*, 15
gigantea, *Batissa*, 3, 13
gigas, *Mactra*, 28
Glaucome, 17, 28
Glycymeris, 17, 28
glypta, *Corbula*, 15
gordoni, *Glycymeris*, 17, 28
gracilis, *Corbicula*, 28
Grandidieria, 24
gravidus, *Unio*, 25, 27
guatulcoensis, *Chione (Chione)*, 14
hainesiana, *Chamberlainia*, 25
hainesianus, *Unio*, 25
Harvella, 21
hastata, *Chlamys*, 20
hendersoni, *Sphaerium*, 23
Hertellina, 23
hopkinsi, *Penitella*, 21
housei, *Unio*, 25
humilis, *Unio*, 25
Hyphantosoma, 21
Hyridella, 26
Hyriopsis, 25, 26
incantata, *Chlamys (Chlamys)*, 3, 28
indentus, *Pecten latiauritus (sic)*, 20
Indonaia, 25, 26
ingallsianus, *Ensidens*, 27
inornatus, *Unio*, 25
inoscularis, *Pseudodon inoscularis*, 18
interflucta, *Nucula (Nucula)*, 19
intermedius, *Donax striatus* var., 15, 16
Iphigenia, 17
iredalei, *Aequipecten (Corymbichlamys)*, 11
iredalei, *Salaputium*, 21
Iridina, 3, 17
iris, *Villosa*, 26
irisans, *Anodontites*, 12
Ischadium, 13
Isocardia, 17
jamesi, *Semele*, 22
japonica, *Mya*, 19
jayana, *Glaucome*, 17
jayensis, *Cyrena*, 15
jayensis, *Unio*, 25
Laevicardium, 14
Laevirostris, 13
Lamelliconcha, 21
Lampsilis, 24
langi, *Egeria tenuicula*, 16
langi, *Mutela*, 19
lata, *Crassinella*, 15
lavalleanus, *Brachidontes*, 13
Legrandina, 17, 18
Leguminaia, 18
lemchei, *Chlamydella favus*, 14
leotaudi, *Anodontites (Anodontites) patagonicus*, 12
Leptopecten, 20, 21
liana, *Phacoides (Lucinisca)*, 21
Limnoscapha, 25, 26
Limopsis, 18
Lioberus, 28
Lucina, 21
Lucinisca, 21
Lunarca, 12
lutens, *Unio*, 26
luteus, *Unio*, 26
luticola, *Sphenia*, 15
lутulentus, *Unio*, 26
lydigiana, *Corbicula*, 28
macmichaeli, *Elliptio*, 16
Macoma, 18
Mactra, 18, 28
magnifica, *Calyptogena (Ectenagena)*, 14

- manhattensis*, *Venus (Gemma)*, 29
mardinensis, *Mondocondylaea*, 18
marella, *Nuculana (Costelloleda)*,
 20
maxima, *Raeta*, 21
maya, *Condylonucula*, 14
mazatlanica, *Arca (Anadara)*, 12
mcmichaeli, *Elliptio*, 16
Meiocardia, 17
menziesi, *Hyridella (Echyridella)*,
 26
Mesafra, 13
mesafricana, *Caelatura*, 13
mexicana, *Nephronais (Sphenon-*
 aia), 24
Mexicardia, 14
Micromactra, 18
minor, *Chione*, 14
modesta, *Semele*, 22
Moerella, 23
Mondocondylaea, 18
Montacuta, 18
montezuma, *Anodonta*, 11
Mulinia, 15
Mutela, 17–19
Mya, 19
Mycetopoda, 19
Mycetopus, 19
myersiana, *Hyriopsis (Limnosca-*
 pha), 25, 26
myersianus, *Unio*, 26
myrae, *Ensis*, 16
nana, *Asaphis*, 12
nana, *Cuspidaria (Tropidomya)*,
 15
Nephronais, 24
nicoiana, *Tellina (Scissula)*, 23
notata, *Corbicula*, 28
novaehollandiae, *Cucumerunio*, 24
novi-eboraci, *Unio*, 26
nucleus, *Unio*, 26
Nucula, 19, 20
Nuculana, 20
Nuculopsis, 20
nux, *Egeria*, 16
occata, *Indonaia*, 26
occatus, *Unio*, 26
olivacea, *Psammobia*, 28
Pacipecten, 20, 21
pallida, *Mulinia*, 15
panamanensis, *Tellina*, 23
Parabornia, 20
Parreysia, 26
parvulum, *Cardium*, 14
pavonia, *Anodonta*, 11
Pecten, 20, 21, 28
Penitella, 21
Perrierina, 18
pertexta, *Alathyria pertexta*, 27
peruviana, *Venerupis*, 28
Petricola, 28
Phacoides, 21, 28
phaselus, *Unio*, 26
Physunio, 25, 27
Pisidium, 21
Pitar, 21
praetenuis, *Mutela garambae*, 19
primei, *Eupera*, 16
procерум, *Trachycardium (Mexi-*
 cardia), 14
Psammacoma, 18
Psammobia, 28
Pseudodon, 18
pseudorecurvus, *Brachidontes*, 13
pugetensis, *Pecten islandicus*, 20
pustula, *Basterotia*, 12
radiata, *Lampsilis (Lampsilis)*, 24
Raeta, 21
rajahensis, *Unio*, 28
richardsoni, *Penitella*, 21
rosaceus, *Brachidontes exustus*
 var., 13
rotula, *Caelatura (Laevirostris?)*,
 13
rudis, *Crassatellites*, 15
rugifera, *Aspatharia (Aspatharia)*
 rugifera, 12
Rupellaria, 28
rustica, *Uniandra contradens*, 27
rusticus, *Unio*, 27
sagittarius, *Unio*, 27
Salaputium, 21
salvadorica, *Volsella (Volsella)*, 28
salvadoricus, *Lioberus*, 28
samarangiae, *Meiocardia*, 17
Scabies, 26, 27
schencki, *Nucula (Nuculopsis)*, 20
Scissula, 23
scobinata, *Scabies*, 27
scobinatus, *Unio*, 27
Semele, 22
senegalensis, *Anodonta*, 28
Septifer, 22
setosum, *Cardium*, 28
shuttleworthi, *Unio*, 27
siliqua, *Macoma*, 18
skoglundae, *Crassinella*, 15
sloataianus, *Unio*, 27
Solen, 22
Solenaia, 19
Somalipecten, 22, 23
spectri, *Macoma (Psammacoma)*
 panamensis, 18
Sphaerium, 23
Sphenia, 15
Sphenonaias, 24
splendens, *Pecten latiauritus (sic)*,
 20
squillina, *Parabornia*, 20
stanleyvillensis, *Caelatura*, 13
striatum, *Sphaerium*, 23
Strophitus, 11
subcylindracea, *Anodonta*, 11
sumatrensis, *Glaconome*, 28
superbus, *Physunio (Physunio)*, 25
superbus, *Unio*, 27
Tellina, 23
Tellinella, 23
Temnoconcha, 23
Tenuicorbula, 15
tenuilineatus, *Tellina*, 23
tenuis, *Corbula (Tenuicorbula)*, 15
tetragona, *Isocardia*, 17
texasanus, *Donax*, 15
tobagoensis, *Glycymeris gordoni*,
 17
Trachycardium, 14
Transenella, 24
trigonus, *Anodontites (Anodon-*
 tites), 11
tropicalis, *Ensis*, 16
Tropidomya, 15
tumbezensis, *Leptopecten (Paci-*
 pecten), 20, 21
tumidula, *Uniandra contradens*,
 25, 28
tumidulus, *Unio*, 27
undulatus, *Strophitus*, 11
Uniandra, 25, 27, 28
Unio, 3, 24–28
Venerupis, 28
Venus, 29
verreauianus, *Unio*, 28
Villosa, 26
violacea, *Corbicula*, 28
Volsella, 28
wheatleyi, *Leguminaia*, 18
yessoensis, *Pecten*, 28
Yoldia, 19

zacae, *Tellina (Tellinella)*, 23
Zairia, 14
zeteiki, *Septifer*, 22

SCAPHOPODA

aberrans, *Cadulus*, 30
acus, *Cadulus*, 29
albicomatus, *Cadulus*, 29
americanum, *Dentalium*, 32
Annulipulsellum, 29
Antalis, 32
arctus, *Cadulus (Platyschides)*, 29
atlanticus, *Cadulus (Gadila)*, 29
bartletti, *Dentalium (Antalis)*, 32
Cadulus, 29–31
californicus, *Cadulus (Gadila)*, 29
candidum, *Fissidentalium*, 33
ceratum, *Dentalium (Antalis)*, 32
Costentalina, 32
dalli, *Cadulus (Polyschides)*, 29
Dentalium, 32, 33
didymum, *Episiphon*, 32
duartei, *Wemersoniella*, 33
Episiphon, 32, 33
euzkadii, *Annulipulsellum*, 29

Fissidentalium, 32
flavum, *Dentalium (Antalis) ceratum*, 32
fusiformis, *Cadulus (Gadila)*, 29
Gadila, 29–31
halius, *Cadulus (Cadulus)*, 30
isaotakii, *Siphonodentalium*, 33
japonicum, *Siphonodentalium*, 33
johnsoni, *Dentalium (Episiphon)*, 32
macleani, *Cadulus (Platyschides)*, 30
mayori, *Cadulus (Gadila)*, 30
miamiensis, *Cadulus (Platyschides)*, 30
nedallisoni, *Cadulus (Polyschides)*, 30
nitidus, *Cadulus (Platyschides)*, 30
okudai, *Siphonodentalium*, 33
panamensis, *Cadulus (Gadila)*, 30
pelliceri, *Dentalium (Episiphon) sowerbyi*, 33
perpusillus, *Cadulus*, 30
platensis, *Cadulus (Cadulus)*, 31
Platyschides, 29–31
Polyschides, 29–31
providensis, *Cadulus (Platyschides)*, 31
Pulsellum, 33
rebeccaense, *Dentalium (Dentalium)*, 32
regularis, *Cadulus (Gadila)*, 31
salishorum, *Pulsellum*, 33
simpsoni, *Cadulus (Platyschides)*, 31
Siphonodentalium, 33
solidum, *Dentalium*, 32
sowerbyi, *Episiphon*, 33
striatus, *Cadulus (Gadila)*, 31
Striocadulus, 29
subagile, *Dentalium (Antalis) agile*, 32
tetraschistus, *Polyschides*, 31
tetrodon, *Cadulus (Polyschides)*, 31
transitorius, *Cadulus (Cadulus transitorius)*, 31
vemae, *Costentalina*, 32
verrilli, *Cadulus (Gadila)*, 31
verrilli, *Dentalium (Fissidentalium) meridionale*, 32
Wemersoniella, 33

Recent issues of the *Novitates* may be purchased from the Museum. Lists of back issues of the *Novitates*, *Bulletin*, and *Anthropological Papers* published during the last five years are available free of charge. Address orders to: American Museum of Natural History Library, Department D, Central Park West at 79th St., New York, N.Y. 10024. TEL: (212) 769-5545. FAX: (212) 769-5009. E-MAIL: scipubs@amnh.org