American Museum Novitates

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY CENTRAL PARK WEST AT 79TH STREET, NEW YORK 24, N.Y.

NUMBER 1700

DECEMBER 20, 1954

The Lectotype of *Megathymus aryxna* Dyar (Lepidoptera, Megathymidae)

By Ernest Layton Bell¹ and Cyril Franklin dos Passos²

Despite the excellent revision of the Megathymidae by Barnes and McDunnough (1912), there has existed considerable confusion for a number of years concerning the nomenclatorial status of certain names in that family. This uncertainty was caused in part by the fact that one name had been proposed from what appeared to be a mixed series, and it was dubious whether the type had in fact been fixed, and if so, by whom. It is to clarify this situation and put to rest the confusion now existing that the present paper is offered. No new name is proposed, and the taxonomic position of the names discussed is not treated, although our studies have led us to definite conclusions respecting some of them.

Before the case of *Megathymus aryxna* Dyar (1905), the subject of this paper, is considered, mention should be made of one earlier name proposed in that genus. This is *neumoegeni* Edwards (1882). This species was described from a series of three specimens taken by Jacob Doll near Prescott, Arizona. The types are in the United States National Museum. A fourth specimen was taken by Doll, but it is not considered a type, although so treated by Strecker (1900). That specimen is in the Chicago Natural History Museum.

Coming then to aryxna we find that the first element of uncertainty

¹ Research Associate, Department of Insects and Spiders, the American Museum of Natural History.

² Research Associate, Department of Insects and Spiders, the American Museum of Natural History; Research Associate, Section of Insects and Spiders, Carnegie Museum.

has been caused by differences of opinion as to what constitutes the type series, and which specimen is the lectotype, because Dyar did not designate a holotype in the original description. Since that description is brief, it seems best to quote it in full:

"M. aryxna, new species.

"This is the form figured in the Biologia Cent.-Am. Lep. Het., III, pl. 69, figs. 3 and 4. It differs from *neumoegeni* in having the fulvous markings considerably reduced, the outer band being broken into spots. I have ten specimens from Arizona from Dr. Barnes and Mr. Poling."

Three different conclusions have been drawn from this description as to what constitutes the type series. First, that it consists of the two specimens figured as neumoegeni in the "Biologia" (figs. 3 and 4), the first of which (fig. 3) was subsequently named drucei by Skinner (1911). Second, that the type series consists only of the 10 specimens actually before Dyar when he drew up the original description. These specimens are in the United States National Museum. Third, that it consists of all 12 specimens, i.e., the two figured in the "Biologia" and the 10 before Dyar. The present authors do not believe that it is necessary to decide this interesting question, because they are of opinion that under the "Règles" the lectotype must be selected from the two specimens figured in the "Biologia." But before proceeding to that question, we shall follow briefly the chronological development of the problem.

As noted above, Skinner in 1911 proposed the name *drucei* for figure 3 in the "Biologia." This certainly eliminated that specimen from consideration, if not under the "Règles" as then existing it had that effect by analogy where the type series consists of two species under paragraph 73.B.h. of the Banks and Caudell Code (1912) which was at that time followed by entomologists when the "Règles" were silent. Therefore, if the type series of *aryxna* consisted only of the two specimens figured in the "Biologia," the remaining specimen by elimination would be the lectotype of *aryxna*. If the type series consisted only of the 10 specimens before Dyar, the result would be different, and the same could be true if it consisted originally of all 12 specimens.

In the aforementioned revision of Megathymidae Barnes and McDunnough partially considered this problem and came correctly to the conclusion that Dyar's 10 specimens consisted of two distinct species, six of one species and four of the other, that the six specimens were the same as those figured in the "Biologia" as neumoegeni, but that they could not hold validly the name, and that Dyar should designate one of the remaining four specimens as the type (lectotype) of aryxna, although, as Barnes and McDunnough admitted, those specimens did not agree very well with

the original description, Dyar having believed apparently that they were females and the other six males. Barnes and McDunnough did not designate any type (lectotype) of aryxna, but at or about the time of the publication of their paper Dyar wrote a holotype label as suggested by them and affixed it to one of the four specimens, but neither he nor anyone else ever published that fact, and it is believed confidently that his action was totally ineffective for that reason and also because none of the 10 specimens was eligible for selection as the holotype.

Skinner and Williams (1924) in their work on the male genitalia of the Hesperiidae of North America considered the problem and fixed definitely the type of aryxna as figure 4 of the "Biologia." This action would have been effective and confirmative of Skinner's earlier action in 1911 when he named figure 3 as drucei if Dyar's labeling of one specimen was ineffective, and would have sunk aryxna as a synonym of neumoegeni, assuming figure 4 to be that species.

It is quite clear from Article 31 of the "Règles," as amended by the Fourteenth International Congress of Zoology at Copenhagen (Hemming, 1953a), that where a new name (aryxna in this case) is published partly as a substitute name (for the "Biologia" figures of neumoegeni) and partly upon specimens before the author (Dyar's 10 specimens) when he published the name, the name is to adhere to the species, whatever it may be (neumoegeni in this case), in respect to which it was published as a substitute, the material which the author had before him when he published his substitute name being only subjectively, and therefore possibly erroneously, identified by him with the species that he renamed (see Hemming, 1953b, MS). The action at Copenhagen was taken as a result of a decision at the Thirteenth International Congress of Zoology held in 1948 at Paris (Hemming, 1950), and based upon documents submitted to the Copenhagen meeting (Hemming, 1953b). Therefore, when we find that Dyar stated "This [aryxna] is the form figured in the Biologia Cent.-Am. Lep. Het., III, pl. 69, figs. 3 and 4. . . . " it is evident that he intended aryxna as a substitute name for the specimens figured as neumoegeni, and that those specimens must be the syntypes from which the lectotype would have to be designated.

The only question left for consideration is which of the two specimens figured was first selected as the lectotype of aryxna. As explained above, Skinner in 1911 made the specimen figured in the "Biologia" as figure 3 the holotype of drucei. Consequently, the specimen, figure 4, automatically became the lectotype of aryxna, that being the only remaining syntype of aryxna after the removal to another species (drucei) of the only other syntype. Under one of the Paris decisions (Hemming, 1950), the ruling

in regard to the type species of genera given in the Commission's Opinion 6 was written into Article 31, which refers to type specimens of species. It is true that at Copenhagen the application to type specimens of the ruling given in Opinion 6 was repealed by the deletion of that provision from Article 31, but at the same time it protected the position where lectotypes in the past had been determined by this provision.

Our conclusion is, therefore, that the lectotype of aryxna is the specimen figured in the Lepidoptera Heterocera section of the "Biologia Centrali-Americana" as figure 4 on plate 69 of volume 3. The specimen is in the British Museum (Natural History). This result is especially fortunate in that it is confirmatory of the works of Skinner, and Skinner and Williams, and permits the recognition of Megathymus evansi Freeman (1950) as a valid name, it being the same species as the four specimens before Dyar. The holotype of evansi is in the American Museum of Natural History.

We desire to express our appreciation to Brigadier W. H. Evans, C.S.I., C.I.E., D.S.O., Honorary Associate, British Museum (Natural History), and Mr. Francis Hemming, C.M.G., C.B.E., of London, England, for giving us their views respecting this nomenclatorial problem, with which we find ourselves in full accord, and to Mr. R. L. Wenzel, Curator, Division of Insects, Chicago Natural History Museum, for lending us Strecker's "type" of M. neumoegeni for dissection.

BIBLIOGRAPHY

BANKS, NATHAN, AND ANDREW NELSON CAUDELL

1912. The entomological code. A code of nomenclature for use in entomology. Washington, D. C., Judd and Detweiler, Inc., 32 pp.

BARNES, WILLIAM, AND JAMES HALLIDAY McDUNNOUGH

1912. Revision of the Megathymidae. Contributions to the natural history of the Lepidoptera of North America. Decatur, Illinois, the Review Press, vol. 1, no. 3, 56 pp., 1 fig., 6 pls.

DRUCE, HERBERT

1881–1900. Insecta. Lepidoptera-Heterocera. *In* Godman, Frederick Du-Cane, and Osbert Salvin (eds.), Biologia Centrali-Americana. London, Taylor and Francis, vol. 2 (1891–1900), [4]+622 pp.; vol. 3 (1881–1900), [4] pp.+101 pls. (colored).

Dyar, Harrison Gray

1905. A review of the Hesperiidae of the United States. Jour. New York Ent. Soc., vol. 13, pp. 111-141.

EDWARDS, WILLIAM HENRY

1882. Description of species of butterflies taken in Arizona by Jacob Doll, 1881. Papilio, vol. 2, pp. 19–29.

FREEMAN, HUGH AVERY

1950. Notes on Megathymus, with description of a new species (Lepidoptera, Rhopalocera, Megathymidae). Field and Lab., vol. 18, pp. 144–146.

HEMMING, FRANCIS (ED.)

1950. International Commission on Zoological Nomenclature. Session held during the Thirteenth International Congress of Zoology, Paris, 21st-27th July, 1948. Conclusions of the Fourth Meeting held at the Sorbonne in the Amphithéâtre Louis-Liard on Thursday, 22nd July, 1948, at 0900 hours. Bull. Zool. Nomenclature, vol. 4, pp. 62-82.

1953a. Copenhagen decisions on zoological nomenclature. Additions to, and modifications of, the Règles Internationales de la Nomenclature Zoologique. Approved and adopted by the Fourteenth International Congress of Zoology, Copenhagen, August, 1953. London, International Trust for Zoological Nomenclature, xxx+136 pp., 2 pls.

1953b. Copenhagen discussions, Case no. 6. Documents relevant to the consideration of the status to be accorded to a trivial name, which when first published, was stated to be a substitute for a previously published name but was in addition applied to a particular species or to particular specimens in cases where the species concerned differs from that to which the earlier name is applicable. Bull. Zool. Nomenclature, vol. 10, pp. 167–198.

[MS] [Abstract of report of Mr. Hemming.] Z.N.(C.)9 15th July, 1953. Reference Z.N.(S.)361. Trivial names published partly as substitute names and partly as names for specified species or specified specimens.

SKINNER, HENRY

1911. The larger boreal American Hesperidae, including Eudamus, Erycides, Pyrrhopyge and Megathymus. Trans. Amer. Ent. Soc., vol. 37, pp. 169–209, pl. 10 (colored).

SKINNER, HENRY, AND ROSWELL CARTER WILLIAMS, JR.

1924. On the male genitalia of the Hesperiidae of North America. Paper 6. Trans. Amer. Ent. Soc., vol. 50, pp. 177-208, figs. 1-26.

STRECKER, FERDINAND HEINRICH HERMAN

 Lepidoptera, Rhopaloceres and Heteroceres, indigenous and exotic. Supplement no. 3. Reading, Pennsylvania, printed for the author, pp. 13-38.

The pagination is as given in a copy of this work in the library of the junior author.

