

AMERICAN MUSEUM NOVITATES

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY
CITY OF NEW YORK MARCH 21, 1953 NUMBER 1615

THE GREGORIO BONDAR TYPES OF SOUTH AMERICAN WEEVILS (COLEOPTERA, CURCULIONIDAE)

BY PATRICIA VAURIE¹

INTRODUCTION

The Bondar collection of South American Curculionidae, some 16,000 specimens, procured for the American Museum of Natural History by David Rockefeller, contains the types or cotypes of 318 species described by Gregorio Bondar. Holotypes were designated by Bondar for only 81 species, the rest being represented by cotype series. Because this is an important collection and for other reasons given below, lectotypes have now been chosen for most of the remaining species. Although it is generally advisable for lectotypes to be designated by the reviser of a group, in the present case because Bondar's species occur throughout the entire enormous family of the Curculionidae and it may be many years before the majority of these groups are revised, and because the bulk of the original series or representatives of the original series are in the American Museum of Natural History, it was thought preferable to choose the lectotypes now.

The present paper lists alphabetically all the species described by Bondar up to December, 1950, and gives the holotype or lectotype locality, the date and host plant, if known, and the number of specimens in the original series, with also the subsequent disposition of these specimens, if known.

From a reading of Bondar's works, most of which appeared in the *Revista de Entomologia* from 1937 to 1950, and from a critical study of his material, it is apparent that he gave little

¹ Assistant, Department of Insects and Spiders.

importance to "so-called types" (Rev. Ent., 1946, vol. 17, p. 81), but was far more interested in the biology of the species and in their relation to and dependence on certain plants. He used the words "type and cotype," "cotype and paratype" virtually as synonyms of one another, and, although he may state that the "type and paratype" are in his collection, the specimens are usually all labeled cotypes, or he may have a unique holotype but the specimen bears a cotype label. There are many metatypes of the more abundant species because he placed cotype labels on specimens after the publication of his description, thus creating far too many cotypes of some species. Lectotypes have therefore been chosen to clarify this situation and to associate one specimen definitely with the description of each species.

Bondar ascribed a few species to "Hustache, *in litt.*," or "Hustache, 1940, n. sp.," and in subsequent papers referred to these as Hustache's species. Since, however, Bondar is the first publisher of the name, he becomes the author "unless it is clear from the contents of the publication that some other person is responsible for said name and its indication, definition, or description" (Article 21 of the Code). In the case of each of the species listed below, Bondar explained in small print above the species he was about to describe that he was describing it but was using the name proposed by Hustache before interruption of their correspondence by the war. The species are *Opseobaris albus*, *Microstrates bondari*, *Phytotribus cocosae*, *Hoplorrhinus copiobae*, *Strongylotes dimorphus*, *Derelomus elaisae*, and *Anopsilus striatipennis*.

ACKNOWLEDGMENTS

Thanks are extended to Dr. Mont A. Cazier, Chairman of the Department of Insects and Spiders of the American Museum of Natural History, who suggested this paper, for his helpful advice in the course of procedure to follow. Dr. Rupert L. Wenzel of the Chicago Natural History Museum has kindly lent specimens of some of the Bondar cotypes in that museum and I wish to thank him for his cooperation. I also wish to thank Miss Rose E. Warner of the United States National Museum for sending me a list of the Bondar cotypes in that institution. I am very grateful to Dr. Bondar himself, who has been most helpful in identifying certain type localities in Brazil and in clarifying some of the questions raised concerning his types or cotypes.

CONDITION OF THE COLLECTION

The specimens, large or small, are generally mounted on their sides on paper points, sometimes many on a point and many points to each pin. Locality labels are often lacking, but each pin bears a number referring to entries in Bondar's notebooks, now in the possession of the American Museum. New labels for locality, holotype, and lectotype have been made for all material.

HOLOTYPES

Fifty-eight of Bondar's 81 holotypes are unique specimens. For the 23 species represented by more than one specimen Bondar may or may not have designated holotypes in his description. When he did do so, he did not necessarily label the specimen as holotype. He has told me (*in litt.*) that in such instances the first specimen on the pin is the holotype, but sometimes there have been two or more pins or two specimens side by side on the same paper point. In such cases, where there is no way of checking which specimen is the holotype, either by its position on the pin or by date or locality, I have had to designate lectotypes from the cotype series, even though the holotype is supposed to be present.

LECTOTYPES

According to Bondar, most of the original specimens of his species are in his collection, but wherever there is doubt, due to the presence of metatypes, no lectotype was designated.

In view of the fact that the male genitalia are proving significant in the classification of many weevils, especially in the Barinae, which include the majority of Bondar's species, I have chosen male lectotypes whenever possible. In some tribes of the Barinae, where secondary sexual characters are feeble or not observable, I have made dissections in order to find a male, but this was not done unless there were at least four cotypes available. In species where he could differentiate the sexes, Bondar usually described them both, sometimes placing the description of the female first, sometimes that of the male, but since he did not designate a holotype, the lectotype could be chosen of either sex.

The lectotypes of four species were designated from material lent by the Chicago Natural History Museum, and they are deposited in that institution.

KEY TO INSTITUTIONS CONTAINING COTYPES

The institutions or private collections that are stated by Bondar to contain additional cotypes of his species are given below. The name of the institution follows the city in which it is located, and only the city is referred to in the text. A key letter has been used for those cities with more than one institution represented.

The private collections are those of Borgmeier in Rio de Janeiro, Brazil; Hustache in Lagny, France (now in all or in part at the Muséum National d'Histoire Naturelle in Paris); Kuschel in Santiago, Chile; Voss in Berlin; Zikan in Rio de Janeiro.

Bahia	Instituto Biologico da Bahia
Berlin	Deutsches Entomologisches Institut
Buenos Aires	Departamento de Zoologia Agricola do Ministerio da Agricultura
Chicago	Chicago Natural History Museum
London	A. Imperial Institute of Entomology B. British Museum (Natural History)
Paris	Muséum National d'Histoire Naturelle
Rio de Janeiro	A. Escola Nacional de Agronomia B. Museu Nacional de Rio de Janeiro C. Ministerio da Agricultura, Seccão de Entomologia D. Instituto de Biologia Vegetal E. Instituto Oswaldo Cruz
São Paulo	A. Departamento de Zoologia de São Paulo B. Instituto Biologico de São Paulo C. Museu Paulista
Tucuman	Instituto Miguel Lillo
Washington	United States National Museum

ANNOTATED LIST OF BONDAR'S TYPES

Species are listed under their original generic names. Localities are given with the smallest unit first; this may be an estate or farm ("fazenda"), a village, town, or city; the state is given last. All localities are in Brazil unless otherwise stated (there are a few species from Argentina, Chile, and Paraguay). All cotypes or paratypes mentioned under each species are from the same locality and bear the same date as the holotype or designated lectotype of that species, unless otherwise stated.

abbreviatus, *Microstrates*. 1949, Rev. Ent., vol. 20, p. 216.

Lectotype, male, from Chapada Diamantina, Mucugê, Bahia, in *Cocos*, designated from 63 cotypes in original series of 74. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

acaciae, Chalcodermus. 1941, Rev. Ent., vol. 20, p. 184.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Acacia*, designated from 10 cotypes in original series of 19. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

acaciae, Thanius. 1949, Rev. Ent., vol. 20, p. 198.

Lectotype, male, from Ondina, Bahia, January, 1946, in *Acacia*, designated from the five original cotypes. One cotype stated by Bondar to be in Chicago, and it is known to be in that institution.

aechmeae, Bromegeraeus. 1942, Rev. Ent., vol. 13, p. 252, figs. 22-23.

Lectotype, sex undetermined, from Nazaré, Bahia, in *Aechmea marmorata*, designated from two cotypes in original series of seven. Other cotypes stated by Bondar to be in Chicago.

albidus, Ancylorrhynchus. 1943, Rev. Ent., vol. 14, p. 362.

Lectotype, male, from Campo Formoso, Bahia, in *Cocos coronata*, designated from six cotypes in original series of eight. Location of other cotypes unknown.

albinotatus, Lispodemus. 1942, Rev. Ent., vol. 13, p. 264.

Holotype, sex undetermined, from Fazenda dos Campos, Virginia, Minas Gerais, 1500 meters. Unique specimen.

albonotatus, Conothorax. 1949, Rev. Ent., vol. 20, p. 179.

Holotype, sex undetermined, from the state of Rio de Janeiro. Location of paratype from São Paulo, November, 1938, unknown. One paratype stated by Bondar to be in London (A).

alboscuteclatus, Polyponus. 1938, Rev. Ent., vol. 8, p. 16, fig. 6.

Holotype, sex undetermined, from Agua Preta, Bahia, labeled cotype in error. "Type in collection of author." Five paratypes stated by Bondar to be in Berlin and Rio de Janeiro (D).

albosignatus, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 321.

Lectotype, sex undetermined, from Ilha dos Frades, Bahia, in Mirtaceae, designated from two cotypes in original series of six. Location of other cotypes unknown.

albosignatus, Torcobius. 1943, Rev. Ent., vol. 14, p. 55.

Lectotype, male, from Bahia, Bahia, in *Rhynchospora*, designated from two cotypes in original series of seven. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

albus, Opseobaris. 1942, Rev. Ent., vol. 13, p. 271.

Lectotype, male, from San Roque, Corrientes, Argentina, February, 1920, in Gramineae, designated from four cotypes in

original series of 13. Other cotypes stated by Bondar to be in Chicago, Rio de Janeiro (A), and São Paulo (A).

amputator, Putatocentrus. 1949, Rev. Ent., vol. 20, p. 203, fig. 12.

Lectotype, male, from Caitité, Bahia, October, 1947, designated from four cotypes in original series of five. Other cotype stated by Bondar to be in Chicago.

andaiae, Phytotribus. 1941, Rev. Ent., vol. 12, p. 292, fig. 11.

Lectotype, male, from Rio Novo, Bahia, in *Attalea compta*, designated from five cotypes in original series of 22. Location of other cotypes unknown. "Type and cotype in collection of author," but all specimens labeled cotypes.

anomala, Liobaridia. 1950, Rev. Ent., vol. 21, p. 469, fig. 15.

Lectotype, male, from Porto Alegre, Rio Grande do Sul, January, 1948, in *Baccharis anomala*, designated from 24 cotypes in original series of 42. Location of other cotypes unknown.

anonae, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 200.

Lectotype, sex undetermined, from Campo Formoso, Bahia, in *Anona coriacea*, designated from one cotype in original series of five. Location of other cotypes unknown.

anonicola, Meroprion. 1939, Rev. Ent., vol. 10, p. 11, fig. 3.

Lectotype, sex undetermined, from Itaberaba, Bahia, in *Anona squamosa*, designated from 10 cotypes (size of original series not stated). "Paratypes" stated by Bondar to be in Berlin. "Type in collection of author," but no specimen is so marked. Three other localities in Bahia given in description, but none is associated with type.

antennalis, Gladosius. 1946, Rev. Ent., vol. 17, p. 333.

Holotype, "possibly male," from Itatiaia, Rio de Janeiro, November, 1933, labeled cotype in error. Unique specimen.

antiquus, Mallerus. 1946, Rev. Ent., vol. 17, p. 86, figs. 7-11.

Lectotype, male, from the state of Santa Catarina, designated from four cotypes in original series of eight. Location of other cotypes unknown.

ariri, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 220.

Lectotype, male, from Lagedo Alto, Bahia, May, 1941, in *Cocos vagans*, designated from nine cotypes in original series of 11. Other cotypes stated by Bondar to be in Chicago.

armatus, Valliopsis. 1948, Rev. Ent., vol. 19, p. 30.

Lectotype, male, from Belmonte, Bahia, October 20, 1944,

in Marantaceae, designated from the two original cotypes. Second cotype is a female.

aterrimus, Tripusus. 1949, Rev. Ent., vol. 20, p. 215, fig. 11.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Diplothemium campestre*, designated from three cotypes in original series of five. Other cotypes stated by Bondar to be in Chicago.

atrata, Bromesia. 1946, Rev. Ent., vol. 17, p. 336.

Lectotype, male, from Vitoria, Espirito Santo, November, 1943, in *Araecoccus*, designated from 32 cotypes in original series of 50. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (B).

atromaculatus, Eurylobus. 1949, Rev. Ent., vol. 20, p. 178.

Holotype, sex undetermined, from Obidos, Amazonas, 1922, labeled cotype in error. Unique specimen.

bahiensis, Apantoteles. 1946, Rev. Ent., vol. 17, p. 84, figs. 3-6.

Holotype, probably male, from Belmonte, Bahia, February 20, 1945. Unique specimen.

bahiensis, Sibariops. 1943, Rev. Ent., vol. 14, p. 75.

Lectotype, male, from Fazenda Pasto Fora, Bahia, Bahia, August 23, 1942, in Ciperaceae, designated from the 14 original cotypes. Although Bondar's description states there were but 13 cotypes, he has 14 listed in the numbered entry in his notebook.

barbiellinii, Cymatobaris. 1949, Rev. Ent., vol. 20, p. 197.

Lectotype, sex undetermined, from Cubatão, São Paulo, designated from one cotype in original series of two. Location of other cotype unknown.

basalis, Glyptobaris. 1946, Rev. Ent., vol. 17, p. 99.

Holotype, sex undetermined, from Agua Preta, Bahia, October 1934, in Convolvulaceae. Unique specimen.

beckeri, Baris. 1949, Rev. Ent., vol. 20, p. 200.

Lectotype, male, from Porto Alegre, Rio Grande do Sul, November, 1947, in *Aspilia setosa*, designated from 12 cotypes in original series of 16. Other cotypes stated by Bondar to be in Chicago.

beckeri, Diorymerus. 1950, Rev. Ent., vol. 21, p. 460, fig. 12.

Holotype, male, from São Gonçalo dos Campos, Bahia, January, 1946, in Malpighiaceae, labeled cotype in error. Location of female paratype unknown.

beckeri, Gladosius. 1949, Rev. Ent., vol. 20, p. 205.

Holotype, "probably female," from Maracás, Bahia, March, 1945, in Bromeliaceae. Unique specimen.

belmonte, Valliopsis. 1948, Rev. Ent., vol. 19, p. 29.

Lectotype, male, from Belmonte, Bahia, October 20, 1944, designated from the two original cotypes.

biancae, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 454.

Lectotype, male, from Fazenda Babassú, São Mateus, Espírito Santo, in *Attalea compta*, designated from two cotypes in original series of nine. Other cotypes stated by Bondar to be in Chicago and São Paulo (C).

bicolor, Gladosius. 1942, Rev. Ent., vol. 13, p. 245.

Holotype, female, from Itatiaia, Rio de Janeiro, November 20, 1931, labeled cotype in error. Unique specimen.

bicolor, Hulpes. 1942, Rev. Ent., vol. 14, p. 374.

Lectotype, male, from the state of Santa Catarina, designated from the 40 original cotypes.

bicolor, Preglyptobaris. 1946, Rev. Ent., vol. 17, p. 100.

Lectotype, male, from Fazenda dos Campos, Virginia, Minas Gerais, November 1, 1917, designated from two cotypes in original series of five. Other cotypes stated by Bondar to be in Chicago and in the collection of Zikan.

bicolor, Scymnoplastophilus. 1947, Rev. Ent., vol. 18, p. 283.

Lectotype, sex undetermined, from Corupá, Santa Catarina, designated from the two original cotypes. Other cotype is from Nova Teutonia, Santa Catarina.

bifasciatus, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 453.

Holotype, female, from São Mateus, Espírito Santo, June 12, 1941, in *Attalea humilis* (?). One female, labeled cotype.

bifasciatus, Sibariops. 1943, Rev. Ent., vol. 14, p. 71.

Holotype, sex undetermined, from Pituba, Bahia, November 30, 1942, in *Rhynchospora cephalotes*, labeled cotype in error. Unique specimen.

binotatus, Derelomus. 1941, Rev. Ent., vol. 12, p. 462, figs. 20-22.

Lectotype, male, from Fazenda Babassú, São Mateus, Espírito Santo, June 12, 1941, in *Attalea compta*, designated from 140 cotypes in original series of "about 390." Other cotypes stated by Bondar to be in Chicago and São Paulo (C).

bleyi, Ancylorrhynchus. 1941, Rev. Ent., vol. 12, p. 464, figs. 23-27.

Lectotype, male, from Vitoria, Espírito Santo, June 26,

1941, in *Cocos picrophylla*, designated from two cotypes in original series of nine. One cotype is in Chicago; the others are stated by Bondar to be in São Paulo (C).

bondari, Microstrates. 1941, Rev. Ent., vol. 12, p. 300, fig. 15.

Lectotype, male, from Nazaré, Bahia, in *Cocos botryophora*, designated from three cotypes in original series of 10. Location of other cotypes unknown. "Type and cotype in collection of author," but all specimens labeled cotypes.

botryophorae, Ancylorrhynchus. 1941, Rev. Ent., vol. 12, p. 295, fig. 13.

Lectotype, female, from the state of Bahia, in *Cocos botryophora*, designated from two cotypes in original series of six. Location of other cotypes unknown. "Type in collection of author," but both specimens labeled cotypes.

braga, Celetes. 1943, Rev. Ent., vol. 14, p. 348.

Lectotype, male, from São Mateus, Espírito Santo, June, 1941, in *Attalea compta*, designated from the three original cotypes. The two other cotypes are females.

brasiliensis, Cavibaris. 1946, Rev. Ent., vol. 17, p. 106.

Lectotype, male, from the state of Rio Grande do Sul, January, 1944, designated from two cotypes in original series of 10. Other cotypes stated by Bondar to be in the collections of Hustache and Zikan.

bridarollii, Borgmeierus. 1945, Rev. Ent., vol. 16, p. 111.

Lectotype, male, from Córdoba, Argentina, January 11, 1930, in *Cyperus*, designated from the three original cotypes.

bromeliae, Gladosius. 1942, Rev. Ent., vol. 13, p. 243.

Holotype, male, from Passa Quatro, Minas Gerais, March 17, 1922, in Bromeliaceae, labeled cotype in error. One female labeled cotype. Holotype is top specimen on pin (Bondar, *in litt.*).

bruniceps, Amphibaris. 1946, Rev. Ent., vol. 17, p. 102.

No lectotype designated. Species described from nine specimens from Agua Preta, Bahia, October, 1934, in Convolvulaceae. Date on our six cotypes, which are probably metatypes, is 1938 instead of 1934. Some cotypes stated by Bondar to be in Chicago, but that institution does not have them at present (Wenzel, *in litt.*)

huri, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 457.

Lectotype, female, from São Mateus, Espírito Santo, June 13, 1941, in *Diplothemium caudescens*, designated from seven

cotypes in original series of 15. Other cotypes stated by Bondar to be in Chicago and São Paulo (C), but there are none in Chicago (Wenzel, *in litt.*).

butia, Palmocentrinus. 1949, Rev. Ent., vol. 20, p. 207.

Lectotype, male, from the state of Santa Catarina, in *Cocos eriospatha*, designated from eight cotypes in original series of nine. Other cotype stated by Bondar to be in Chicago.

butia, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 222.

Lectotype, male, from Curitiba, Paraná, in *Cocos eriospatha*, designated from 15 cotypes in original series of 21. Other cotypes stated by Bondar to be in Chicago.

cachoeiranus, Phytotribus. 1943, Rev. Ent., vol. 14, p. 353.

Lectotype, male, from Cachoeira, Bahia, in *Attalea borge-siana*, designated from five cotypes in original series of 17. Location of other cotypes unknown.

caerulescens, Thanius. 1949, Rev. Ent., vol. 20, p. 198.

Lectotype, male, from Ponta Grossa, Paraná, designated from the eight original cotypes.

caldeirai, Thoracocyphus. 1949, Rev. Ent., vol. 20, p. 174, fig. 2.

Lectotype, male, from Riacho de Santa Ana, Bahia, October 21, 1947, in Leguminosae, designated from four cotypes in original series of 13. Other cotypes stated by Bondar to be in Chicago.

calvus, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 217.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Cocos*, designated from 16 cotypes in original series of 21. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

cambotae, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 323.

Lectotype, sex undetermined, from Agua Preta, Bahia, in *Thalissia* (?), designated from two cotypes in original series of six. Location of other cotypes unknown.

camelodeformis, Stegotes. 1946, Rev. Ent., vol. 17, p. 317.

Lectotype, male, from Fazenda Ermida, Belmonte, Bahia, October, 1944, in *Acacia*, designated from the four original cotypes.

caminha, Heterosternus. 1943, Rev. Ent., vol. 14, p. 112, fig. 23.

Lectotype, male, from Usina Aliança, Bahia, March, 1943, in Labiaceae, designated from 31 cotypes in original series of

70. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

campestris, Celetes. 1949, Rev. Ent., vol. 20, p. 180.

Lectotype, male, from Mucugê, Bahia, 1000 meters, October, 1948, in *Diplothemium campestre*, designated from 41 cotypes in original series of 104. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

campestris, Dialomia. 1942, Rev. Ent., vol. 13, p. 16.

Holotype, male, from Jiqui, Barra da Estiva, central Bahia, May, 1941, in *Cocos campestris*. One female paratype.

campestris, Hustachea. 1942, Rev. Ent., vol. 13, p. 33.

Lectotype, male, from Jiqui, Barra da Estiva, central Bahia, May, 1941, in *Cocos campestris*, designated from 11 cotypes in original series of 18. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B).

campestris, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 218.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Diplothemium campestre*, designated from 86 cotypes in original series of 109. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

campestris, Remetrus. 1949, Rev. Ent., vol. 20, p. 211, fig. 14.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Diplothemium campestre*, designated from 30 cotypes in original series of 38. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

camposi, Ancylorrhynchus. 1941, Rev. Ent., vol. 12, p. 466, fig. 28.

Lectotype, female, from Jiqui, Bahia, May, 1941, in *Cocos campestris*, designated from five cotypes in original series of 15. Two cotypes are in Chicago, the others stated by Bondar to be in São Paulo (C).

camposi, Chalcodermus. 1939, Rev. Ent., vol. 10, p. 13, fig. 5.

Holotype, sex undetermined, from Agua Preta, Bahia, in "cacaoeiro." One paratype stated by Bondar to be in Rio de Janeiro (C).

canavaliae, Chalcodermus. 1941, Rev. Ent., vol. 12, p. 433, figs. 7-8.

Holotype, sex undetermined, from Bahia, Bahia, in *Canavalia obtusifolia*, labeled cotype in error. "Type and paratype in collection of author," but location of paratype unknown.

capichaba, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 51.

Holotype, sex undetermined, from Vitoria, Espirito Santo, labeled cotype in error. Unique specimen.

capichaba, Gladosius. 1942, Rev. Ent., vol. 13, p. 345.

Lectotype, female, from Vitoria, Espirito Santo, April 2, 1941, designated from one cotype in original series of two. Location of other cotype (state of Bahia, no date) unknown.

capichaba, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 451, fig. 15.

Holotype, male, from Fazenda Babassú, São Mateus, Espirito Santo, June 12, 1941, in *Attalea compta*. One male labeled cotype. Location of female paratype unknown.

caseariae, Spermobaris. 1949, Rev. Ent., vol. 20, p. 202, fig. 10.

Lectotype, male, from Porto Alegre, Rio Grande do Sul, September, 1946, in *Casearia sylvestris*, designated from nine cotypes in original series of 11. Other cotypes stated by Bondar to be in Chicago.

caseyanus, Sibariops. 1943, Rev. Ent., vol. 14, p. 69.

Lectotype, male, from Bahia, Bahia, August, 1942, in Ciperaceae, designated from the nine original cotypes.

caseyi, Linonotus. 1948, Rev. Ent., vol. 19, p. 25.

Holotype, female, from Corupá, Santa Catarina. Unique specimen.

cearanus, Microstrates. 1942, Rev. Ent., vol. 13, p. 29.

Lectotype, male, from the state of Ceara (Bondar, *in litt.*) in *Cocos picrophylla*, designated from 46 cotypes in original series of 110. Other cotypes stated by Bondar to be in Chicago and São Paulo (D, E).

cecropiae, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 193.

Lectotype, male, from Agua Preta, Bahia, January 20, 1930, in *Cecropia cinerea*, designated from eight cotypes in original series of 19. Two cotypes are known to be lost; location of others unknown. Above locality is from cotype labels, as no locality was given in original description.

cesalpiniae, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 326.

No lectotype designated. Species described from three specimens from Lagedo Alto, Bahia, in *Cesalpinia microphylla*. We have 12 specimens, same locality, labeled as cotypes, all mounted on paper cards on one pin; three of these might be the original specimens and the others metatypes, but as they are mounted in pairs it is impossible to know if this is true.

chusqueae, Notolomus. 1949, Rev. Ent., vol. 20, p. 182.

Lectotype, male, from Cordillera Pemehue, Malleco, Chile,

January, 1946, in *Chusquea culeou*, designated from 17 cotypes in original series of 43. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

cissophagus, Dinomorphus. 1948, Rev. Ent., vol. 19, p. 5, figs. 6-11.

Lectotype, male, from Bom Jesus da Lapa, Bahia, October, 1947, in *Cissus*, designated from 17 cotypes in original series of 93. Other cotypes stated by Bondar to be in Chicago, London (A), Paris, and Rio de Janeiro (A).

cocois, Microstrates. 1941, Rev. Ent., vol. 12, p. 301, fig. 16.

Lectotype, male, from Florianopolis, Santa Catarina, in *Cocos Romanzoffiana*, designated from 96 cotypes in original series of "about 150." Location of other cotypes unknown. "Type, cotype, and paratypes in collection of author," but all specimens are labeled cotypes.

cocosae, Phytotribus. 1941, Rev. Ent., vol. 12, p. 446, figs. 10, 12.

Lectotype, male, from Nazaré, Bahia, in *Cocos botryophora*, designated from nine cotypes in original series of 42. Lectotype and eight cotypes in Chicago. Other cotypes stated by Bondar to be in São Paulo (C) and collection of Hustache. Although cotypes were also stated by Bondar to be in his collection, our 24 specimens are not so labeled.

cocos-campestris, Microstrates. 1942, Rev. Ent., vol. 13, p. 28.

Lectotype, male, from Jiqui, Barra da Estiva, central Bahia, May 6, 1941, in *Cocos campestris*, designated from 32 cotypes in original series of 50. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B).

complanatus, Monteus. 1943, Rev. Ent., vol. 14, p. 46.

Holotype, "probably female," from Itatiaia, Rio de Janeiro. Location of paratype unknown.

copaiferae, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 197.

Lectotype, sex undetermined, from Agua Preta, Bahia, in *Copaifera Langsdorffii*, designated from 10 cotypes in original series of 20. Location of other cotypes unknown except for two which Bondar says in his notebook were sent to London (B).

copiobae, Hoplorrhinus. 1941, Rev. Ent., vol. 12, p. 285.

Lectotype, male, from the state of Bahia, in *Attalea*, designated from 23 cotypes in original series of "abundant material." Two cotypes are in Washington (Warner, *in litt.*); others stated by Bondar to be in Berlin and in collection of Hustache. "Type

and cotype in collection of author," but all specimens are labeled cotypes.

corymbosae, Parallelosomus. 1950, Rev. Ent., vol. 21, p. 477.

Lectotype, male, from Porto Alegre, Rio Grande do Sul, in *Rhynchospora corymbosa*, designated from five cotypes in original series of six. Other cotypes [metatypes?] stated by Bondar to be in Chicago and Rio de Janeiro (A).

costalimai, Angelocentris. 1949, Rev. Ent., vol. 20, p. 213, fig. 13.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Cocos*, designated from six cotypes in original series of eight. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

costalimai, Parallelosomus. 1943, Rev. Ent., vol. 14, p. 41.

Lectotype, male, from Pituba, Bahia, May to July, 1942, in *Rhynchospora scaberrima*, designated from 75 cotypes in original series of 200. Other cotypes stated by Bondar to be in Chicago, Rio de Janeiro (A), São Paulo (A), and Buenos Aires.

costalimai, Polpones. 1943, Rev. Ent., vol. 14, p. 50.

Lectotype, male, from Aliança, Bahia, in either *Rhynchospora cyperoides* or *R. scaberrima*, designated from 88 cotypes, although original series included but 63. Some of these cotypes are undoubtedly metatypes, but as there are 10 to 12 specimens on each pin, it is impossible to know where the original series of 63 specimens ends. Other cotypes stated by Bondar to be in Chicago, Rio de Janeiro (A), and São Paulo (A, B).

crotoni, Spermobaris. 1943, Rev. Ent., vol. 14, p. 114, figs. 24-26.

Lectotype, male, from Feira de Santa Ana, Bahia, in *Croton*, designated from 29 cotypes in original series of 46. Other cotypes stated by Bondar to be in Chicago and São Paulo (A). The lectotype locality is taken from Bondar's notebook; no locality was given in description.

cruciferae, Radamus. 1946, Rev. Ent., vol. 17, p. 104.

Lectotype, male, from Belmonte, Bahia, March, 1945, in *Crucifera*, designated from 31 cotypes in original series of 50. Other cotypes stated by Bondar to be in Chicago and in the collection of Hustache.

curtus, Phytotribus. 1941, Rev. Ent., vol. 12, p. 447, fig. 11.

Holotype, male, from São Mateus, Espírito Santo, June 12, 1941, in *Attalea compta*. Location of female paratype unknown.

cydoniae, Apion. 1950, Rev. Ent., vol. 21, p. 454, fig. 9.

Lectotype, male, from Jacobina, Bahia, April, 1949, in *Cydonia*

oblonga, designated from 34 cotypes in original series of 84. Other cotypes stated by Bondar to be in Bahia, Chicago, and Rio de Janeiro (A).

dahlgreni, Hoplorrhinus. 1941, Rev. Ent., vol. 12, p. 290.

Holotype, female, from Nazaré, Bahia, in *Cocos botryophora*, labeled cotype in error. One male paratype. Location of female paratype unknown. Holotype is top specimen on pin (Bondar, *in litt.*).

decemmaculatus, Polpones. 1945, Rev. Ent., vol. 16, p. 111.

Lectotype, probably male, from Rio Claro, São Paulo, October, 1934, designated from the six original cotypes.

decolor, Celetes. 1943, Rev. Ent., vol. 14, p. 347.

Lectotype, male, from Bahia, Bahia, in *Cocos coronata*, designated from 61 cotypes in original series of "abundant material."

depressa, Stereobaris. 1949, Rev. Ent., vol. 20, p. 196.

Lectotype, male, from the state of Santa Catarina, December, 1942, designated from the two original cotypes.

dimorphus, Strongylotes. 1942, Rev. Ent., vol. 13, p. 269.

Lectotype, male, from Mafra, Santa Catarina, January, 1942, in Gramineae, designated from four cotypes in original series of eight. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

diplothemii, Dialomia. 1942, Rev. Ent., vol. 13, p. 15.

Lectotype, male, from São Mateus, Espírito Santo, June, 1941, in *Diplothemium caudescens*, designated from the three original cotypes.

discordis, Tripestes. 1946, Rev. Ent., vol. 17, p. 91.

Holotype, sex undetermined, from the state of Paraná. Unique specimen.

dispar, Amphibaris. 1946, Rev. Ent., vol. 17, p. 101.

Lectotype, sex undetermined, from Jacaranda, Canavieiras, Bahia, in Convolvulaceae, designated from the two original cotypes.

divergens, Stegotes. 1946, Rev. Ent., vol. 17, p. 325.

Holotype, male, from Funil, Minas Gerais. Unique specimen.

drewsi, Laemosaccus. 1947, Rev. Ent., vol. 18, p. 293.

Holotype, "probably" female, from Fazenda Campos, Virginia, Minas Gerais, labeled cotype in error. Unique specimen.

dubius, Sibariops. 1943, Rev. Ent., vol. 14, p. 77.

Lectotype, sex undetermined, from Bahia, Bahia, in *Cocos schizophylla*, designated from the two original cotypes.

efraimi, Metriophilus. 1942, Rev. Ent., vol. 13, p. 4.

Lectotype, sex undetermined, from Santa Teresinha, Bahia, in Leguminosae, designated from eight cotypes in original series of 46. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B).

elaisae, Derelomus. 1941, Rev. Ent., vol. 12, p. 461, figs. 17-19.

Lectotype, male, from the state of Bahia, in *Elaeis guineensis*, designated from 22 cotypes in original series of 100. Twenty-five cotypes are in Washington (Warner, *in litt.*); others stated by Bondar to be in Chicago, São Paulo (C), and the collection of Hustache.

elenaee, Gladosius. 1942, Rev. Ent., vol. 13, p. 247.

Holotype, sex undetermined, from Itatiaia, Rio de Janeiro, March 19, 1926, labeled cotype in error. Unique specimen.

eriospathae, Ancyloirrhynchus. 1943, Rev. Ent., vol. 14, p. 364.

Lectotype, male, from Curitiba, Paraná, in *Cocos eriospatha*, designated from 10 cotypes in original series of 31. Location of other cotypes unknown.

eucalypti, Miremorphus. 1950, Rev. Ent., vol. 21, p. 451, figs. 1-6.

Lectotype, sex undetermined, from Caraiba, Goyaz, in *Eucalyptus*, designated from five cotypes in original series of 10. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

externus, Dioryserratus. 1950, Rev. Ent., vol. 21, p. 466, fig. 14.

Holotype, sex undetermined, from the state of Paraná. Unique specimen.

fariai, Hoplorrhinus. 1941, Rev. Ent., vol. 12, p. 289, fig. 9.

Lectotype, male, from Cachoeira, Bahia, in *Attalea humilis*, designated from five cotypes in original series of seven. Location of other cotypes unknown. "Type and cotype in collection of author," but all specimens labeled cotypes.

farinarius, Plectrodontus. 1938, Rev. Ent., vol. 8, p. 7, figs. 7, 8.

Holotype, sex undetermined, from Agua Preta, Bahia, labeled cotype in error. Holotype is top specimen on pin (Bondar, *in litt.*). One paratype. About 38 paratypes stated by Bondar to be in Berlin, London (A), Rio de Janeiro (D, E), and in the collection of Voss.

fasciatus, Thaliabaris. 1945, Rev. Ent., vol. 16, p. 109.

Lectotype, probably female, from Rio Bravo, Entre Rios

Argentina, February 6, 1943, in *Thalia multiflora*, designated from five cotypes in original series of 10. Other cotypes stated by Bondar to be in the collection of Kuschel.

fernando-costai, Balanephagus. 1941, *Rev. Ent.*, vol. 12, p. 297, fig. 18.

Holotype male, from Fazenda Copiobá, Nazaré, Bahia, November 20, 1940, in *Diplothemium caudescens*, labeled cotype in error. "Type and cotype in collection of author," but one female "cotype" is in Chicago, and location of eight other paratypes is unknown.

ficus, Metoposoma. 1950, *Rev. Ent.*, vol. 21, p. 456, figs. 10-11.

Lectotype, male, from Boa Nova, Bahia, May 12, 1949, in *Ficus benjamina*, designated from 48 cotypes in original series of 132. Other cotypes stated by Bondar to be in Bahia, Chicago, London (B), and Rio de Janeiro (A).

fiedleri, Conotrachelus. 1944, *Rev. Ent.*, vol. 15, p. 195.

Lectotype, sex undetermined, from Ilha dos Frades, Bahia, in Mirtaceae, designated from two cotypes in original series of three. Location of other cotype unknown.

flavipennis, Thaliabaris. 1943, *Rev. Ent.*, vol. 14, p. 133.

Lectotype, male, from Fazenda Bom Gosto, Santo Amaro, Bahia, March 4, 1943, in *Thalia geniculata* or *T. angustifolia*, designated from 58 cotypes in original series of 80. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

flavolineatus, Orissus. 1946, *Rev. Ent.*, vol. 17, p. 332.

Lectotype, male, from the state of Paraná, designated from the two original cotypes, both males.

flavoscutellatus, Prionomerus. 1947, *Rev. Ent.*, vol. 18, p. 280.

Holotype, sex undetermined, from Ponta Grossa, Paraná. Unique specimen.

fontenellei, Hoplorhinus. 1941, *Rev. Ent.*, vol. 12, p. 450, fig. 14.

Lectotype, male, from Fazenda Babassú, São Mateus, Espírito Santo, in *Attalea compta*, designated from 15 cotypes in original series of 48. Two cotypes are in Washington (Warner, *in litt.*), and others stated by Bondar to be in Chicago and São Paulo (C).

franciscanus, Prodinus. 1950, *Rev. Ent.*, vol. 21, p. 464.

Holotype, sex undetermined, from the banks of Rio São Francisco, Lapa, Bahia, October 20, 1947, in Malpighiaceae. Unique specimen.

fuirenae, Sibariops. 1943, *Rev. Ent.*, vol. 14, p. 68.

Lectotype, male, from Pituba, Bahia, March to August, 1942, in *Fuirena umbellata*, designated from 16 cotypes in original series of 150. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

fulvopilosus, Peridinetus. 1949, Rev. Ent., vol. 20, p. 188, fig. 5.

Lectotype, male, from Canavieiras, Bahia, October, 1946, in *Piperacea arbustiva*, designated from four cotypes in original series of five. Other cotypes stated by Bondar to be in Chicago.

funebri, Centrinus. 1943, Rev. Ent., vol. 14, p. 128, fig. 31.

Lectotype, male, from Itatiaia, Rio de Janeiro, January 13, 1927, designated from seven cotypes in original series of 11. This date does not agree with the date given by Bondar (December, 1933), but it may have been overlooked at the time of description. All but two of the missing cotypes are accounted for in entry in Bondar's notebook (one given to Dr. Costa Lima, one to London), and it is assumed the above are part of the original series. One of the 1927 cotypes is in Chicago.

gallicola, Baptoharinus. 1949, Rev. Ent., vol. 20, p. 190, fig. 8.

Lectotype, male, from Ponta Grossa, Paraná, designated from 19 cotypes in original series of 25. Other cotypes stated by Bondar to be in Bahia, Chicago, and Rio de Janeiro (A).

gaucho, Geraeus. 1946, Rev. Ent., vol. 17, p. 334.

Lectotype, male, from Porto Alegre, Rio Grande do Sul, August, 1935, designated from the four original cotypes (two males, two females).

gibber, Justus. 1946, Rev. Ent., vol. 17, p. 94, fig. 12.

Lectotype, male, from Ponta Grossa, Paraná, designated from three cotypes in original series of five. Location of other cotypes unknown.

gibbifrons, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 52.

Holotype, sex undetermined, from Belmonte, Bahia, October, 1945. Unique specimen.

gica, Balanephagus. 1940, Rev. Ent., vol. 11, p. 852.

Holotype, male, from Bomfim, Bahia, in *Cocos coronata*, labeled cotype in error. "Holotype in collection of author." One paratype stated by Bondar to be in Berlin, one in Washington. One paratype is in Chicago.

gica, Thaliabaris. 1943, Rev. Ent., vol. 14, p. 132, figs. 35-40.

Lectotype, male, from Fazenda Bom Gosto, Santa Amaro, Bahia, March 4, 1943, in *Thalia angustifolia* or *T. geniculata*, designated from 56 cotypes in original series of 120. Other

cotypes stated by Bondar to be in Chicago and São Paulo (A).

giganteus, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 220.

Lectotype, male, from Agua Limpa, Minas Gerais, January, 1946, designated from three cotypes in original series of four. Other cotype stated by Bondar to be in Chicago.

guabirobae, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 196.

Lectotype, sex undetermined, from Agua Preta, Bahia, in *Britoa rugosa*, designated from two cotypes in original series of four. Location of other cotypes unknown.

guapevae, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 201.

Holotype, sex undetermined, from Agua Preta, Bahia, in fruits of Sapotaceae. Unique specimen.

guareae, Scymnoplastophilus. 1938, Rev. Ent., vol. 8, p. 12, figs. 11, 13.

Lectotype, sex undetermined, from Agua Preta, Bahia, January 31, 1937, in *Guarea rosea*, designated from eight cotypes in original series of 25. "Type in collection of author," but all labeled cotypes. Other cotypes stated by Bondar to be in London (A) and Rio de Janeiro (D, E).

gudimiri, Prionomerus. 1938, Rev. Ent., vol. 8, p. 5, fig. 5.

Holotype, sex undetermined, from Agua Preta, Bahia, in *Persea gratissima*, labeled cotype in error. Unique specimen.

hannibali, Polpones. 1943, Rev. Ent., vol. 14, p. 51.

Holotype, female, from Fazenda dos Campos, Virginia, Minas Gerais. Unique specimen.

hatschbachi, Ancylorrhynchus. 1943, Rev. Ent., vol. 14, p. 363.

Lectotype, male, from Curitiba, Paraná, in *Cocos eriospatha*, designated from 29 cotypes in original series of 59. Location of other cotypes unknown.

hatschbachi, Microstrates. 1943, Rev. Ent., vol. 14, p. 370.

Lectotype, male, from Curitiba, Paraná, in *Cocos eriospatha*, designated from 19 cotypes in original series of 20. Location of other cotype unknown.

hatschbachi, Thoracocyphus. 1949, Rev. Ent., vol. 20, p. 174, fig. 1.

Holotype, sex undetermined, from Curitiba, Paraná. Unique specimen.

heringeri, Eurhinus. 1948, Rev. Ent., vol. 19, p. 18, fig. 16.

Lectotype, male, from the state of Minas Gerais, September 25, 1946, designated from six cotypes in original series of 14.

Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

heringeri, Trachybaris. 1946, Rev. Ent., vol. 17, p. 109.

Lectotype, sex undetermined, from the state of Minas Gerais, in *Calliclamys latifolia*, designated from seven cotypes in original series of 22. Other cotypes stated by Bondar to be in Chicago and in collection of Hustache.

heterocolor, Stegotes. 1946, Rev. Ent., vol. 17, p. 322.

Lectotype, male, from Fazenda Ermida, Belmonte, Bahia, October, 1944, in *Acacia*, designated from the two original cotypes.

heteropteri, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 50.

Lectotype, sex undetermined, from Porto Alegre, Rio Grande do Sul, April, 1946, in *Heteropteris aenea*, designated from one cotype in original series of three. Other cotypes stated by Bondar to be in Chicago.

heteropteris, Prodinus. 1950, Rev. Ent., vol. 21, p. 465.

Lectotype, sex undetermined, from Porto Alegre, Rio Grande do Sul, April, 1947, in *Heteropteris aenea*, designated from three cotypes in original series of four. Lectotype and one cotype in Chicago. We have seven additional specimens labeled cotypes, but with the date April 12, 1950, which are undoubtedly meta-

horni, Piazorhinus. 1938, Rev. Ent., vol. 8, p. 15.

Holotype, sex undetermined, from Agua Preta, Bahia, in Leguminosae. Two paratypes. Twenty-one paratypes stated by Bondar to be in Berlin, London (A), and Rio de Janeiro (D, E).

humilis, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 455.

Lectotype, male, from the state of Espirito Santo, in *Attalea humilis* (?), designated from 10 cotypes in original series of 54. Other cotypes stated by Bondar to be in Chicago and São Paulo (C).

hustachei, Apocnemidophorus. 1947, Rev. Ent., vol. 18, p. 292.

Lectotype, male, from Ponta Grossa, Paraná, in *Crinum*, designated from two male cotypes in original series of four. Location of other cotypes unknown.

hustachei, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 286, fig. 8.

Lectotype, male, from Fazenda Copiobá, Nazaré, Bahia, November 20, 1940, in *Diplothemium caudescens*, designated from six cotypes in original series of 35. "Type and cotype in collection of author," but all specimens labeled cotypes. There are

now eight cotypes in Washington (Warner, *in litt.*); location of others unknown.

hustachei, Liobaridia. 1950, Rev. Ent., vol. 21, p. 470.

Lectotype, female, from Ponta Grossa, Paraná, designated from two cotypes in original series of three. One of the two cotypes, also a female, is in Chicago; location of other cotype (male) unknown.

inermis, Microstegotes. 1946, Rev. Ent., vol. 17, p. 329.

Holotype, "possibly female," from Rio Negro, Amazonas, November, 1927. Unique specimen.

intermedius, Gladosius. 1942, Rev. Ent., vol. 13, p. 244.

Holotype, "probably female," from Virginia, Minas Gerais, December 1, 1920, labeled cotype in error. Unique specimen.

internus, Dioryserratus. 1950, Rev. Ent., vol. 21, p. 466, fig. 13.

Holotype, sex undetermined, from São Gonçalo dos Campos, Bahia, January, 1946, in Malpighiaceae. Unique specimen.

jacilo, Linonotus. 1943, Rev. Ent., vol. 14, p. 125.

Lectotype, female, from Agua Preta, Bahia, September 8, 1937, in Marantaceae, designated from the two original cotypes, both females.

jiquiensis, Phytotribus. 1941, Rev. Ent., vol. 12, p. 448, fig. 9.

Lectotype, male, from Jiqui, Bahia, May, 1941, in *Cocos campestris*, designated from six cotypes in original series of 32. Other cotypes stated by Bondar to be in Chicago and São Paulo (C).

justus, Dioryserratus. 1950, Rev. Ent., vol. 21, p. 467.

Holotype, sex undetermined, from Ponta Grossa, Paraná, November, 1944. Unique specimen.

justus, Linonotus. 1948, Rev. Ent., vol. 19, p. 22.

Lectotype, female, from Ponta Grossa, Paraná, designated from the three original cotypes, all females.

justus, Meroption. 1947, Rev. Ent., vol. 18, p. 282.

Holotype, sex undetermined, from Ponta Grossa, Paraná, September, 1945. Unique specimen.

justus, Naupactus. 1949, Rev. Ent., vol. 20, p. 176, fig. 4.

Lectotype, male, from Ponta Grossa, Paraná, designated from two cotypes in original series of 11. Other cotypes stated by Bondar to be in Chicago.

justus, Stegotes. 1946, Rev. Ent., vol. 17, p. 322.

Lectotype, male, from Ponta Grossa, Paraná, designated from the three original cotypes.

kuscheli, Borgmeierus. 1945, Rev. Ent., vol. 16, p. 110.

Lectotype, male, from Rio Bravo, Entre Rios, Argentina, February 6, 1943, in *Cyperus giganteus*, designated from the eight original cotypes.

kuscheli, Megavallius. 1948, Rev. Ent., vol. 19, p. 28.

Holotype, female, from Porto Alegre, Rio Grande do Sul, October 15, 1936, labeled cotype in error. Unique specimen.

kuscheli, Thaliabaris. 1945, Rev. Ent., vol. 16, p. 105.

Lectotype, male, from Rio Bravo, delta of Rio Paraná, Argentina, February, 1943, in *Thalia multiflora*, designated from two cotypes in original series of five. Other cotypes stated by Bondar to be in collection of Kuschel.

lamellicornis, Saranthebaris. 1945, Rev. Ent., vol. 16, p. 317.

Lectotype, male, from Fazenda Bom Futuro, Belmonte, Bahia, October 20, 1944, in *Saranthe*, designated from 37 cotypes in original series of 65. Other cotypes stated by Bondar to be in London (B) and São Paulo (A).

landeiroi, Elytrocoptus. 1942, Rev. Ent., vol. 13, p. 235, fig. 14.

Lectotype, sex undetermined, from Vitoria, Espirito Santo, in *Psidium goyava*, designated from the three original cotypes.

landeiroi, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 456.

Lectotype, male, from Vitoria, Espirito Santo, June 5, 1941, in *Cocos picrophylla*, designated from 61 cotypes in original series of 280. Other cotypes stated by Bondar to be in Chicago and São Paulo (C). One cotype is in Washington (Warner, *in litt.*).

landeiroi, Rhyssomatus. 1942, Rev. Ent., vol. 13, p. 3, figs. 1-3.

Lectotype, probably male, from Agua Preta, Bahia, designated from two cotypes in original series of eight. Other cotypes, two of which are from the state of Espirito Santo, stated by Bondar to be in Chicago and São Paulo (A, B).

lanei, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 322.

Lectotype, sex undetermined, from Itabuna, Bahia, in Labiaceae, designated from one cotype in original series of two. Location of other cotype unknown.

lanei, Polpones. 1943, Rev. Ent., vol. 14, p. 51.

Lectotype, male, from Bahia, Bahia, in *Rhynchospora cephalotes*, designated from the three original cotypes. Two other cotypes are females.

langei, Celetes. 1943, Rev. Ent., vol. 14, p. 345.

Lectotype, male, from Curitiba, Paraná, designated from 24

cotypes in original series of 50. Location of other cotypes unknown. Fourteen of the 24 are from the state of Santa Catarina. **lapanus, Prodinus.** 1950, Rev. Ent., vol. 21, p. 463.

Lectotype, male, from valley of Rio São Francisco, Lapa, Bahia, October 20, 1947, designated from three cotypes in original series of six. Other cotypes stated by Bondar to be in Chicago.

lauracei, Meroprion. 1947, Rev. Ent., vol. 18, p. 282.

Lectotype, sex undetermined, from Dias Davila, Bahia, October 10, 1940, in Lauraceae, designated from two cotypes in original series of four. Location of other cotypes unknown.

leiospathae, Ancylorrhynchus. 1950, Rev. Ent., vol. 21, p. 454, fig. 7.

Lectotype, male, from Pirassununga, São Paulo, October, 1949, in *Cocos leiospatha*, designated from the two original cotypes. Lectotype is in Chicago.

leiospathae, Tripusus. 1950, Rev. Ent., vol. 21, p. 474.

Holotype, sex undetermined, from Pirassununga, São Paulo, October, 1949, in *Cocos leiospatha*, labeled cotype in error. Unique specimen.

lepagei, Sibariops. 1943, Rev. Ent., vol. 14, p. 70.

Lectotype, probably male, from Fazenda Pasto Fora, Bahia, Bahia, in Ciperaceae, designated from five cotypes in original series of eight. Location of other cotypes unknown.

levi, Sibariops. 1943, Rev. Ent., vol. 14, p. 74.

Lectotype, male, from Fazenda Pasto Fora, Bahia, Bahia, August 23, 1942, in Ciperaceae, designated from the 10 original cotypes.

lineatus, Ancylorrhynchus. 1943, Rev. Ent., vol. 14, p. 363.

Lectotype, male, from the state of Santa Catarina, in *Cocos Romanzoffiana*, designated from five cotypes in original series of 12. Location of other cotypes unknown.

lineatus, Borgmeierus. 1943, Rev. Ent., vol. 14, p. 57.

Lectotype, male, from the state of Bahia, in *Diclidium lenticulare*, designated from 30 cotypes in original series of 60. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B). Name of plant was later corrected to *Cyperus ferax* by Bondar.

littoralis, Geraeus. 1942, Rev. Ent., vol. 13, p. 248, figs. 16-18.

Lectotype, male, from Vila Velha, Espírito Santo, March, 1941, in *Araecoccus*, designated from 17 cotypes in original series

of 98. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B); 11 of the above 17 cotypes and the lectotype are in Chicago. We have 20 metatypes, labeled as cotypes, dated 1943.

longirostris, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 327.

Holotype, "probably female," from Itatiaia, Rio de Janeiro, labeled cotype in error. Unique specimen.

longulus, Thaliabaris. 1945, Rev. Ent., vol. 16, p. 107.

Lectotype, probably female, from Rio Bravo, Entre Rios, Argentina, February 6, 1943, in *Thalia multiflora*, designated from one cotype in original series of two. Location of other cotype unknown.

lunatus, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 46.

Lectotype, male, from Ponta Grossa, Paraná, designated from one cotype in original series of five. Other cotypes, two of which are from Tucuman, Argentina, stated by Bondar to be in Chicago and (entry in Bondar's notebook) in collection of Hustache.

maculatus, Peridinetus. 1949, Rev. Ent., vol. 20, p. 188.

Lectotype, sex undetermined, from Canavieiras, Bahia, October, 1946, in *Piper*, designated from two cotypes in original series of three. Location of other cotype unknown.

maculatus, Thaliabaris. 1945, Rev. Ent., vol. 16, p. 107.

Lectotype, male, from Rio Bravo, Entre Rios, Argentina, February 6, 1943, in *Thalia multiflora*, designated from six cotypes in original series of nine. Other cotypes stated by Bondar to be in collection of Kuschel.

malleri, Celetes. 1943, Rev. Ent., vol. 14, p. 344.

Lectotype, male, from the state of Santa Catarina, in *Cocos Romanzoffiana*, designated from two cotypes in original series of five. Location of other cotypes, one of which is from Curitiba, Paraná, March, 1942, unknown.

malleri, Linonotus. 1943, Rev. Ent., vol. 14, p. 126.

Holotype, male, from the state of Santa Catarina, labeled cotype in error. Unique specimen.

malleri, Liobaridia. 1950, Rev. Ent., vol. 21, p. 471.

Lectotype, male, from the state of Santa Catarina, designated from three cotypes in original series of six. Other cotypes stated by Bondar to be in Chicago.

malleri, Nertus. 1942, Rev. Ent., vol. 13, p. 266.

Lectotype, probably male, from Mafra, Santa Catarina,

December, 1941, designated from two cotypes in original series of three. Other cotype stated by Bondar to be in Chicago.

malleri, Palmocentrinus. 1949, Rev. Ent., vol. 20, p. 207.

Lectotype, sex undetermined, from the state of Santa Catarina, November, 1943, designated from three cotypes in original series of four. Other cotype stated by Bondar to be in Chicago.

malpighiaceae, Chalcodermus. 1949, Rev. Ent., vol. 20, p. 183.

Lectotype, female, from Andaraí, Bahia, October, 1948, in Malpighiaceae, designated from two cotypes in original series of three. Other cotype stated by Bondar to be in Chicago.

marantacei, Centrinus. 1943, Rev. Ent., vol. 14, p. 129, fig. 32.

Holotype, male, from Bahia, Bahia, February 3, 1929, in Marantaceae, labeled cotype in error. Unique specimen.

marginata, Hustachea. 1942, Rev. Ent., vol. 13, p. 34.

Lectotype, sex undetermined, from Santa Teresinha, Bahia, May, 1941, in *Cocos vagans*, designated from the two original cotypes.

mariae-helenae, Derelomus. 1941, Rev. Ent., vol. 12, p. 281, fig. 5.

Lectotype, probably male, from Florianópolis, Santa Catarina, September 10, 1940, in *Cocos eriospatha*, designated from eight cotypes in original series of 15. Two cotypes in Washington (Warner, *in litt.*). Location of other cotypes unknown. "Type and cotype in collection of author," but all specimens labeled cotypes.

maritimus, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 221.

Lectotype, male, from the state of Espírito Santo "near the coast," April, 1941, in *Diplothemium maritimum*, designated from 18 cotypes in original series of 21. Other cotypes stated by Bondar to be in Chicago.

marizae, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 452, fig. 16.

Lectotype, male, from Fazenda Babassú, São Mateus, Espírito Santo, June 12, 1941, in *Attalea compta*, designated from 25 cotypes in original series of 120. Other cotypes stated by Bondar to be in Chicago and São Paulo (C). One cotype in Washington (Warner, *in litt.*).

marshalli, Chalcodermus. 1929, Correio Agric., vol. 8, p. 126.

Lectotype, female, from the state of Bahia, designated from two cotypes in original series of 12. Other cotypes stated by Bondar to be in London (B).

marshalli, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 319.

Lectotype, sex undetermined, from Campo Formoso, Bahia, November 5, 1943, in *Anona coriacea*, designated from five cotypes in original series of 20. Other cotypes stated by Bondar to be in London (B) and São Paulo (A).

marshalli, Dinomorphus. 1950, Rev. Ent., vol. 21, p. 449.

Holotype, sex undetermined, from Feira de Santa Ana, Bahia, labeled cotype in error. Unique specimen.

marshalli, Linonotus. 1945, Rev. Ent., vol. 16, p. 316.

Lectotype, male, from the state of Santa Catarina, designated from the two original cotypes. Other cotype is female.

marshalli, Scymnoplastophilus. 1938, Rev. Ent., vol. 8, p. 11.

Holotype, sex undetermined, from Agua Preta, Bahia, in *Guarea*, labeled cotype in error. Three paratypes. Holotype is first specimen on pin (Bondar, *in litt.*). About 48 paratypes stated by Bondar to be in Berlin, London (A), and Rio de Janeiro (D).

massarandubae, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 202.

Lectotype, male, from Agua Preta, Bahia, in Sapotaceae, designated from the two original cotypes.

mendonca, Spermologus. 1943, Rev. Ent., vol. 14, p. 100.

Lectotype, male, from Campo Formoso, Bahia, April, 1943, in *Anona coriacea*, designated from four cotypes in original series of 26. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

metallescens, Baptobarinus. 1949, Rev. Ent., vol. 20, p. 190.

Lectotype, sex undetermined, from Agua Preta, Bahia, designated from three cotypes in original series of four. Other cotype stated by Bondar to be in Chicago.

metallica, Willinkia. 1949, Rev. Ent., vol. 20, p. 191, figs. 6-7.

No lectotype designated. Species described from seven specimens from Vila Guilhermina, Tucuman, Argentina, 1944 and 1945. Our four specimens, labeled cotypes, have dates of February 16-25, 1948, and must be metatypes. Two specimens from Chicago, labeled cotypes, are from Villa Ana, Santa Fé, Argentina, February 4-18, 1946. Some cotypes stated by Bondar to be in Argentina.

mimica, Bondariella.¹ 1942, Rev. Ent., vol. 13, p. 21, figs. 4-7.

Lectotype, male, from Bahia, Bahia, in *Cocos schizophylla*, designated from 123 cotypes in original series of "about 400."

¹ Described by Hustache and Bondar.

Other cotypes stated by Bondar to be in Chicago, São Paulo (B), and in collection of Hustache.

mimosae, Spermobaris. 1949, Rev. Ent., vol. 20, p. 201, fig. 9.

Lectotype, male, from Curitiba, Paraná, in *Mimosa*, designated from three cotypes in original series of five. Other cotypes stated by Bondar to be in Chicago.

minimus, Ancylorrhynchus. 1950, Rev. Ent., vol. 21, p. 455, fig. 8.

Lectotype, female, from Corupá, Santa Catarina, designated from two cotypes in original series of three. One cotype is in Chicago; location of other unknown.

minimus, Anopsilus. 1942, Rev. Ent., vol. 13, p. 257.

Holotype, sex undetermined, from the state of Bahia, in *Canna denudata*. Unique specimen.

minor, Phytotribus. 1941, Rev. Ent., vol. 12, p. 447, fig. 13.

Lectotype, male, from São Mateus, Espírito Santo, June 12, 1941, in *Attalea compta*, designated from six cotypes in original series of 14. Other cotypes stated by Bondar to be in Chicago and São Paulo (C). Locality of cotypes was not given in description but was taken from entry #2852 in Bondar's notebook.

modicus, Linonotus. 1948, Rev. Ent., vol. 19, p. 25.

Holotype, female, from Tucuman, Argentina. Unique specimen.

montei, Linonotus. 1945, Rev. Ent., vol. 16, p. 315.

Lectotype, male, from Fazenda Bom Futuro, Belmonte, Bahia, October 20, 1944, in *Saranthe*, designated from 29 cotypes in original series of 80. Other cotypes stated by Bondar to be in London (B) and São Paulo (A).

montei, Metriophilus. 1942, Rev. Ent., vol. 13, p. 5.

Lectotype, probably male, from Santa Teresinha, Bahia, in Leguminosae, designated from five cotypes in original series of 30. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B).

montei, Sibariops. 1943, Rev. Ent., vol. 14, p. 76.

Lectotype, male, from Fazenda Pasto Dentro, Bahia, Bahia, designated from five cotypes in original series of eight. Other cotypes stated by Bondar to be in Chicago and São Paulo (B). Locality of cotypes was not given in description but was taken from entry #2957 in Bondar's notebook.

morei, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 324.

Lectotype, probably male, from Itu, São Paulo, February,

1939, designated from two cotypes in original series of six. Location of other cotypes unknown. Locality taken from specimens; none given in description.

mourei, Erirrhinoides. 1943, Rev. Ent., vol. 14, p. 350.

Lectotype, male, from Curitiba, Paraná, designated from two cotypes in original series of three. Location of other cotype unknown.

mourei, Pseudocentrinus. 1943, Rev. Ent., vol. 14, p. 368.

Lectotype, male, from Curitiba, Paraná, March, 1942, in *Cocos Romanzoffiana*, designated from the 10 original cotypes, five of which are from the state of Santa Catarina.

mucugeana, Bondariella. 1949, Rev. Ent., vol. 20, p. 208.

Lectotype, male, from Mucugê, Bahia, October, 1948, in *Cocos*, designated from 286 cotypes in original series of "about 350." Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

mundururu, Teratonychus. 1949, Rev. Ent., vol. 20, p. 186, figs. 15-19.

Lectotype, male, from Andaraí, Bahia, October, 1948, in Melastomaceae, designated from two cotypes in original series of five. Other cotypes stated by Bondar to be in Chicago.

murtinus, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 198.

Lectotype, male, from Ilha dos Frades, Bahia, in Mirtaceae, designated from 11 cotypes in original series of 35. Other cotypes stated by Bondar to be in London (B).

mutilatus, Bignonibaris. 1946, Rev. Ent., vol. 17, p. 110, figs. 13-14.

Lectotype, sex undetermined, from Ponta Grossa, Paraná, in *Anemopaegma*, designated from 20 cotypes in original series of 70. Lectotype and cotypes in Chicago. Our 160 specimens, same locality, probably contain representatives of original series, but bear no cotype labels. Some cotypes stated by Bondar to be in collection of Hustache.

nasutus, Sibariops. 1943, Rev. Ent., vol. 14, p. 75.

Lectotype, male, from Bahia, Bahia, in *Rhynchospora cephalotes*, designated from 32 cotypes in original series of 40. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

nicki, Linonotus. 1948, Rev. Ent., vol. 19, p. 24.

Lectotype, male, from Jabaquara, São Paulo, October 29, 1941, designated from the two original cotypes.

niger, Thaliabaris. 1945, Rev. Ent., vol. 16, p. 108.

Lectotype, male, from Rio Bravo, Entre Rios, Argentina, February 6, 1943, in *Thalia multiflora*, designated from the two original cotypes. Other cotype is a female.

nigriventris, Stegotes. 1946, Rev. Ent., vol. 17, p. 327.

Lectotype, male, from the state of Santa Catarina, designated from two cotypes in original series of three. Location of other cotype unknown.

oiti, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 203.

Holotype, "evidently male," from Agua Preta, Bahia, in *Conepia excelsa*, labeled cotype in error. Unique specimen.

olyrae, Lispodemus. 1942, Rev. Ent., vol. 13, p. 262.

Lectotype, male, from Bahia, Bahia, April 1-12, 1942, in *Olyra latifolia*, designated from five cotypes in original series of 16. Other cotypes stated by Bondar to be in Buenos Aires, Chicago, Rio de Janeiro (A), and São Paulo (A).

oncidi, Montella. 1948, Rev. Ent., vol. 19, p. 35.

Holotype, male, from Curitiba, Paraná, October, 1943, in *Oncidium concolor*. One paratype, female.

orlandoi, Linonotus. 1948, Rev. Ent., vol. 19, p. 22.

Lectotype, male, from Fazenda Terra Firme, Canavieiras, Bahia, October 4, 1946, in *Sarante*, designated from the two original cotypes.

pallidus, Celetes. 1943, Rev. Ent., vol. 14, p. 346.

Lectotype, male, from the state of Paraná, in *Cocos*, designated from the two original cotypes. Other cotype, a female, is from state of Santa Catarina.

palmarum, Phytotribus. 1943, Rev. Ent., vol. 14, p. 356.

Lectotype, male, from the state of Espirito Santo, in *Attalea*, designated from two cotypes in original series of four. Location of other cotypes unknown.

paranaensis, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 222.

Lectotype, male, from Ponta Grossa, Paraná, designated from 19 cotypes in original series of 28. Other cotypes stated by Bondar to be in Bahia and Chicago.

paranaensis, Stegotes. 1946, Rev. Ent., vol. 17, p. 318.

Holotype, "apparently female," from Curitiba, Paraná, labeled cotype in error. Unique specimen.

partitus, Diorymerus. 1950, Rev. Ent., vol. 21, p. 462.

Holotype, sex undetermined, from Virginia, Minas Gerais, 1500 meters, November 25, 1914, labeled cotype in error. "Type and paratype in collection of author"; location of paratype, dated October 11, 1915, unknown.

paspali, Coelonertinus. 1949, Rev. Ent., vol. 20, p. 226.

Lectotype, male, from Porto Alegre, Rio Grande do Sul, January, 1948, in *Paspalum urvillei*, designated from 42 cotypes in original series of 47. Other cotypes stated by Bondar to be in Chicago.

paulistanus, Megavallius. 1948, Rev. Ent., vol. 19, p. 27.

Lectotype, male, from Itatiaia, Rio de Janeiro, designated from two cotypes in original series of four. Location of other cotypes unknown.

pedrito, Sibariops. 1943, Rev. Ent., vol. 14, p. 70.

Lectotype, sex undetermined, from Fazenda Pasto Fora, Bahia, Bahia, August 23, 1942, in Ciperaceae, designated from nine cotypes in original series of 13. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

pedritosilvai, Sibariops. 1943, Rev. Ent., vol. 14, p. 72.

Lectotype, probably male, from Agua Preta, Ilheos, Bahia, August, 1942, in *Diclidium lenticulare*, designated from five cotypes in original series of 12. Other cotypes stated by Bondar to be in Chicago, Rio de Janeiro (A), and São Paulo (A).

perseae, Prionopus. 1937, Rev. Ent., vol. 7, p. 480, figs. 3-4.

Holotype, sex undetermined, from Agua Preta, Bahia, December 30, 1936, in *Persea gratissima*, labeled cotype in error. "Type in collection of author." One paratype. Other paratypes stated by Bondar to be in Berlin and London (A).

pilosiventris, Megavallius. 1948, Rev. Ent., vol. 19, p. 26.

Holotype, male, from Jequié, Bahia, December, 1922, in Marantaceae, labeled cotype in error. Unique specimen.

pindobae, Phytotribus. 1941, Rev. Ent., vol. 12, p. 293, fig. 10.

Lectotype, male, from Fazenda Copiobá, Nazaré, Bahia, in *Attalea humilis*, designated from four cotypes in original series of 13. Location of other cotypes unknown. "Type and cotype in collection of author," but all specimens labeled cotypes.

piperi, Cyrionyx. 1943, Rev. Ent., vol. 14, p. 105, figs. 17-19.

Lectotype, sex undetermined, from Bahia, Bahia, March 8, 1942, in *Piper*, designated from seven cotypes in original series of 20. Location of other cotypes unknown.

pitubae, Conotrachelus. 1945, Rev. Ent., vol. 16, p. 320.

Lectotype, sex undetermined, from Pituba, Bahia, in Sapotaceae, designated from three cotypes in original series of 15. Other cotypes stated by Bondar to be in London (B) and São Paulo (A).

planithorax, Hoplorrhinus. 1941, Rev. Ent., vol. 12, p. 288.

Lectotype, male, from Jequié, Bahia, in *Attalea compta*, designated from 21 cotypes. Original series was "about 10." "Type and cotype in collection of author," but all labeled cotypes. Some of our cotypes may be metatypes, but it is impossible to know where the original series ends.

plaumanni, Linonotus. 1943, Rev. Ent., vol. 14, p. 123.

Holotype, female, from Nova Teutonia, Santa Catarina. Unique specimen.

plaumanni, Lispodemus. 1950, Rev. Ent., vol. 21, p. 475, fig. 16.

Lectotype, sex undetermined, from Nova Teutonia, Santa Catarina, designated from the two original cotypes.

plaumanni, Revena. 1943, Rev. Ent., vol. 14, p. 383.

Lectotype, male, from Nova Teutonia, Santa Catarina, designated from the two original cotypes. Other cotype a female.

polyphaga, Dialomia. 1942, Rev. Ent., vol. 13, p. 14.

Lectotype, male, from environs of Bahia, Bahia, in *Attalea junifera*, designated from nine cotypes in original series of 16. Six of the nine are from state of Bahia. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

psychotriæ, Abrachybaris. 1946, Rev. Ent., vol. 17, p. 105.

Lectotype, sex undetermined, from Belmonte, Bahia, in Rubiaceae, designated from five cotypes in original series of 20. Other cotypes stated by Bondar to be in Chicago and in collection of Hustache.

ramosi, Sibariops. 1943, Rev. Ent., vol. 14, p. 67.

Lectotype, male, from Pituba, Bahia, designated from 30 cotypes in original series of 50. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

romanzoffianæ, Phytotribus. 1943, Rev. Ent., vol. 14, p. 355.

Lectotype, male, from Nova Teutonia, Santa Catarina, in *Cocos Romanzoffiana*, designated from 12 cotypes in original series of 17. Location of other cotypes unknown.

ruber, Stegotes. 1946, Rev. Ent., vol. 17, p. 326.

Holotype, "evidently female," from Hansa [now Corupá], Santa Catarina, December, 1942. Unique specimen.

rubicundus, Linonotus. 1948, Rev. Ent., vol. 19, p. 24.

Holotype, female, from Corupá, Santa Catarina, labeled cotype in error. Unique specimen.

rufa, Euterpia. 1942, Rev. Ent., vol. 13, p. 25.

Lectotype, sex undetermined, from Nazaré, Bahia, November

20, 1940, in *Euterpe edulis*, designated from four cotypes in original series of eight. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B). Locality of cotypes was not given in description, but was taken from entry #2751 in Bondar's notebook.

ruficollis, Lydamis. 1946, Rev. Ent., vol. 17, p. 89.

Lectotype, male, from the state of Paraná, designated from two cotypes in original series of 12. Location of other cotypes unknown.

rufipennis, Sibariops. 1943, Rev. Ent., vol. 14, p. 73.

Lectotype, male, from Fazenda Pasto Fora, Bahia, Bahia, August 23, 1942, designated from 10 cotypes in original series of 16. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

rufipes, Montebaris. 1949, Rev. Ent., vol. 20, p. 193.

Lectotype, male, from Curitiba, Paraná, designated from the two original cotypes.

rufipes, Montella. 1948, Rev. Ent., vol. 19, p. 34.

Lectotype, male, from Florestal, Rio de Janeiro, October, 1944, in *Sternorhynchus australis*, designated from eight cotypes in original series of 11. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

rufirostris, Bromesia. 1946, Rev. Ent., vol. 17, p. 337.

Lectotype, probably male, from Inhambupé, Bahia, February, 1943, in *Araecoccus*, designated from five cotypes in original series of six. Location of other cotype unknown.

rufirostris, Loboderinus. 1948, Rev. Ent., vol. 19, p. 43, figs. 26-27, 30.

Lectotype, male, from the state of Bahia, designated from 80 cotypes in original series of 102. Other cotypes stated by Bondar to be in Chicago, London (A), and Rio de Janeiro (A).

rufomaculatus, Mesothanius. 1949, Rev. Ent., vol. 20, p. 199.

Lectotype, male, from the state of Minas Gerais, designated from four cotypes in original series of five, two of the four being from Morumbi, São Paulo, January 25, 1945. Other cotype from Morumbi stated by Bondar to be in Chicago.

rufus, Strongylothes. 1942, Rev. Ent., vol. 13, p. 269.

Holotype, female, from Nazaré, Bahia, in *Andropogon condensatus*, labeled cotype in error. Unique specimen.

rugicollis, Telemus. 1950, Rev. Ent., vol. 21, p. 458.

Holotype, sex undetermined, from Morumbi, São Paulo. Unique specimen.

ruschiana, Bondariella. 1942, Rev. Ent., vol. 13, p. 24.

Lectotype, male, from Colatina, Espirito Santo, June, 1941, in *Cocos ruschiana*, designated from 149 cotypes in original series of "about 200." Other cotypes stated by Bondar to be in Chicago and São Paulo (B).

ruschianae, Ancylorrhynchos. 1941, Rev. Ent., vol. 12, p. 468, figs. 29-31.

This species is not present in Bondar's collection. Described from two specimens from Colatina, Espirito Santo, June 17, 1941, in *Cocos ruschiana*.

sampsoni, Ceratopus. 1947, Rev. Ent., vol. 18, p. 288.

Lectotype, male, from Bahia, Bahia, in *Ficus*, designated from one cotype in original series of two. Location of other cotype (male) unknown.

saranthae, Loboderes. 1948, Rev. Ent., vol. 19, p. 40, figs. 23-24.

Holotype, sex undetermined, from Belmonte, Bahia, October 20, 1944, in *Saranthae*. One paratype.

schubarti, Angelocentris. 1950, Rev. Ent., vol. 21, p. 473.

Lectotype, male, from Pirassununga, São Paulo, in *Cocos leiospatha*, designated from four cotypes in original series of six. Other cotypes stated by Bondar to be in Chicago and in collection of Borgmeier.

scleriae, Polpones. 1943, Rev. Ent., vol. 14, p. 51.

Lectotype, male, from Itatiaia, Rio de Janeiro, in *Scleria bracteata*, designated from the two original cotypes. Other cotype (female) is from Fazenda Copiobá, Nazaré, Bahia.

scleriae, Torcus. 1943, Rev. Ent., vol. 14, p. 79.

Lectotype, male, from Bahia, Bahia, in *Scleria bracteata*, designated from eight cotypes in original series of 10. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

scutellatus, Polpones. 1943, Rev. Ent., vol. 14, p. 52.

Lectotype, sex undetermined, from the state of Santa Catarina, designated from the two original cotypes.

semirufus, Diorymerus. 1950, Rev. Ent., vol. 21, p. 461.

Holotype, sex undetermined, from São Gonçalo dos Campos, Bahia, January, 1946, in Malpighiaceae. Unique specimen.

serjaniae, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 47.

Lectotype, male, from Soure, Bahia, June 20, 1946, in *Serjania*, designated from 11 cotypes in original series of 22. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

sexgibbosus, Prionomerus. 1938, Rev. Ent., vol. 8, p. 5, fig. 4.

Holotype, sex undetermined, from Agua Preta, Bahia, in *Persea gratissima*, labeled cotype in error. Unique specimen.

signatus, Glyptobaris. 1946, Rev. Ent., vol. 17, p. 98.

Lectotype, sex undetermined, from Jacaranda, Canavieiras, Bahia, March, 1945, in Convolvulaceae, designated from the two original cotypes.

signatus, Gravatageraeus. 1942, Rev. Ent., vol. 13, p. 251, figs. 20-21.

Lectotype, male, from Bahia, Bahia, in *Hohenbergia*, designated from six cotypes in original series of 14. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B). We have five metatypes, labeled cotypes, from Pituba, Bahia.

silvai, Glyptobaroides. 1949, Rev. Ent., vol. 20, p. 194.

Lectotype, possibly female, from Guaratiba, Distrito Federal, Rio de Janeiro, October, 1940, designated from five cotypes in original series of eight. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

similis, Craspedotus. 1948, Rev. Ent., vol. 19, p. 13, figs. 12-14.

Lectotype, male, from Belmonte, Bahia, designated from 18 cotypes in original series of 27. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

similis, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 458.

Holotype, female, from São Mateus, Espírito Santo, April 12, 1941, in *Attalea compta*. Unique specimen.

similis, Sibariops. 1943, Rev. Ent., vol. 14, p. 64.

Lectotype, male, from Pituba, Bahia, March to September, 1942, in *Fuirena umbellata*, designated from 32 cotypes in original series of "about 50." Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

sincora, Megalobaris. 1949, Rev. Ent., vol. 20, p. 195.

Holotype, possibly male, from Serra do Sincorá, Mucugê, interior of Bahia, October, 1948. Unique specimen.

sparsepilosus, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 48.

Lectotype, probably male, from Soure, Bahia, June, 1946, designated from three cotypes in original series of seven. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

spiniger, Stegotes. 1946, Rev. Ent., vol. 17, p. 320.

Lectotype, male, from Itatiaia, Rio de Janeiro, designated from the three original cotypes.

spinus, Hoplorhinus. 1941, Rev. Ent., vol. 12, p. 457.

Holotype, male, from the state of Espirito Santo, in *Diplothemium caudescens*. Unique specimen.

splendidus, Eurhinus. 1948, Rev. Ent., vol. 19, p. 19, figs. 17-18.

Lectotype, male, from the state of Minas Gerais, in *Cissus cicyoides*, designated from 13 cotypes in original series of 24. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

squamosus, Phytotribus. 1943, Rev. Ent., vol. 14, p. 354.

Lectotype, male, from Nova Teutonia, Santa Catarina, in *Cocos*, designated from five cotypes in original series of 11. Location of other cotypes unknown.

squamosus, Stegotes. 1946, Rev. Ent., vol. 17, p. 319.

Holotype, "probably male," from the state of Santa Catarina. Unique specimen.

squamulosus, Phytotribus. 1943, Rev. Ent., vol. 14, p. 354.

Lectotype, male, from Curitiba, Paraná, in *Cocos*, designated from 10 cotypes in original series of 15. Location of other cotypes unknown.

sternicornis, Anopsilus. 1942, Rev. Ent., vol. 13, p. 256.

Lectotype, male, from Bahia, Bahia, in *Canna flacida* or *C. denudata*, designated from 10 cotypes in original series of 20. Other cotypes stated by Bondar to be in Chicago and São Paulo (A). Locality of cotypes was not given in description but was taken from entry #1113 in Bondar's notebook.

sternicornis, Derelomus. 1941, Rev. Ent., vol. 12, p. 277, figs. 2-3.

Lectotype, male, from Fazenda Copiobá, Nazaré, Bahia, November 20, 1940, in *Diplothemium caudescens*, designated from 29 cotypes in original series of "hundreds of specimens." Twenty-one cotypes in Washington (Warner, *in litt.*), location of other cotypes unknown. "Type and paratype in collection of author," but all specimens labeled cotypes.

stigmatophylli, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 49.

Lectotype, sex undetermined, from Itaberaba, Bahia, July, 1947, in *Stigmatophyllum*, designated from two cotypes in original series of three. Other cotype stated by Bondar to be in Chicago.

striatipennis, Anopsilus. 1942, Rev. Ent., vol. 13, p. 255.

Lectotype, male, from Bahia, Bahia, in *Canna denudata*, des-

ignated from two cotypes in original series of five. Other cotypes stated by Bondar to be in Chicago.

stromantheae, Linonotus. 1943, Rev. Ent., vol. 14, p. 124, figs. 29–30.

Lectotype, male, from Bahia, Bahia, July 5, 1942, in *Stromanthe*, designated from the four original cotypes.

subglaber, Strongylotes. 1949, Rev. Ent., vol. 20, p. 224.

Lectotype, male, from Tucuman, Argentina, designated from the two original cotypes. Other cotype is from Misiones, Argentina.

suturalis, Palmoderes. 1948, Rev. Ent., vol. 19, p. 38, figs. 21–22, 28.

Lectotype, sex undetermined, from Corupá, Santa Catarina, October, 1943, designated from 12 cotypes in original series of 15. Other cotypes stated by Bondar to be in Chicago.

tatianae, Derelomus. 1941, Rev. Ent., vol. 12, p. 280.

Lectotype, sex undetermined, from Nazaré, Bahia, in *Attalea funifera*, designated from 37 cotypes in original series of "numerous specimens." Twenty-one cotypes in Washington (Warner, *in litt.*); location of other cotypes unknown. "Type and paratype in collection of author," but all specimens labeled cotypes.

tenebrosus, Gladosius. 1946, Rev. Ent., vol. 17, p. 334.

Holotype, "probably female," from the state of Minas Gerais, January, 1925. Unique specimen.

tenuirostris, Eurypages. 1946, Rev. Ent., vol. 17, p. 92.

Holotype, "probably female," from Itatiaia, Rio de Janeiro. Unique specimen.

thaliae, Thaliabaris. 1943, Rev. Ent., vol 14, p. 133.

Lectotype, male, from Fazenda Bom Gosto, Santo Amaro, Bahia, March 4, 1943, in *Thalia geniculata* or *T. angustifolia*, designated from 75 cotypes in original series of 120. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

tibialis, Stegotes. 1949, Rev. Ent., vol. 20, p. 204.

Holotype, sex undetermined, from Maracás, Bahia, January, 1945, in Bromeliaceae. Unique specimen.

tiririca, Atorcus. 1943, Rev. Ent., vol. 14, p. 58.

Lectotype, male, from environs of Bahia, in *Scleria bracteata*, designated from 43 cotypes in original series of 52. Eleven of the 43 cotypes in Chicago. Other cotypes stated by Bondar to be in Rio de Janeiro (A) and São Paulo (A). We have 16 additional specimens labeled cotypes which, since they are dated March

9, 1943, only four months prior to publication of the description, are no doubt metatypes.

tiririca, Polpones. 1943, Rev. Ent., vol. 14, p. 52.

Holotype, sex undetermined, from Ilheos, Bahia, in Ciperaceae. Unique specimen.

titara, Dahlgrenia. 1942, Rev. Ent., vol. 13, p. 36.

Lectotype, sex undetermined, from Rio Braço do Norte, São Mateus, Espirito Santo, June, 1941, in *Desmoncus polyacanthos*, designated from 173 cotypes in original series of 410. Other cotypes stated by Bondar to be in Chicago and São Paulo (A, B).

torresi, Bondariella. 1942, Rev. Ent., vol. 13, p. 22.

Lectotype, male, from Lagedo Alto, district of Santa Luzia, Bahia, in *Cocos vagans*, designated from 248 cotypes in original series of "about 300." Other cotypes stated by Bondar to be in Chicago and São Paulo (B). Some of these others may have locality of Santa Tereza, Bahia.

trapia, Craspedotus. 1948, Rev. Ent., vol. 19, p. 14.

Lectotype, probably female, from Tucano, Bahia, December, 1946, designated from 24 cotypes in original series of 45. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

trithrinacis, Celetes. 1943, Rev. Ent., vol. 14, p. 348.

Lectotype, male, from Tucuman, Argentina, in *Trithrinax campestris*, designated from four cotypes in original series of nine. Location of other cotypes unknown.

tuberosus, Justus. 1946, Rev. Ent., vol. 17, p. 95.

Lectotype, male, from Corupá, Santa Catarina, designated from the two original cotypes.

tucum, Parisoschoenus. 1949, Rev. Ent., vol. 20, p. 219.

Lectotype, male, from Curitiba, Paraná, March, 1942, designated from 41 cotypes (16 from Curitiba, 25 from Bahia, Bahia, same date) in original series of 50. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

tucumanensis, Stegotes. 1946, Rev. Ent., vol. 17, p. 324.

Lectotype, male, from Tucuman, Argentina, designated from the two original cotypes.

ubacahy, Conotrachelus. 1944, Rev. Ent., vol. 15, p. 196.

Lectotype, male, from Santa Fé, Argentina, December, 1940, in *Eugenia edulis*, designated from three cotypes in original series of six. Location of other cotypes unknown.

vagans, Dialomia. 1942, Rev. Ent., vol. 13, p. 18.

Lectotype, male, from the interior of the state of Bahia, in *Cocos vagans*, designated from 11 cotypes in original series of 16. Other cotypes stated by Bondar to be in Chicago and São Paulo (A).

vagans, Revena. 1943, Rev. Ent., vol. 14, p. 382, figs. 1-7.

Lectotype, male, from Bahia, Bahia, in *Cocos vagans*, designated from the 20 original cotypes.

variabilis, Parallelosomus. 1943, Rev. Ent., vol. 14, p. 43.

Lectotype, male, from Pituba, Bahia, August 15, 1942, in *Rhynchospora cyperoides*, designated from seven cotypes in original series of 16. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A).

versicolor, Thaliabaris. 1945, Rev. Ent., vol. 16, p. 106.

Lectotype, male, from Rio Bravo, Entre Rios, Argentina, February 6, 1943, in *Thalia multiflora*, designated from seven cotypes in original series of nine. Other cotypes stated by Bondar to be in collection of Kuschel.

vochysiae, Chalcodermus. 1948, Rev. Ent., vol. 19, p. 51.

Lectotype, probably male, from the state of Minas Gerais, July 7, 1946, in Vochysiaceae, designated from one cotype in original series of four. Other cotypes stated by Bondar to be in Chicago.

vossi, Piazorhinus. 1938, Rev. Ent., vol. 8, p. 14, fig. 14.

Holotype, sex undetermined, from Agua Preta, Bahia, in *Coccoloba*, labeled cotype in error. "Type in collection of author." One paratype. Thirteen paratypes stated by Bondar to be in Berlin, London (A), and Rio de Janeiro (D, E).

willinki, Eurhinus. 1948, Rev. Ent., vol. 19, p. 20.

Lectotype, male, from Tucuman, Argentina, January 21, 1945, designated from one cotype in original series of two. Location of other cotype unknown.

willinki, Trachybaris. 1950, Rev. Ent., vol. 21, p. 468.

Lectotype, sex undetermined, from Paraguay, January 1, 1946, in Aristolochiaceae, designated from one cotype in original series of three. Other cotypes stated by Bondar to be in Chicago. Date in description was given as 1949, a typographical error.

xixim, Homalocerus. 1947, Rev. Ent., vol. 18, p. 277.

Lectotype, sex undetermined, from Ponta Grossa, Paraná, designated from the two original cotypes.

yvensi, Chalcodermus. 1941, Rev. Ent., vol. 12, p. 429, figs. 2-6.

Lectotype, sex undetermined, from Vitoria, Espirito Santo, June 9, 1941, in *Phaseolus vulgaris*, designated from eight cotypes in original series of 45. Other cotypes stated by Bondar to be in Chicago and São Paulo (C). One cotype in Washington (Warner, *in litt.*).

zikani, Anopsilus. 1942, Rev. Ent., vol. 13, p. 258.

Holotype, "probably female," from Itatiaia, Maromba, 1100 meters, Rio de Janeiro, October 8, 1931. Unique specimen.

zikani, Gladosius. 1942, Rev. Ent., vol. 13, p. 246.

Holotype, sex undetermined, from Itatiaia, Rio de Janeiro, November 26, 1929, in Bromeliaceae, labeled cotype in error. One paratype from Ilheos, Bahia, May 25, 1942.

zikani, Homalocerus. 1947, Rev. Ent., vol. 18, p. 276.

Holotype, sex undetermined, from the state of Rio de Janeiro. Unique specimen.

zikani, Linonotus. 1943, Rev. Ent., vol. 14, p. 126.

Holotype, "probably female," from Itatiaia, Rio de Janeiro, December 21, 1933, labeled cotype in error. Unique specimen.

zikani, Lispedemus. 1942, Rev. Ent., vol. 13, p. 263.

Lectotype, sex undetermined, from Fazenda dos Campos, Virginia, Minas Gerais, 1500 meters, December 12, 1919, designated from one cotype in original series of two. Location of other cotype unknown.

zikani, Orissus. 1946, Rev. Ent., vol. 17, p. 330.

Lectotype, male, from Itatiaia, Rio de Janeiro, February, 1931, designated from three cotypes in original series of four. Location of other cotype unknown.

zikani, Parallelosomus. 1943, Rev. Ent., vol. 14, p. 43.

Holotype, female, from the state of Rio de Janeiro, labeled cotype in error. Unique specimen.

zikani, Sibariops. 1943, Rev. Ent., vol. 14, p. 62.

Holotype, male, from São Gabriel, Rio Negro, Amazonas, November 9, 1927, labeled cotype in error. Unique specimen.

zikani, Strongylothes. 1949, Rev. Ent., vol. 20, p. 223.

Lectotype, male, from the state of Minas Gerais, January, 1947, designated from 27 cotypes in original series of 39. Other cotypes stated by Bondar to be in Chicago and Rio de Janeiro (A). The state of Rio de Janeiro, given as the locality in the original description, is an error; the specimens were merely sent by Zikan who lives in Rio de Janeiro (Bondar, *in litt.*).

