

# REVISION OF THE PLANT BUG GENUS *CORIDROMIUS* SIGNORET (INSECTA: HETEROPTERA: MIRIDAE)

NIKOLAI J. TATARNIC

*School of Biological Sciences  
University of Sydney  
Department of Entomology  
Australian Museum*

GERASIMOS CASSIS

*School of Biological, Earth and Environmental Sciences  
University of New South Wales*

BULLETIN OF THE AMERICAN MUSEUM OF NATURAL HISTORY  
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, NY 10024  
Number 315, 95 pp., 21 figures, 8 maps, 2 tables  
Issued August 29, 2008

## CONTENTS

Abstract . . . . .	3
Introduction . . . . .	3
Materials and Methods . . . . .	3
Specimens . . . . .	3
Homology and Terminology . . . . .	7
Measurements . . . . .	7
Systematic Placement . . . . .	7
Biology . . . . .	17
Habitats and Microhabitats . . . . .	17
Host Plant Associations . . . . .	19
Sexual Behavior . . . . .	20
Taxonomy . . . . .	20
<i>Coridromius</i> . . . . .	20
Key to Species of <i>Coridromius</i> . . . . .	25
<i>Coridromius bicolor</i> . . . . .	27
<i>Coridromius boianotum</i> . . . . .	31
<i>Coridromius bulbopella</i> . . . . .	32
<i>Coridromius carinatus</i> . . . . .	33
<i>Coridromius chenopoderis</i> . . . . .	35
<i>Coridromius chinensis</i> . . . . .	45
<i>Coridromius confusus</i> . . . . .	48
<i>Coridromius crassus</i> . . . . .	50
<i>Coridromius declivipennis</i> . . . . .	52
<i>Coridromius drepanopennis</i> . . . . .	53
<i>Coridromius ephippius</i> . . . . .	55
<i>Coridromius epithema</i> . . . . .	58
<i>Coridromius falsicoleus</i> . . . . .	59
<i>Coridromius hermosus</i> . . . . .	61
<i>Coridromius lestoni</i> . . . . .	63
<i>Coridromius marmoreus</i> . . . . .	64
<i>Coridromius minusculus</i> . . . . .	65
<i>Coridromius monotocopsis</i> . . . . .	67
<i>Coridromius nakatanii</i> . . . . .	71
<i>Coridromius neoguineanus</i> . . . . .	72
<i>Coridromius nigrus</i> . . . . .	73
<i>Coridromius pilbarensis</i> . . . . .	74
<i>Coridromius prolixipenis</i> . . . . .	77
<i>Coridromius pteraulos</i> . . . . .	78
<i>Coridromius punctatus</i> . . . . .	79
<i>Coridromius ruwenzorii</i> . . . . .	82
<i>Coridromius schuhi</i> . . . . .	82
<i>Coridromius sommelieri</i> . . . . .	82
<i>Coridromius tahitiensis</i> . . . . .	85
<i>Coridromius testaceous</i> . . . . .	87
<i>Coridromius thalame</i> . . . . .	89
<i>Coridromius variegatus</i> . . . . .	89
<i>Coridromius zetteli</i> . . . . .	92
Discussion . . . . .	93
Acknowledgments . . . . .	94
References . . . . .	94

## ABSTRACT

The plant bug genus *Coridromius* Signoret is revised. Twenty-one new species from Australia, the South Pacific, Asia and Africa are described (*bicolor*, *boianotum*, *bulbopella*, *chenopoderis*, *confusus*, *crassus*, *drepanopenis*, *ephippius*, *epithema*, *falsicoleus*, *hermosus*, *lestoni*, *marmoreus*, *monotocopsis*, *pilbarensis*, *prolixipenis*, *pteraulos*, *ruwenzorii*, *sommelieri*, *tahitiensis*, and *thalame*). Ten previously described species are redescribed (*schuhi* is excluded, as the type material is presumed lost and Linnauvori's original description and illustrations [Linnauvori, 1994] are not sufficient to distinguish *schuhi* from most other species). The genus *Coridromoides* Carvalho is synonymized with *Coridromius*, and its sole species, *Coridromoides carinatus* Carvalho, 1956, is transferred to *Coridromius*. A key to species, color habitus figures, illustrations of male genitalia, photographs of paragenital dissections, and scanning electron micrographs of structural details are provided. Detailed locality information and distribution maps of all species are included. Host plant data is provided for some species.

## INTRODUCTION

*Coridromius* Signoret (Heteroptera: Miridae: Orthotylinae: Halticinae) is an enigmatic plant bug genus readily recognized by its stout body and muscular jumping metathorax, which typically exhibit dark brown diagonal striping on their outer surface (figs. 1–3). This genus is noteworthy because it is the only plant bug known to engage in traumatic insemination, analogous to that found in the Cimicoidea (Tatarnic et al., 2006).

The genus was originally described by Montrouzier (1861) under the name *Ocyphus*, for a single species from New Caledonia (*O. variegatus*). Signoret (1862) recognized this as a junior homonym of the coleopteran genus *Ocyphus* Leach and proposed *Coridromius* as a replacement name.

*Coridromius* is found throughout the Old World tropics and subtropics, where 11 species have been described thus far. Aside from the type species, the genus is composed of four species from New Guinea (*minusculus* Carvalho; *neoguineanus* Carvalho; *nigrus* Carvalho, 1987; and *punctatus* Carvalho, 1987), one from Japan (*declivipennis* Miyamoto and Yasunaga, 1999), two from China (*chinensis* Liu and Zhao, 1999, and *testaceous* Liu and Zhao, 1999), two from Laos (*zetteli* Chérot et al., 2004, and *nakatanii* Chérot et al., 2004), and one from the Ivory Coast (*schuhi* Linnauvori 1994: all specimens of this species are presumed lost). An Australian species has also been introduced into Arizona, Hawaii, Texas, Florida, California, New Mexico, and Mexico.


Until recently this genus was considered to be relatively rare in collections (e.g., Schuh, 1974) and very few miridologists had seen specimens. Over the past 25 years an increasing number of specimens have been collected throughout the Old World tropics and subtropics. This has resulted in the description of several new species by various authors (Carvalho, 1987; Linnauvori, 1994; Liu and Zhao, 1999; Miyamoto and Yasunaga, 1999; Chérot et al., 2004), yet there has not been any real attempt to provide a synthesis of the genus.

This project was designed to address this taxonomic impediment and provide a monographic treatment of the genus. Phylogenetic and biogeographic analyses will be published separately. This work, the result of a larger effort aimed at documenting the mirid subfamilies Orthotylinae and Phylinae on a global basis, is part of the U.S. National Science Foundation–funded project, the Planetary Biodiversity Inventory (PBI).

## MATERIALS AND METHODS

### SPECIMENS

This study is based on specimens from existing museum collections as well as new material collected during the PBI fieldwork program. We have collected specimens of *variegatus* from New Caledonia; *chinensis*, *testaceous*, and *zetteli* from southern China and Laos; and *chenopoderis*, *monotocopsis*, and *pilbarensis* from across Australia, in part supported by our colleagues Randall T. Schuh from the American Museum of Natural History, Christiane Weirauch from



A–C Nikolet Tatarnic; D Chris Darling (Royal Ontario Museum); E–F Matthew Bulbert (Australian Museum)

Fig. 1. **A.** An unidentified species of *Phyllanthus* (Euphorbiaceae), a host plant of *C. variegatus* in New Caledonia. **B.** Adult *C. variegatus* on *Phyllanthus* bud. **C.** An unidentified species of *Macaranga* (Euphorbiaceae) in southern China, from which *C. chinensis*, *testaceous*, and *zetteli* were collected. **D.** Close-up of *C. chinensis* on *Macaranga* flowers. **E–F.** *C. monotocopsis* on *Monotoca elliptica* (Epacridaceae) in Sydney, Australia.



A-C Nikolai Tatarnic; D Tomohide Yasunaga

Fig. 2. **A.** *Rhagodia preissi* (Chenopodiaceae), one of many host plants of *C. chenopoderis*. **B.** Dark and light morphs of *C. chenopoderis* on *R. preissi*. **C.** *C. chenopoderis* on *R. preissi* flower. **D.** *C. testaceous*.

the University of California, Riverside, Celia Symonds from the Australian Museum, and Michael Wall from the San Diego Museum of Natural History. Michael Schwartz from Agriculture and Agri-food Canada, Thomas Henry from the National Museum of Natural

History, and Alfred Wheeler from Clemson University have also collected numerous specimens of *chenopoderis* from southern North America, while Dan Polhemus from the Hawaii Department of Land and Natural Resources has collected this same species in




Fig. 3. Illustration of female *C. pteraulos*, with insets indicating some alternate character states as found in other species: **A**. Metatibial spines may be either long and thick, as in *pteraulos*, or short and thin. **B**. The posterior margin of abdominal sternite II may be distinctly angular or relatively straight. **C**. The preepisternum may be either bilobed or unilobed. **D**. The modified lateral margin of the hemelytron, unique to *pteraulos* females.

Hawaii. Host plant records were obtained from many of these collecting events.

All specimens examined over the course of this project have been labeled with matrix code labels, giving each specimen a “unique specimen identifier” (USI). USI codes all begin with the PBI project code “AMNH\_PBI” followed by a unique eight-digit number (e.g., “AMNH\_PBI 00018973” is the USI for the holotype of *C. variegatus*). In a few cases, where species are represented by a large series of specimens, not all individuals were provided with USI codes; only specimens with USI codes are referred to in this work. USI codes are found in the lists of examined specimens, which also include all available collecting data as provided on labels, including host plant identification and georeference data whenever available. When coordinates were not originally provided, this information was obtained from gazetteers, atlases and other sources. These georeferencing data were used to generate all

the distribution maps included herein. Additional information on specimens and on the PBI project can be found on the PBI website (<http://research.amnh.org/pbi>).

Acronyms for the institutions from which we examined specimens are as follows:

AM	Australian Museum, Sydney
AMNH	American Museum of Natural History, New York
ANIC	Australian National Insect Collection, CSIRO, Canberra
BMNH	Natural History Museum, London
BPBM	Bishop Museum, Honolulu
CNC	Canadian National Collection of Insects, Agriculture and Agri-Food Canada, Ottawa
NHMW	Naturhistorisches Museum, Vienna
NKUM	Nankai University, Tianjin
QM	Queensland Museum, Brisbane
SAMA	South Australian Museum, Adelaide

SANC	Plant Protection Research Institute, Pretoria
TAMU	Texas A&M University, Department of Entomology, College Station
TYCN	Tomohide Yasunaga Collection, Nagasaki
ULB	Université Libre de Bruxelles, Brussels
USNM	United States National Museum of Natural History, Smithsonian Institution, Washington
ZMUC	Danish Natural History Museum, Copenhagen

#### HOMOLOGY AND TERMINOLOGY

Terminology for modifications of the male pygophore has been adopted from Miyamoto and Yasunaga (1999). Every attempt was made to dissect and illustrate each species using pen and ink drawings, photographs, and scanning electron micrographs (SEMs). For some species examinations were limited due to a paucity of specimens. Male genitalia were imaged *in situ* using SEM and were illustrated from dissections. In all species of *Coridromius* the aedeagus is reduced to a simple tube, which is routed through a gutter in the left paramere (see fig. 17A); thus, it is the shape of the paramere—not that of the aedeagus—that is of greater taxonomic importance. For this reason only the parameres are illustrated. Since *Coridromius* mate via traumatic insemination, the female genitalia are nonfunctional during mating and are therefore considerably reduced, with the posterior wall entirely membranous and the seminal depository reduced or absent (fig. 20A). These structures are not taxonomically informative and are not treated in this review. In contrast, the female external paragenitalia are variable between species and are described here in detail for the first time.

#### MEASUREMENTS

All measurements are maximum lengths in millimeters. Measurements include: clypeus to cuneus length, head length and width, pronotal length and width, scutellar length

and width, cuneus length, interocular distance, and the lengths of antennal segments I through IV. Because the hemelytra in *Coridromius* are strongly declivous caudal to the cuneal fracture, the total body length is inferred from measuring from the clypeus to the cuneal fracture, rather than to the end of the wing membrane. All measurements are shown in table 1.

#### SYSTEMATIC PLACEMENT

Owing to its unusual, compact body shape and unique genitalia, the tribal placement of *Coridromius* has proven problematic. Schuh (1974) initially placed *Coridromius* within the tribe Orthotylini, while acknowledging its similarity to the Hawaiian genus *Nesidiorchestes* Kirkaldy, the latter placed in the Halticini by Carvalho (1952). Schuh suggested that the genitalia of these two genera needed comparison to determine whether they are in fact closely related and to determine their tribal placement. He noted that although the Halticini are sometimes difficult to distinguish from the Orthotylini on external characters alone, the lack of interramal lobes on the posterior wall of the female genitalia always distinguishes the Halticini (Schuh, 1974). The placement of *Coridromius* in the Orthotylini persisted, with Linnauori (1994) noting that it differed significantly from other Orthotylini by its small, compact body, *Halticus*-like legs and unique male genitalia, features that suggest the genus may require its own tribe. He thought *Coridromius* to be related to *Coridromoides* and perhaps also *Nesidiorchestes*, both of which belonged to the Halticini.

After examining the female genitalia, Cassis and Gross (1995) and Schuh (1995) placed *Coridromius* within the Halticini. While they did not explicitly justify their taxonomic placement, there are nonetheless several characters to support this decision: the tarsal claws lack pulvilli, the aedeagus is small with the vesica membranous and lacking spiculi, and most telling of all, the posterior wall lacks interramal lobes.

In light of what we now know of the mating biology of *Coridromius* (discussed below), any inferences of the homology of genitalic structures should be made with

TABLE 1  
Measurements of *Coridromius* species

TABLE 1  
*(Continued)*



TABLE 1  
*(Continued)*

TABLE 1  
(Continued)



TABLE 1  
(Continued)



TABLE 1  
(Continued)

TABLE 1  
(Continued)

Species	Length				Width				AI	AIIV	AIV
	Clyp-Cun	Head	Pronotum	Scutellum	Cuneus	Head	Pronotum	Scutellum			
<i>zetteli</i>	<b>Mean</b>	<b>1.80</b>	<b>0.14</b>	<b>0.61</b>	<b>0.39</b>	<b>0.45</b>	<b>0.80</b>	<b>1.24</b>	<b>0.59</b>	<b>0.49</b>	<b>0.15</b>
	SD	0.07	0.03	0.04	0.04	0.03	0.01	0.02	0.01	0.02	0.02
	Range	0.15	0.09	0.11	0.09	0.06	0.03	0.06	0.02	0.03	0.03
	Min	1.72	0.11	0.57	0.33	0.42	0.79	1.22	0.58	0.48	0.13
	Max	1.87	0.19	0.68	0.42	0.48	0.82	1.27	0.60	0.51	0.17
	Count	5	5	5	5	5	5	5	5	4	3

caution. For example, the aedeagus in *Coridromius*, while membranous as in other halticinae, is coupled with the left paramere to form a highly specialized intromittent organ unlike that of any other Halticinae (or indeed any other Miridae). Furthermore, while the posterior wall of the female genitalia lacks interramal lobes, it also lacks any sclerotization whatsoever, a point of distinction.

## BIOLOGY

### HABITATS AND MICROHABITATS

*Coridromius* has a largely paleotropical distribution and is found in a wide range of habitats, from arid landscapes to wet tropical forests. Some of the species have widespread distributions and range across a number of habitat types; for example, *chenopoderis* is known from arid and semiarid shrubland and open woodland across temperate Australia. Many other species are known only from the type locality, such as the rainforest species *nigrus*, *prolixipenis*, *drepanopenis*, *epithema*, *ephippius*, and *sommelieri*, but generalizations about restricted habitat range cannot be made without more extensive collecting. While many species are known from either single or very few specimens, others have been collected in very large numbers (e.g., the Australian species *chenopoderis*, *monotocopsis*, and *pilbarensis*, the northern Sulawesi species *confusus*, and the type species *variegatus*).

*Coridromius* is thought to be phytophagous, with a number of species found most commonly on the reproductive components of their host plants, both on buds and mature flowers, but also on leaves and stems (e.g., *chenopoderis*, *chinensis*, *monotocopsis*, *variegatus*; figs. 1 and 2). There are occurrences of multiple species of *Coridromius* co-occurring on the same host plants without evidence of resource partitioning. For example, in southern China one of us (NT) collected *chinensis*, *zetteli*, and *testaceus* all on the same *Macaranga* tree (Euphorbiaceae) at the same time. Similarly, there are several species that are known from the same collecting locality, sometimes even the same collecting event. For example, *drepanopenis*, *falsicoleus*, *pteraulos*, and *sommelieri* were all collected by T.

TABLE 2  
Host plants of various *Coridromius* species

Species	Host plant genus and species	Plant family	Plant Order	Number of specimens collected
bicolor	<i>Macaranga</i> sp.	Euphorbiaceae	Malpighiales	5
bicolor	<i>Mallotus</i> sp.	Euphorbiaceae	Malpighiales	7
bicolor	<i>Mangifera</i> sp.	Anacardiaceae	Sapindales	2
boianotum	<i>Acalypha</i> sp.	Euphorbiaceae	Malpighiales	1
boianotum	unknown sp.	Urticaceae	Rosales	1
boianotum	<i>Pipturus</i> sp.	Urticaceae	Rosales	10
bulbopella	<i>Berkheya bipinnatifida</i>	Asteraceae	Asterales	6
chenopoderis	<i>Olearia axillaris</i>	Asteraceae	Asterales	3
chenopoderis	<i>Olearia muelleri</i>	Asteraceae	Asterales	22
chenopoderis	<i>Lepidium oleraceum</i>	Brassicaceae	Brassicales	7
chenopoderis	unknown sp.	Chenopodiaceae	Caryophyllales	16
chenopoderis	<i>Atriplex amnicola</i>	Chenopodiaceae	Caryophyllales	47
chenopoderis	<i>Atriplex isatidea</i>	Chenopodiaceae	Caryophyllales	60
chenopoderis	<i>Atriplex nummularia omissa</i>	Chenopodiaceae	Caryophyllales	3
chenopoderis	<i>Chenopodium auricomum</i>	Chenopodiaceae	Caryophyllales	79
chenopoderis	<i>Chenopodium murale</i>	Chenopodiaceae	Caryophyllales	140
chenopoderis	<i>Einadia nutans</i> subsp. <i>nutans</i>	Chenopodiaceae	Caryophyllales	15
chenopoderis	<i>Enchytraea tomentosa</i> var. <i>tomentosa</i>	Chenopodiaceae	Caryophyllales	3
chenopoderis	<i>Halosarcia indica</i> <i>bidens</i>	Chenopodiaceae	Caryophyllales	20
chenopoderis	<i>Maireana oppositifolia</i>	Chenopodiaceae	Caryophyllales	25
chenopoderis	<i>Rhagodia</i> sp.	Chenopodiaceae	Caryophyllales	7
chenopoderis	<i>Rhagodia baccata</i>	Chenopodiaceae	Caryophyllales	88
chenopoderis	<i>Rhagodia baccata</i> <i>baccata</i>	Chenopodiaceae	Caryophyllales	13
chenopoderis	<i>Rhagodia latifolia</i> subsp. <i>latifolia</i>	Chenopodiaceae	Caryophyllales	11
chenopoderis	<i>Rhagodia parabolica</i>	Chenopodiaceae	Caryophyllales	34
chenopoderis	<i>Rhagodia preissi</i>	Chenopodiaceae	Caryophyllales	60
chenopoderis	<i>Rhagodia preissii</i> <i>preissii</i>	Chenopodiaceae	Caryophyllales	53
chenopoderis	<i>Rhagodia preissii</i> subsp. <i>obovata</i>	Chenopodiaceae	Caryophyllales	34
chenopoderis	<i>Rhagodia spinescens</i>	Chenopodiaceae	Caryophyllales	9
chenopoderis	<i>Salsola tragus</i> subsp. <i>tragus</i>	Chenopodiaceae	Caryophyllales	128
chenopoderis	<i>Suaeda australis</i>	Chenopodiaceae	Caryophyllales	18
chenopoderis	<i>Frankenia</i> sp.	Frankeniaceae	Caryophyllales	1
chenopoderis	<i>Muehlenbeckia florulenta</i>	Polygonaceae	Caryophyllales	14
chenopoderis	<i>Acacia</i> sp.	Fabaceae	Fabales	20
chenopoderis	<i>Acacia acinacea</i>	Fabaceae	Fabales	1
chenopoderis	<i>Acacia oswaldii</i>	Fabaceae	Fabales	6
chenopoderis	<i>Acacia longifolia</i>	Fabaceae	Fabales	2
chenopoderis	<i>Allocasuarina humilis</i>	Casuarinaceae	Fagales	1
chenopoderis	<i>Dicrastylis fulva</i> forma <i>fulva</i>	Lamiaceae	Lamiales	1
chenopoderis	<i>Teucrium racemosum</i>	Lamiaceae	Lamiales	1
chenopoderis	<i>Eremophila forrestii</i>	Scrophulariaceae	Lamiales	1
chenopoderis	<i>Eremophila ionantha</i>	Scrophulariaceae	Lamiales	1
chenopoderis	<i>Eremophila mitchellii</i>	Scrophulariaceae	Lamiales	4
chenopoderis	<i>Myoporum insulare</i> ?	Scrophulariaceae	Lamiales	6
chenopoderis	<i>Alyogyne pinoniana</i>	Malvaceae	Malvales	2
chenopoderis	<i>Pimelea ferruginea</i>	Thymelaeaceae	Malvales	2
chenopoderis	<i>Pimelea rosea</i>	Thymelaeaceae	Malvales	1
chenopoderis	<i>Agonis parviceps</i>	Myrtaceae	Myrales	1
chenopoderis	<i>Melaleuca brevifolia</i>	Myrtaceae	Myrales	1
chenopoderis	<i>Melaleuca uncinata</i>	Myrtaceae	Myrales	3
chenopoderis	<i>Scholtzia capitata</i>	Myrtaceae	Myrales	19
chenopoderis	<i>Scholtzia drummondii</i>	Myrtaceae	Myrales	2

TABLE 2  
(Continued)

Species	Host plant genus and species	Plant family	Plant Order	Number of specimens collected
<i>chenopoderis</i>	<i>Hakea leucoptera</i>	Proteaceae	Proteales	110
<i>chenopoderis</i>	<i>Hakea mitchellii</i>	Proteaceae	Proteales	1
<i>chenopoderis</i>	<i>Malus</i> sp.	Rosaceae	Rosales	16
<i>chenopoderis</i>	<i>Prunus</i> sp.	Rosaceae	Rosales	2
<i>chenopoderis</i>	<i>Prunus</i> sp.	Rosaceae	Rosales	9
<i>chenopoderis</i>	<i>Amyema cambagei</i>	Loranthaceae	Santalales	1
<i>chenopoderis</i>	<i>Amyema quandang</i> var. <i>quandang</i>	Loranthaceae	Santalales	1
<i>chenopoderis</i>	<i>Exocarpos aphyllus</i>	Santalaceae	Santalales	19
<i>chenopoderis</i>	<i>Nitraria billardierei</i>	Nitrariaceae	Sapindales	19
<i>chenopoderis</i>	<i>Flindersia maculosa</i>	Rutaceae	Sapindales	6
<i>chenopoderis</i>	<i>Nitraria billardierei</i>	Nitrariaceae	Sapindales	30
<i>chinensis</i>	<i>Macaranga</i> sp.	Euphorbiaceae	Malpighiales	57
<i>chinensis</i>	<i>Mallotus japonicus</i>	Euphorbiaceae	Malpighiales	4
<i>chinensis</i>	<i>Pentaspodon motleyi</i>	Anacardiaceae	Sapindales	1
<i>minusculus</i>	unknown sp.	Cyperaceae	Poales	1
<i>monotocopsis</i>	<i>Chrysanthemoïdes monolifera</i>	Asteraceae	Asterales	1
<i>monotocopsis</i>	<i>Brachyloma daphnoides</i>	Ericaceae	Ericales	8
<i>monotocopsis</i>	<i>Monotoca</i> sp.	Ericaceae	Ericales	16
<i>monotocopsis</i>	<i>Monotoca elliptica</i>	Ericaceae	Ericales	364
<i>monotocopsis</i>	<i>Acacia longifolia</i>	Fabaceae	Fabales	1
<i>pilbarensis</i>	<i>Scaevola crassifolia</i>	Goodeniaceae	Asterales	1
<i>pilbarensis</i>	<i>Atriplex bunburyana</i>	Amaranthaceae	Caryophyllales	16
<i>pilbarensis</i>	<i>Gastrolobium spinosum</i>	Fabaceae	Fabales	1
<i>pilbarensis</i>	<i>Agonis linearifolia</i> ?	Myrtaceae	Mytales	1
<i>pilbarensis</i>	<i>Melameuca hamulosa</i>	Myrtaceae	Mytales	6
<i>pilbarensis</i>	<i>Melaleuca rhamphophylla</i>	Myrtaceae	Mytales	1
<i>pilbarensis</i>	<i>Melaleuca sheathiana</i>	Myrtaceae	Mytales	3
<i>pilbarensis</i>	<i>Melaleuca uncinata</i>	Myrtaceae	Mytales	21
<i>pilbarensis</i>	<i>Santalum lanceolatum</i>	Santalaceae	Santalales	4
<i>pilbarensis</i>	<i>Euroschinus falcata</i>	Anacardiaceae	Sapindales	5
<i>pilbarensis</i>	<i>Schinus areira</i>	Anacardiaceae	Sapindales	2
<i>pilbarensis</i>	<i>Dodonaea viscosa angustissima</i>	Sapindaceae	Sapindales	28
<i>prolixipenis</i>	<i>Homalanthus</i> sp.	Euphorbiaceae	Malpighiales	1
<i>punctatus</i>	<i>Rubus</i> sp.	Rosaceae	Rosales	1
<i>testaceous</i>	<i>Macaranga</i> sp.	Euphorbiaceae	Malpighiales	7
<i>testaceous</i>	unknown sp.	Cyperaceae	Poales	2
<i>variegatus</i>	<i>Phyllanthus</i> sp.	Euphorbiaceae	Malpighiales	25
<i>variegatus</i>	unknown sp.	Sapindaceae	Sapindales	24
<i>zetteli</i>	<i>Macaranga</i> sp.	Euphorbiaceae	Malpighiales	2

C. Maa in Tenompok, Sabah, between the 10th and 19th of September 1959. Unfortunately no host data were recorded for these collecting events.

#### HOST PLANT ASSOCIATIONS

Host plant associations are known for 12 of the 32 species documented in this work (see table 2), with the vast majority of host records coming from the Australian species

*chenopoderis*, which is undoubtedly polyphagous.

It is well-known that many Heteroptera are associated with higher angiosperm clades such as the rosids and asterids (Cassis and Gross, 2002; Cassis and Vanags, 2006; Cassis et al., 2007); this also appears to be the case for *Coridromius*. There are two records of *Coridromius* species on monocots, with *minusculus* (in Papua New Guinea) and *testaceous* (in Laos) collected from undetermined

Cyperaceae species, but in both cases the records refer to only one or two specimens and may only be sitting records.

Excluding the extensive list of host plant associations known for *chenopoderis* and the single host record of *minusculus*, 11 of the species of *Coridromius* with host records are known from rosid angiosperms. The majority of these have been collected from the family Euphorbiaceae (*bicolor*, *boianotum*, *prolixipennis*, *testaceus*, *variegatus*, and *zetteli*). Only two species are known from asterids: *bulbopella* from South Africa and *monotocopsis* from eastern Australia.

The ubiquitous Australian species *chenopoderis*, which has also been discovered extrazonally in Hawaii and parts of southern North America, is most commonly associated with chenopods (Chenopodiaceae; 17 species, see table 2). However, on two separate occasions it has been taken on the proteaceous genus *Hakea*, with over 100 specimens at one location. Another Australian species, *pilbarensis*, has also been collected from a chenopod at a single location. Although chenopods are common hosts for *Coridromius* within Australia, no other species of *Coridromius* are known from chenopods elsewhere.

#### SEXUAL BEHAVIOR

The genitalia of *Coridromius* males are unique among the Miridae, with the left paramere and aedeagus coupled to form a scyphelike copulatory organ, analogous to that found in the Cimicidae, Polycenidae, and some Anthocoridae. Members of these three families mate by a process termed “traumatic insemination”, whereby males stab females in the abdomen with their needlelike genitalia and ejaculate into their body cavity (Carayon, 1966, 1977). It is thought that this form of mating has evolved as a means for males to circumvent female pre- or postmating resistance (Stutt and Siva-Jothy, 2001; Arnqvist and Rowe, 2005). Not surprisingly, this form of mating is costly to females in terms of physical damage (Morrow and Arnqvist, 2003) and increased risk of infection (Reinhardt et al., 2003). In response, females of many of these species have evolved internal and external paragen-

tal structures, collectively known as the “spermalege” (Carayon, 1959, 1966, 1977) to receive the intromittent organ and sperm, and to reduce the costs of mating (Stutt and Siva-Jothy, 2001; Morrow and Arnqvist, 2003; Reinhardt et al., 2003).

As in the above taxa, *Coridromius* is also now known to practice traumatic insemination, with the females of many species exhibiting elaborate paragenitalia (Tatarnic et al., 2006). We have observed the mating behavior of four species in captivity (*chenopoderis*, *chinensis*, *monotocopsis*, and *variegatus*). When several specimens of both sexes were kept in small test tubes with host plant clippings, males were observed pouncing on the dorsum of nearby individuals, generally females but occasionally other males and even nymphs. In all cases mounted individuals would shake vigorously, jump about, and kick in an apparent attempt to dislodge their would-be suitors. During these struggles mounting males curled their abdomens over to the right side of their mate’s body, extended their left paramere and attempted to stab their partner anteriorly along the right side of the abdomen. Such mating attempts were very brief, never lasting more than about 30 seconds.

#### TAXONOMY

##### *Coridromius* Signoret

*Ocypus* Montrouzier, 1861: 67, junior homonym of *Ocypus* Leach, in Samouelle, 1819: 172 (Coleoptera). Type species: *Ocypus variegatus* Montrouzier, 1861, by monotypy.

*Coridromius* Signoret, 1862: 5 (nom. nov. for *Ocypus* Montrouzier 1861).

*Neocyprus* Distant, 1914: 378 (unnecessary nom. nov. for *Ocypus* Montrouzier, 1861). Carvalho, 1987: 61; Linnauvori, 1994: 15; Schuh, 1995: 46; Cassis and Gross, 1995: 185; Liu and Zhao, 1999: 55; Miyamoto and Yasunaga, 1999: 33; Chérot et al., 2004: 57.


*Coridromoides* Carvalho, 1956: 54 (new genus). Type species: *Coridromoides carinatus* Carvalho, 1956, by original designation.

**DIAGNOSIS:** *Coridromius* is recognized by the following character states: body compact and stout (figs. 1–5); small, with most species between 2–3 mm in length (table 1); short and broad head with substylate eyes (figs. 4–

6); hemelytra sharply deflected at cuneal fracture; greatly enlarged metafemora, frequently marked with dark brown diagonal banding (figs. 1–3); femora with recessed bothria; aedeagus simple, membranous, without sclerotizations (figs. 12A, 17A); left paramere larger than right and generally scythe-shaped, with gutter running from base to apex, coupled with aedeagus to form piercing intromittent organ (figs. 12A, 17A); right paramere smaller than left, triangulate to club-shaped, sometimes with short and rounded (e.g., figs. 10B–C, 10G–H) or long and sharp (e.g., figs. 7O–P) apical process; female genitalia reduced, with the posterior wall entirely membranous (fig. 20A); female paragenitalia sometimes present (figs., 8C–D, 9C–D, 11C–D, 11G–H, 13A–B, 13D–F, 16C–D, 18F–G, 19B–C, 19G, 20B–C, 21A). *Coridromius* is readily distinguished from all other Halticini (and all other Miridae) by its unique male genitalia.

**REDESCRIPTION:** COLORATION: Variable, but typically brown with various brown, yellow, and cream-colored markings; frequently with dark brown vittae on frons and transverse stripes on outer surface of metafemur (e.g., figs. 1–3). SURFACE AND VESTITURE: Head, pronotum, scutellum, propleuron, metepimeron, and hemelytra impunctate to punctate; broadly covered with short or long simple, white, decumbent setae. STRUCTURE: Head (fig. 6): Transverse, short; eyes substylate; frons tumescent medially, shallowly depressed adjacent to eyes; vertex usually with a pair of slightly raised tubercles adjacent to eyes, bordered posteriorly by shallow foveae; posterior margin of head weakly to strongly medially rounded; mandibular and maxillary plates well defined; gena sometimes swollen; clypeus prominent; buccula arcuate and flared; labium reaching slightly beyond hind coxae. Antenna: AI very short and cylindrical (approximately five times shorter than width of vertex); AII about one-fifth longer than AI and apically clavate; AIII short, about twice as long as AI; AIV about as long as AI (table 1). Thorax: Pronotum broad and convex, anterolateral margins usually somewhat upturned, submarginal region of humeral angles shallowly excavate, collar present and rounded, visible from above, collar

region either undifferentiated or weakly defined, lateral margins sometimes carinate; mesoscutum steeply declivent, sometimes visible, scutellum flat or swollen and rounded, sometimes with rounded lobe at apex (e.g., fig. 1D); proepisternum lobate, with either one (figs. 3C, 13C) or two (figs. 3C, 18A) lobes projecting laterally; metathoracic spiracle prominent; metepimeron platelike, with posterior margin either truncate (e.g., figs. 8E, 11C) or extending partly over abdomen (e.g., figs. 9E–F, 11G–H, 13E–F); metanotum sometimes raised into a vertical transverse plate separating thorax from abdomen, with lateral margins prominent and flared (figs. 9C–D, 16C–D). Metathoracic gland: External efferent system reduced, peritreme elongate, fingerlike, along posterior margin of metepisternum, bounded anteriorly by a thin strip of evaporative bodies. Hemelytra: Macropterous; steeply declivent at cuneal fracture; embolium thin and platelike, sometimes flared; cuneus small and triangular. Legs: Metafemur swollen, laterally compressed near apex; caudal margin of hind tibia with two rows of minute spines interspersed with longer spines, sometimes noticeably thickened (fig. 3A). MALE GENITALIA: Pygophore trapeziform, posterior margin typically biconcave, sometimes with fold (figs. 8A, 8H, 14D, 16A, 21C–D), groove (figs. 11E–F, 14G–H, 19H) or mesal suture on left side (figs. 9A, 9H, 11A, 13H, 14F, 16E, 19D, 21A), sometimes with ventral apical process projecting from right side (figs. 9A, 9G–H, 11A, 13H, 14E–F, 16A–B, 16E, 18D, 19D); parameres heavily sclerotized; right paramere always smaller than left, usually either triangular and somewhat elongate (figs. 7B–C, 7E–G, 7L–M, 10J, 12B–C, 12E–F, 12K–L, 12O–P, 15B–C, 17B–D, 17M–O, 17Q–R), boot-shaped (figs. 7I–J, 15H–I), or club-shaped with a long, spinelike (figs. 7O–P, 7R–S, 12N–O, 17F–G) or short and more rounded (figs. 10B–D, 10F–H, 15E–F, 17B–D, 17P–Q) apical process; left paramere generally curved and scyphelike, sometimes with one to several tight coils (e.g., figs. 7H, 8F, 17H, 17S, 19E–F, 21D), apically acuminate, with cleft gutter running entire length, sometimes completely enclosed until apex (figs. 7Q, 9G–H, 15D, 14E–F); aedeagus simple, flagellate, with



Nikolai Tatarnic

Fig. 4. Habitus views of *Coridromius* spp.: *bicolor*–*lestoni*.




Fig. 5. Habitus views of *Coridromius* spp.: *marmoreus*–*zetteli*.



Nikolai Tatarnic

Fig. 6. Anterior view of head of *Coridromius* spp.

membranous to weakly sclerotized phalotheca and phallobase, permanently routed through channel of left paramere (e.g., fig. 17A). FEMALE GENITALIA: Simple and reduced, with posterior wall membranous, interramal lobes absent (e.g., fig. 20A); abdomen sometimes with visible external paragenital modifications.

**REMARKS:** The genus *Coridromius* is easily recognized by the unique body shape, extremely swollen hind femora and highly specialized male genitalia. While several species are easily identifiable by coloration and body shape, others are sometimes more difficult to place. For most species the male genitalia can typically be examined *in situ*, but often one must remove female specimens from points or remove the hemelytra in order to expose possible paragenital modifications, which have always been found on the right side of the abdomen. The following key is intended to be used with both sexes, except in those species that are known by one sex.

#### KEY TO SPECIES OF *CORIDROMIUS*

1. Proepisternum bilobed (figs. 3C, 18A); metanotum prominent and flared (fig. 16C–D) 2
- Proepisternum unilobed (figs. 3C, 13C); metanotum not prominent and flared (fig. 13D) . . . . . 17
2. Metepimeron punctate (figs. 13A, 16G, 19A–B); right paramere club-shaped with short, rounded process, appearing U-shaped from above (figs. 10B–D, 10F–H, 15E–F, 17B–D, 17I–K, 17T–U). . . . . 3
- Metepimeron impunctate (figs. 8E, 9E–F, 11G, 13E, 14B, 18C, H, 19G); right paramere not club-shaped, not appearing U-shaped when viewed from above . . . . . 6
3. Costal margin of hemelytron straight and weakly carinate (fig. 5: *sommelieri*, *testaceus*, *zetteli*) . . . . . 4
- Costal margin of hemelytron weakly to strongly flared posteriorly (fig. 4: *crassus*, *declivipennis*, *falsicoleus*; fig. 5: *nakatanii*, *pteraulos*) . . . . . 11
4. Male left paramere tightly corkscrewed (figs. 17H, 17S, 19E–F, 21D); female possibly with corkscrewed paragenital opening on right side of abdomen (fig. 19B–C) . . . . . 5
- Known only from females, female without any visible paragenitalia; male suspected to not have corkscrewed left paramere (South China, Laos, Nepal, Sumatra, Vietnam) . . . *testaceus*
5. Apex of male left paramere perpendicular to its axis, pygophore with triangular ventral apical process (figs. 17H, 19D–F); female with corkscrewed paragenital opening on right side of body (figs. 19B–C, 20B) (Sabah) . . . . . *sommelieri*
- Apex of male left paramere almost parallel to its axis (figs. 17S, 21D); pygophore without apical process (fig. 21D); female unknown (South China, Laos, Vietnam) . . . . . *zetteli*
6. Dorsum with mottled cream and brown coloration (figs. 1D, 4: *chinensis*, *confusus*; fig. 5: *marmoreus*, *minusculus*, *punctatus*) . . . . . 7
- Not with mottled cream and brown coloration . . . . . 15
7. Both lobes of proepisternum of equal length (e.g., fig. 18A); lateral margins of pronotum and hemelytra without brown spots; right paramere rounded with long, sharp, triangular apical process (figs. 7O–P, 7R–S, 9A, 12N–O, 18D–E) . . . . . 8
- Anterior lobe of proepisternum shorter than posterior lobe (fig. 14A); lateral margins of pronotum and hemelytra marked with small brown spots (fig. 5: *marmoreus*); right paramere without sharp, triangular apical process (fig. 12K–L) (Philippines) . . . . . *marmoreus*
8. Apex of scutellum greatly swollen and rounded (figs. 1D, 4: *chinensis* 1, *chinensis* 2); AII without dark brown postmedial band (figs. 1D, 4: *chinensis*); gutter of left paramere open over entire length (figs. 7N, 9A) (India, Nepal, South China, Laos, Brunei, Japan, Taiwan, Vietnam) . . . . . *chinensis*
- Apex of scutellum only slightly swollen (fig. 4: *confusus*; fig. 5: *minusculus*, *punctatus*); AII with dark brown postmedial band; left paramere either entirely open or closed until apex . . . . . 9
9. AI dark brown with pale base and apex; AIII and AIV entirely dark brown; left paramere open over entire length (fig. 17E) (Papua New Guinea) . . . . . *punctatus*
- AI pale with broad, dark brown annulation; AIII and AIV dark brown, yellowish white basally; left paramere open or closed. . . . . 10
10. Left paramere a thin, closed tube, open at apex (figs. 9G–H, 7Q) (Sulawesi, Solomon Islands) . . . . . *confusus*
- Left paramere open over entire length (fig. 12M) (Irian Jaya and Papua New Guinea) . . . . . *minusculus*
11. Costal margin of hemelytron weakly flared posteriorly (fig. 4: *declivipennis*, *falsicoleus*, fig. 5: *pteraulos*) . . . . . 12
- Costal margin of hemelytron strongly flared posteriorly (fig. 4: *crassus*; fig. 5: *nakatanii*). . . . . 14

12. Body dark brown with yellow-orange markings (fig. 4: *falsicoleus*), female with prominent paragenital opening on right side of abdominal sternite II (figs. 20A–B); male unknown (India, Sabah) . . . . . *falsicoleus*
- Body color variable, not as above; female without prominent paragenital opening on right side of body . . . . . 13
13. Posterolateral flaring of hemelytron rounded (fig. 5: *pteraulos*); anterolateral margin of hemelytron on female strongly recurved, forming a distinct gutter (figs. 3D, 16H); left posterior ventral margins of male abdominal segments VII and VIII with patch of strong, caudally directed bristlelike setae (fig. 16E–F) (Papua New Guinea, Sabah) . . . . . *pteraulos*
- Posterolateral flaring of hemelytra angular (fig. 4: *declivipennis*); anterolateral margin of hemelytra on female not forming a curved tube; patch of setae on male abdominal segments VII and VIII much less distinct than in *pteraulos* (Japan) . . . . . *declivipennis*
14. Dark brown apical annulation on AII separated from postmedial dark brown band by thin white annulation (fig. 5: *nakatanii*); left paramere long, thin, and enclosed until apex (figs. 14E–F, 15D), female with prominent paragenital opening on right side within abdominal sternite II (Laos, Vietnam). . . . . *nakatanii*
- AII with dark brown apical annulation and subapical whiteannulation, but no dark post-medial band (fig. 4: *crassus*); left paramere thick and relatively long, not enclosed until apex (fig. 10A) (North Sulawesi). . . . . *crassus*
15. Body small, brown, and punctate; posterior margin of embolium strongly flared outward, cuneal fracture deep (fig. 4: *lestoni*); dorso-lateral margins of abdomen grooved (Ghana). . . . . *lestoni*
- Uniform brownish yellow with two large dark brown spots on scutellum (fig. 4: *bulbopella*; fig. 5: *ruwenzorii*); posterior margin of vertex broadly rounded (fig. 6). . . . . 16
16. Head smooth and impunctate; female and male with “cup and bulb” structure on right abdominal tergites II and III, hidden by hemelytra (fig. 8C–D); male left paramere long and thin with single twist at base (fig. 8F) (South Africa, Kenya, Tanzania). . . . . *bulbopella*
- Head smooth and weakly punctate; female with small lobed projection on right abdominal tergites II and III, hidden by hemelytra (fig. 18F–G); male unknown (Uganda). . . . . *ruwenzorii*
17. Anterolateral margins of pronotum projecting as rounded lobes, uniform brownish yellow with two dark brown spots on scutellum (fig. 4: *boianotum*); posterior margin of pygophore weakly bisinuate, without deep groove, fold, or suture on left side (fig. 8B) (Solomon Islands, Vanuatu) . . . . . *boianotum*
- Anterolateral margins of pronotum not projecting as rounded lobes; coloration variable; posterior margin of pygophore with or without groove, fold or mesal suture on left side . . . . . 18
18. Pronotum campanulate, much wider posteriorly; deeply punctate; anterior of humeral angles projecting forward; brownish yellow (fig. 4: *carinatus*) (Palau) . . . . . *carinatus*
- Pronotum not campanulate; anterior of humeral angles not projecting forward; coloration and texture variable . . . . . 19
19. Light tan in coloration, with paired dark brown spots on pronotum behind eyes, on pronotal disc, and on scutellum, hemelytra with brown transverse chevron pattern (fig. 5: *neoguineanus*); head with irregular dark brown band on clypeus, sometimes extending upward onto frons, middle of vertex with a single dark brown spot, frons without vittae (fig. 6: *neoguineanus*); right paramere boot-shaped (fig. 15H–I); posterior margin of pygophore with deep U-shaped groove right of centre (fig. 14G–H) (Papua New Guinea, Irian Jaya) . . . . . *neoguineanus*
- Coloration not as above; right paramere triangulate; groove or fold on left posterior margin of pygophore . . . . . 20
20. Hemelytra always with dark brown “Y” along inner margin of clavus (fig. 5: *tahitiensis* ♂, *tahitiensis* ♀); left margin of pygophore with a deep U-shaped groove (fig. 19H); female with prominent intersegmental paragenital opening on right side of abdomen between segments II and III (figs. 26G, 27C) (Tahiti). . . . . *tahitiensis*
- Hemelytra without dark brown “Y” pattern along inner margin of clavus; folded groove on posterior margin of pygophore deep or shallow; female without paragenitalia as in *tahitiensis* . . . . . 21
21. Metepimeron extended posteriorly as a long winglike lobe (figs. 11G, 13E); costal margin of hemelytron flared anteriorly, abruptly angled inward at midpoint (figs. 4: *epithema*, *hermosus*, 5: *nigrus*, *prolixipenis*) . . . . . 22
- Metepimeron truncate or only slightly lobed; lateral margin of embolium not shaped as above . . . . . 25
22. Head with contrasting dark brown or black coloration above and yellow below, beginning at base of clypeus; yellow piping along

- posterior margin of vertex and adjacent to eyes (fig. 6: *hermosus*, *prolixipenis*) . . . . . 23
- Head not colored dark above and yellow below; with or without yellow piping . . . . . 24
23. Body orange-brown with two conspicuous black markings on anterior portion of pronotum, scutellum often bright orange (fig. 4: *hermosus*); head black above, yellow below (fig. 6: *hermosus*); metepimeron extended as a long, triangular lobe (fig. 13E); male left paramere thick and relatively long (figs. 12D, 13G); posterior margin of pygophore with broad ventral apical process (fig. 13H); right dorsal laterotergites II and III of female swollen and desclerotized (fig. 13D–F) (Papua New Guinea) . . . . . *hermosus*
- Body dark reddish brown above (fig. 5: *prolixipenis*); head dark reddish brown above and yellow below, with yellow piping (fig. 6: *prolixipenis*); fore- and middle legs yellow basally, hindlegs uniform dark reddish-brown; left paramere extremely long and thin (fig. 15M–N); posterior margin of pygophore without ventral apical process (fig. 15N); female unknown (Papua New Guinea) . . . . . *prolixipenis*
24. Body uniformly dark reddish brown to black with faint orange piping along margins of pronotum (fig. 5: *nigrus*); posterior margin of head, outer ocular margins and mandibular plates yellow-orange (fig. 6: *nigrus*); metafemur uniformly dark; metepimeron extended as a long, triangular lobe (as in fig. 13E); right abdominal laterotergites II and III of female swollen and desclerotized (as in fig. 13D–F); male unknown (Papua New Guinea) . . . . . *nigrus*
- Body uniformly orange-brown (fig. 4: *epithema*); head and pronotum with faint orange piping (fig. 6: *epithema*); metafemur uniformly dark, metepimeral lobe extremely long and apically rounded (fig. 11G); right abdominal laterotergites II and III of female swollen, membranous and saddle-shaped (fig. 11G–H); male unknown (Papua New Guinea) . . . . . *epithema*
25. Body uniformly dark brown (fig. 5: *thalame*) with orange-yellow legs and head (fig. 6: *thalame*); right abdominal segment II of female irregularly swollen, conspicuous paragenital opening between segments II and III (fig. 21A); male unknown (Sabah) . . . . . *thalame*
- Body not uniformly dark brown; female without paragenitalia as in *thalame* . . . . . 26
26. Body nearly uniform brownish orange (fig. 4: *drepanopenis*); left paramere extremely long and thin (figs. 10I, 10K); posterior margin of pygophore with folded groove (fig. 10K); female unknown (Sabah) . . . . . *drepanopenis*
- Body not colored as above; left paramere not extremely long and thin; groove on pygophore variable . . . . . 27
27. Dorsolateral margin of abdominal segment II of females extremely swollen, with laterally projecting tumescence at posterior margin of segment (fig. 11B–D); male with relatively long and thick left paramere (figs. 11E–F, 12A); left posterior margin of pygophore with shallow depression (Philippines) . . . . . *ephippius*
- Females without visible external paragenitalia; right posterior margin of pygophore with distinct folded groove . . . . . 28
28. Posterior margin of metepimeron with small but distinct rounded lobe projecting from upper corner (as in fig. 14B); folded groove on posterior margin of pygophore shallow (fig. 21B–C) (New Caledonia, Fiji) *variegatus*
- Posterior margin of metepimeron truncate; folded groove in pygophore shallow (fig. 21D) or deep (figs. 8H, 16A) . . . . . 29
29. Body broad, generally light brown in color with darker brown markings (fig. 5: *monotocopsis*); pygophore with shallow folded groove (fig. 14D); left paramere marginally longer than in other Australian species (fig. 15A); found primarily on *Monotoca elliptica* (eastern Australia) . . . . . *monotocopsis*
- Body broad or narrow, either large and orange-brown with brown and yellow markings (fig. 5: *pilbarensis*) or small, ranging from almost black to olive in coloration (figs. 2B–C, 4: *chenopoderis* 1, *chenopoderis* 2); folded groove on pygophore deep (figs. 8H, 16A) . . . . . 30
30. Body large and broad, orange-brown with brown and yellow markings (fig. 5: *pilbarensis*); pygophore with small tubercle to right of deep, folded groove (fig. 16A–B) (Australia) . . . . . *pilbarensis*
- Body small and narrow, coloration varying from almost black to olive or light brown (figs. 2B–C, 4: *chenopoderis* 1, *chenopoderis* 2); folded groove on pygophore deep, without small tubercle on right side (fig. 8H); found primarily on Chenopodiaceae (Australia, New Zealand, Norfolk Island, Mexico, Arizona California, Florida, Hawaii, New Mexico, Texas) . . . . . *chenopoderis*

***Coridromius bicolor*, new species**  
Figures 4, 6, 7A–C, 8A, map 1

**HOLOTYPE:** Male: **VANUATU:** Shefa: Shepherd Group: Tongariki Island, 16.8°S 168.5°E, 0–300 m, 29 Aug 1979, W.C.






Fig. 7. Left and right parameres of *Coridromius* spp.: A–C: *C. bicolor*. D–G: *C. boianotum*. H–J: *C. bulbopella*. K–M: *C. chenopoderis*. N–P: *C. chinensis*. Q–S: *C. confusus*.



Sue Lindsay, Australian Museum

Fig. 8. Scanning electron micrographs: **A.** *C. bicolor* ♂ genitalia, posterior view. **B.** *C. boianotum* ♂ genitalia, posterior view. **C–F:** *C. bulbopella*: **C.** ♀ dorsolateral view, right hindwing and hemelytron removed to expose paragenital "cup and bulb" (arrow). **D.** Detail of "cup and bulb". **E.** ♀ thorax and abdomen, right lateral view. **F.** ♂ genitalia, dorsal view. **G.** *C. carinatus* ♂ genitalia, dorsal view. **H.** *C. chenopoderis* ♂ genitalia, posterior view. FG = folded groove on pygophore. LP = left paramere, MES = mesepimeron, MET = metepimeron P = peritreme, RP = right paramere.



Map 1. Distribution map of *C. bicolor*, *boianotum*, *tahitiensis*, and *variegatus*.

Gagne, *Macaranga* sp. (Euphorbiaceae) (AMNH\_PBI 00041470) (BPBM type# 16724) (BPBM).

**DIAGNOSIS:** Recognized by the following characters: flat and broad shape (fig. 4); posterior margin of pygophore with shallow folded groove to left (fig. 8A); short left paramere (figs. 7A, 8A); small, triangular right paramere with rounded apex (fig. 7B-C); most specimens dark brown with the anterior half of the hemelytra pale. *C. bicolor* is similar in appearance to *variegatus*, *chenopoderis*, *monotocopsis*, and *pilbarensis* but is noticeably wider across the hemelytra, has a slightly flatter pronotum, different coloration, and a unique right paramere shape.

**DESCRIPTION: COLORATION** (figs. 4, 6): Orange-brown to dark chocolate brown. Head: Mostly yellow- to orange-brown; frons with dark brown vittae; paired transverse depressions and midpoint of vertex dark brown, in some cases these markings join together as a single band; vertex sometimes with yellow piping along posterior margin; clypeus with two submedial, vertical brown stripes; mandibular and maxillary plates yellow to orange, buccula brown to orange with yellow ventral margin; labrum dark brown, labium orange-brown, brown apically. Antennae: AI yellow, apex sometimes with faint brown ring; AII mostly pale with dark brown annulation; AIII and AIV dark brown. Thorax: Pronotal collar yellow, anterior margin brown; pronotum orange-brown to dark brown with faint medial stripe, collar region sometimes mottled with faint, coalescing brown spots, anterolateral

corners and humeral angles generally paler; mesoscutum yellow- to orange-brown; scutellum yellow- or orange-brown to dark brown, sometimes with a faint brown medial stripe, apex generally cream colored; thoracic pleura yellow- or orange-brown with posterior margins of each sclerite yellowed. Hemelytra: Mostly dark brown with the anterior half of the hemelytra sometimes pale, or mostly yellow to orange-brown with claval commissure dark brown and corium with faint symmetrical dark markings; the clavus always darker, sometimes somewhat reddened at costal fracture and apex of cuneus. Legs: Yellow-orange; metafemur with 7-8 diagonal brown stripes. Abdomen: yellow to brown. **SURFACE AND VESTITURE** (figs. 4, 6): Head and pronotum shallowly punctate; propleuron, metepimeron, scutellum and hemelytra impunctate; dorsum with sparse distribution of short, white, decumbent setae. **STRUCTURE:** (figs. 4, 6) Head: Frons broadly tumescent medially; vertex with two slightly raised tubercles adjacent to eyes, bordered posteriorly by shallow depressions, posterior margin somewhat carinate, slightly rounded. Thorax: Pronotum very broad, weakly convex, margins carinate; anterolateral margins only weakly upturned, submarginal region of humeral angles weakly excavate, collar region undifferentiated; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared. Hemelytra: Costal margin thin and platelike, flared along entire length and weakly sinuate. Legs: Metatibial spines thin and short. Abdomen: When

viewed laterally, posterior margin of abdominal sternite II not sharply angular. MALE GENITALIA (figs. 7A–C, 8A): Right paramere somewhat triangular, curved, and rounded (fig. 7B–C); left paramere sickle-shaped, with a broad gutter on inner surface (figs. 7A; 8A); posterior margin of pygophore with shallow groove on left formed by folding of the pygophore margin (fig. 8A). FEMALE PARAGENITALIA: No visible external paragenitalia.

**ETYMOLOGY:** Named for the coloration typical of most specimens.

**HOST:** This species has been collected on *Macaranga* and *Mallotus* spp. (Euphorbiaceae), as well as from mango blossoms (Anacardiaceae).

**DISTRIBUTION:** Known only from Vanuatu (map 1).

**PARATYPES:** VANUATU: Shefa: Shepherd Group: Tongariki Island, 16.8°S 168.5°E, 0–300 m, 29 Aug 1979, W.C. Gagne, *Macaranga* sp. (Euphorbiaceae), 2♂ (AMNH\_PBI 00041471, AMNH\_PBI 00041472), 2♀ (AMNH\_PBI 00041473, AMNH\_PBI 00041474) (BPBM). Torba: Banks Islands: Vanoua Lava Island, 2 km W of Sola, 13.83333°S 167.4667°E, 0–30 m, 17 Sep 1979, G.A. Samuelson, *Mallotus* sp. (Euphorbiaceae), 6♂ (AMNH\_PBI 00042134–AMNH\_PBI 00042139), 1♀ (AMNH\_PBI 00041475) (BPBM).

**OTHER SPECIMENS EXAMINED:** VANUATU: Shefa: Shepherd Group: Tongariki Island, 16.8°S 168.5°E, 0–300 m, 29 Aug 1979, W.C. Gagne, *Mangifera* sp. (Anacardiaceae), 2 nymphs (AMNH\_PBI 00042140, AMNH\_PBI 00042141) (BPBM).

#### *Coridromius boianotum*, new species

Figures 4, 6, 7D–G, 8B, map 1


**HOLOTYPE:** Male: SOLOMON ISLANDS: Malaita: Andalimu-Ngarafata (SW Fiu River), 8.71666°S 160.68333°E, 1 m, 19 Sep 1957, J.L. Gressitt, *Acalypha* sp. (Euphorbiaceae) (AMNH\_PBI 00041490) (BPBM type# 16725) (BPBM).

**DIAGNOSIS:** Distinguished by the flared anterolateral margins of the pronotum (fig. 4). *C. boianotum* shares similar coloration with *bulbopella* and *ruwenzorii*, but can be readily distinguished from these by its

unilobate proepisternum and lack of visible paragenitalia. It is also similar to *carinatus*, though with much less pronounced anterolateral projections of the pronotum. Both species have similar coloration and texture, a weakly curved left paramere, and a lack of any groove, fold, or suture along the posterior margin of the pygophore.

**DESCRIPTION: COLORATION** (figs. 4, 6): Head: Pale yellow-orange; frons without vittae; labium pale, brown apically. Antenna: Pale, apex of AII slightly brown. Thorax and abdomen: Concolorous yellow-orange, paler ventrally; scutellum with two prominent lateral brown spots. Legs: Concolorous yellow-orange, metafemur without dark diagonal stripes. **SURFACE AND VESTITURE** (figs. 4, 6): Head smooth and impunctate; pronotum and scutellum glossy with deep, widely spaced punctuation; hemelytra, propleuron, and metepimeron impunctate; setae on dorsum short, white, and simple, most setae knocked off examined specimens. **STRUCTURE:** (figs. 4, 6) Head: Frons and vertex forming a confluent broad medial tumescence, becoming shallowly depressed adjacent to eyes; vertex without tubercles adjacent to eyes; posterior margin of head medially rounded, not carinate, without transverse depressions adjacent to eyes. Thorax: Pronotum broad, lateral and posterior margins thin and carinate, anterolateral margins strongly flared and lobed, submarginal region of humeral angles weakly excavated, posterior margin weakly upturned, collar region weakly defined; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared; scutellum swollen. Hemelytra: Costal margin evenly rounded and thinly carinate. Legs: Metatibial spines short and thin. Abdomen: When viewed laterally, abdominal sternite II not angular. **MALE GENITALIA** (figs. 7D–G, 8B): Right paramere small, broadly triangular (fig. 7E–G); left paramere weakly curved (figs. 7D, 8B); ventral margin of pygophore simple and weakly biconvex, without fold, groove, or mesal suture on left side (fig. 8B). **FEMALE PARAGENITALIA:** No visible external paragenitalia.

**ETYMOLOGY:** The epithet *boianotum* is derived from Latin (*boia*, “collar”) and



Map 2. Distribution map of *C. bulbopella*, *lestoni*, and *ruwenzorii*.

Greek (*noton*, “back”) and reflects the expanded anterolateral projections of the pronotum.

**HOST:** Known only from the genus *Pipturus* (Urticaceae).

**DISTRIBUTION:** Known from the Solomon Islands and Vanuatu (map 1).

**PARATYPES: SOLOMON ISLANDS:**  
**Guadalcanal:** Roroni, 35 km E. of Honiara, 9.45°S 160.23333°E, 10 m, 08 May 1964, R. Straatman, 1♂ (AMNH\_PBI 00041492) (BPBM). Malaita: Auki, 8.767°S 160.7°E, 21 Sep 1957, J.L. Gressitt, *Pipturus* sp. (Urticaceae), 1♀ (AMNH\_PBI 00041491) (BPBM). **Western Province: New Georgia Islands:** New Georgia Island, Munda, 8.31666°S 157.25°E, 15 Jul 1959, J.L. Gressitt, 1♂ (AMNH\_PBI 00041493), 1♀ (AMNH\_PBI 00041494) (BPBM). **VA-NUATU: Shefa: Epi Island:** Lowekewou, 16.6°S 168.1667°E, 31 Aug 1979, W.C.

Gagne, (Urticaceae), 1♀ (AMNH\_PBI 00041506) *Pipturus* sp. (Urticaceae), 6♂ (AMNH\_PBI 00041495–AMNH\_PBI 00041500), 3♀ (AMNH\_PBI 00041501–AMNH\_PBI 00041503) (BPBM). Vaemali, 16.567°S 168.183°E, 08 Aug 1967, J. & M. Sedlacek, 2♀ (AMNH\_PBI 00041504, AMNH\_PBI 00041505) (BPBM).

*Coridromius bulbopella*, new species  
 Figures 4, 6, 7H–J, 8C–F, map 2

**HOLOTYPE:** Male: **SOUTH AFRICA:**  
**KwaZulu-Natal:** Ngome Forest Station, 27°S 31°E, 1100 m, 05–08 Feb 1996, M. Stiller, *Berkheya bipinnatifida* (Asteraceae) (AMNH\_PBI 00186395) (SANC).

**DIAGNOSIS:** Easily identified by a single twist in the male’s left paramere and the “cup and bulb” paragenital structure on the right abdominal laterotergites II and III of both

females and males. *C. bulbopella* is most similar to *ruwenzorii*, however in the latter the “cup and bulb” structure is much reduced.

**DESCRIPTION: COLORATION** (figs. 4, 6): Head: Concolorous brownish yellow, slightly paler ventrally; vertex without brown vittae. Antenna (fig. 4): Mostly yellow; AI with wide brown subapical band; AII with brown annulation; AIII and AIV brown. Thorax: Pronotal collar yellow; pronotum orange-yellow with pale yellow medial stripe, edges somewhat pale; mesoscutum orange-yellow; scutellum orange-yellow to yellow with two large brown markings; thoracic pleura orange-yellow, becoming yellow along posterior margins. Hemelytra: Mostly yellow- to orange-brown, clavus somewhat darker, becoming more brown along claval commissure and adjacent to scutellum; corium with faint brown transverse chevron pattern; membrane pale brown with darker brown veins. Abdomen: Orange-yellow. Legs: Orange-yellow; metafemur without transverse stripes. **STRUCTURE AND VESTITURE** (figs. 4, 6, 8E): Head: Frons broadly tumescent; vertex without tubercles adjacent to eyes; posterior margin of head distinctly rounded. Thorax: Pronotum broad and evenly rounded, margins thinly carinate, anterolateral margins adjacent to callar region depressed, submarginal region of humeral angles weakly excavated, posterior margin weakly upturned, callar region weakly developed; proepisternum bilobed; posterior margin of metepimeron truncate (fig. 8E); metanotum prominent and flared; scutellum not greatly swollen. Hemelytra: Embolium flared posteriorly, carinate, cuneus small. Legs: Metatibial spines long and thick. Abdomen: When viewed from the side, posterior margin of abdominal sternite II not sharply angular (fig. 8E). **MALE GENITALIA** (figs. 7H–J, 8F): Right paramere boot-shaped (figs. 7I–J, 8F); left paramere tightly coiled once basally, very long, thin and scythe-shaped, evenly curved and slightly twisted along its axis to apex (figs. 7H, 8F);

posterior margin of pygophore weakly bi-convex, without fold, groove, mesal suture, or ventral apical process (fig. 8F). **FEMALE PARAGENITALIA** (fig. 8C–D): Right laterotergites II and III modified into cup-and-bulb-shaped structure, also present in males.

**ETYMOLOGY:** The species name *bulbopella* combines the Latin *bulbus* (“bulb”) with the Greek *pella* (“wooden cup”), and refers to the cup-and-bulb-shaped paragenital structure of this species.

**REMARKS:** *C. bulbopella* is the only species of *Coridromius* in which external paragenitalic structures are found in both sexes. Nothing is known of the function of these unusual structures.

**HOST:** Collected on *Berkheya bipinnatifida* (Asteraceae).

**DISTRIBUTION:** Found in southeastern Africa (map 2).

**PARATYPES: SOUTH AFRICA: KwaZulu-Natal:** Ngome Forest Station, 27°S 31°E, 1100 m, 05–08 Feb 1996, M. Stiller, *Berkheya bipinnatifida* (Asteraceae), 2♂ (AMNH\_PBI 00041476, AMNH\_PBI 00186396), 4♀ (AMNH\_PBI 00186397–AMNH\_PBI 00186400) (SANC).

**OTHER SPECIMENS EXAMINED: KENYA: Central Province:** Limuru, 1.1°S 36.65°E, 2240 m, Sep 1954, Unknown, 10♂ (AMNH\_PBI 00041480–AMNH\_PBI 00041489), 3♀ (AMNH\_PBI 00041477–AMNH\_PBI 00041479) (BPBM). **TANZANIA: East Usambara,** Amani, 5.36667°S 39.71667°E, 1000 m, 06 Feb 1977, H. Enghoff, O. Lomholdt & O. Martin, 1♀ (AMNH\_PBI 00000261) (ZMUC).


***Coridromius carinatus* (Carvalho 1956),  
new combination**

Figures 4, 6, 8G, map 3

*Coridromoides carinatus* Carvalho, 1956: 54–55 (n. sp.).

**DIAGNOSIS:** Recognized by the swollen callar region, anterior projections of the humeral angles (fig. 4) and stylate eyes (figs. 4, 6). *C. carinatus* is most similar to *boianotum*, with both species sharing similar pronotal projections, coloration, texture, and male genitalia. Female unknown.


**REDESCRIPTION: COLORATION** (figs. 4, 6): Head: Pale brownish-orange; frons without brown vittae. Antenna: Pale yellow-



Map 3. Distribution map of *C. carinatus*, *confusus*, *crassus*, *ephippius*, and *marmoreus*.

orange, apex of AI and AII, base of AIII, and all of AIV slightly brown. Thorax: Mostly brownish-orange, paler ventrally; pronotal collar yellowish orange, brown laterally; pronotum brownish orange with pale medial stripe bordered by darker brownish-orange fields; mesoscutum dark brownish orange; scutellum brownish orange with two broad lateral brown spots. Hemelytra: Somewhat translucent brownish yellow, becoming darker orange on clavus, especially along scutellar margin. Legs: Yellowish orange, metafemur with five or six light brown diagonal stripes. Abdomen: Brownish yellow. SURFACE AND VESTITURE (figs. 4, 6): Head and scutellum smooth, glossy and impunctate; pronotum glossy and highly sculptured with deep, irregular punctures; propleuron, metepimeron, and hemelytra impunctate; head and dorsum clothed in

short, white, decumbent setae. STRUCTURE (figs. 4, 6): Head: Wide, eyes stylate; frons broadly tumescent medially; vertex without paired tumescences adjacent to eyes; posterior margin of head strongly rounded medially and carinate along lateral margins. Thorax: Pronotum highly sculptured and campanulate, anterolateral margins thin and carinate, projecting forward as distinct lobes, collar region greatly swollen, posterior margin flared upward and strongly carinate, humeral angles broadly rounded; proepisternum unilobed; posterior margin of metepimeron truncate; scutellum swollen. Hemelytra: Costal margin straight and narrowly carinate; cuneus narrow and long. Legs: Metatibial spines short and thin. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not angular. MALE GENITALIA (fig. 8G): Right para-



Map 4. Distribution map of *C. chenopoderis*, *monotocopsis* and *pilbarensis*. (Not shown on map: *C. chenopoderis* is also found in Hawaii and parts of southern North America.)

mere simple and rounded; left paramere thin and weakly curved; posterior margin of pygophore without groove, fold, or mesal suture. FEMALE PARAGENITALIA: Unknown.

**REMARKS:** Although the pronotal shape and stylate eyes of *carinatus* are unique to this species, all other characters—including the male genitalia—are typical of *Coridromius*. We do not find that this species differs enough from *Coridromius* to justify its own genus, and therefore synonymize *Coridromoides* as a junior synonym of *Coridromius*.

**HOST:** Unknown.

**DISTRIBUTION:** Known only from a single specimen from Palau (map 3).

**HOLOTYPE:** Male: **PALAU:** Northwest Auluptagel Island, Pacific Science Building, 7.30833°N 134.4778°E, 25 m, 13 Dec 1952, J.L. Gressitt, Light Trap (AMNH\_PBI 00185355) (USNM).

*Coridromius chenopoderis*, new species  
Figures 2B–C, 4, 6, 7K–M, 8H, map 4

**HOLOTYPE:** Male: **AUSTRALIA: Western Australia:** 13.2 km N of jct of Agana Kalbarri Rd and Brand Hiway, Galena River Bridge, 27.82917°S 114.6873°E, 190 m, 24

Oct 2004, Cassis, Wall, Weirauch, Symonds, *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6988431 (AMNH\_PBI 00015692) (AM).

**DIAGNOSIS:** Recognized by the following combination of characters: proepisternum unilobed; metanotum not prominently flared; posterior margin of pygophore biconvex, with prominent fold on left side forming deep U-shaped groove, without small apophysis on right margin of groove. *C. chenopoderis*, *monotocopsis*, *pilbarensis*, and *variegatus* are all very similar and often difficult to tell apart. Of the four species, *chenopoderis* exhibits the most phenotypic variation in both size and coloration (e.g., fig. 2B–C) and has the widest variety of known host plants of all *Coridromius*. It has a somewhat narrower body shape and is typically smaller than the others, though size ranges do overlap. Males can be distinguished from the other species by the relatively shorter left paramere, the much deeper U-shaped fold on the posterior margin of the pygophore, by the lack of a small apical process on the right margin of this fold (found only in *pilbarensis*) and by the lack of a small rounded lobe on the posterior margin of the metepimeron (as in fig. 14B) as found in *variegatus*.

**DESCRIPTION: COLORATION** (figs. 2B–C, 4, 6): Extremely variable, ranging from light tan or olive to brown or black with brown markings (fig. 2B–C). Head: Mostly brownish yellow to orange-brown with dark brown vittae on frons; lateral margins bordering eyes and posterior margin of vertex yellow; paired tubercles on vertex yellow, bordered posteriorly with dark brown markings, sometimes merging with dark brown spot in middle of vertex; clypeus with irregular median and lateral dark brown markings; mandibular and maxillary plates darkened along ventral margins; buccula pale; labrum light brown; labium yellow, brown apically; in some specimens head mostly dark brown. Antenna: AI yellow-brown to orange-brown with thin brown apical and basal annulations; AII yellow-brown to orange-brown, sometimes slightly brown apically; AIII and AIV orange-brown to brown. Thorax: Pronotal collar yellow with brown anterior and posterior margins, uniformly dark brown in darker specimens; pronotum light tan, olive, brown, or dark brown, often with pale medial stripe, usually with irregular dark brown or black markings on collar region; mesoscutum orange-brown to dark brown; scutellum ranging from mostly dark brown with yellow corners to mostly light brown or tan with darker brown markings; thoracic pleura light brown to dark brown, often with yellow markings; mesopleuron brown to black ventrally. Hemelytra: Light tan or olive to brown or black, sometimes with a brown chevron on corium, sometimes darker along claval commissure; membrane faint brown with darker veins. Legs: Coxae yellow to brown; pro- and mesofemora yellow to dark brown, sometimes with faint, subapical brown markings; hind femur yellow to orange-brown with 8 or 9 dark brown diagonal stripes on outer surface; tibiae yellow to brown; tarsi brown apically. Abdomen: Yellow to dark brown or black, sometimes with lateral brown markings around spiracles, males sometimes with reddish markings on genital capsule. **SURFACE AND VESTITURE** (figs. 4, 6): Head glossy and shallowly punctate; pronotum shallowly punctate; propleuron, metepimeron, scutellum, and hemelytra impunctate; head and dorsum with sparse distribution of

short, simple, white, decumbent setae. **STRUCTURE** (figs. 4, 6): Head: Wide; frons broadly tumescent medially; paired tubercles adjacent to eyes; transverse depressions above tubercles shallow; posterior margin of head slightly rounded medially but otherwise flat. Thorax: Pronotum broad and weakly rounded, margins thinly carinate, anterolateral margins weakly upturned, posterior margin weakly medially cleft, submarginal region of humeral angles weakly excavated; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared. Hemelytra: Embolium thin and evenly rounded, nearly straight. Legs: Metatibial spines thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. **MALE GENITALIA** (figs. 7K–M, 8H): Right paramere triangular and elongate (figs. 7L–M); left paramere sickle-shaped, apex acute (figs. 7K, 8H); posterior margin of pygophore biconvex, with distinct fold on left forming a deep U-shaped groove (fig. 8H). **FEMALE PARAGENITALIA:** No visible external paragenitalia.

**ETYMOLOGY:** This name was chosen to reflect this species' preference for chenopodiaceous plants.

**HOSTS:** This species is known from several plant families but is most commonly collected on chenopods (Chenopodiaceae).

**DISTRIBUTION:** The most widely distributed of the Australian species, *chenopoderis* is found throughout the country. It has also been collected in New Zealand and Norfolk Island, where it has probably been introduced (map 4). This species has also recently been discovered in North America by Michael D. Schwartz; it has been collected off chenopods in Mexico, Arizona, New Mexico, southern California, Florida, and Texas. Additionally, specimens of *Coridromius* have been collected from Hawaii and previously identified as *variegatus*. Examination of these specimens, collected from *Chenopodium murale*, confirms that they are in fact *chenopoderis*.

**PARATYPES: AUSTRALIA: New South Wales:** 2.9 km W of Nyangan, 31.55001°S 147.1553°E, 202 m, 18 Oct 2001, Cassis, Silveira, Wall, *Einadia nutans* subsp. *nutans* (R.Br.) A. J. Scott (Chenopodiaceae), det.

NSW staff NSW658277, 7♂ (AMNH\_PBI 00015526–AMNH\_PBI 00015532), 3♀ (AMNH\_PBI 00012117–AMNH\_PBI 00012119) *Rhagodia spinescens* R.Br. (Chenopodiaceae), det. NSW staff NSW658274, 1♂ (AMNH\_PBI 00012105), 5♀ (AMNH\_PBI 00012106–AMNH\_PBI 00012110) (AM). 42 km SW Bourke toward Louth, 30.33334°S 144.5833°E, 100 m, 27 Oct 1995, Schuh and Cassis, *Flindersia maculosa* (Lindley) Benth. (Rutaceae), det. P. H. Weston 1996 NSW395939, 1♂ (AMNH\_PBI 00002613), 5♀ (AMNH\_PBI 00002614–AMNH\_PBI 00002618) (AMNH). 9.5 km E of Balranald on Sturt Hwy, 34.702°S 143.615°E, 20 Oct 1996, Schuh and Cassis, *Nitraria billardierei* DC. (Zygophyllaceae), det. Perth staff PERTH05095247, 16♂ (AMNH\_PBI 00006354–AMNH\_PBI 00006369), 7♀ (AMNH\_PBI 00006370–AMNH\_PBI 00006376) *Nitraria billardierei* DC. (Zygophyllaceae), det. Perth staff PERTH05095247, 1♂ (AMNH\_PBI 00123417) (AM). *Atriplex nummularia omissa* Aellen (Chenopodiaceae), det. Perth staff PERTH-05054680, 2♂ (AMNH\_PBI 00002699, AMNH\_PBI 00002700), 1♀ (AMNH\_PBI 00002701) (AMNH). Comerong Island Nature Reserve, nr. Nowra, Seven Mile Beach, on dunes, 34.882°S 150.735°E, 09 Apr 2006, C. Symonds, (Chenopodiaceae), 5♂ (AMNH\_PBI 00012442–AMNH\_PBI 00012446), 3♀ (AMNH\_PBI 00012447–AMNH\_PBI 00012449) (AM). Myall Lakes NP, 32.57916°S 152.29083°E, 16 Nov 1996, L. Wilkie, *Acacia longifolia* (Mimosaceae), 2♂ (AMNH\_PBI 00012401, AMNH\_PBI 00012402) (AM). Warrumbungle National Park, Wambelong Campground, 31.21666°S 149.08333°E, 550 m, 25 Oct 1995, Schuh and Cassis, *Amyema cambagei* (Blakely) Danser (Loranthaceae), det. B.M. Wiecek 1996 NSW395934, 1♀ (AMNH\_PBI 00002731) (AMNH). Northern Territory: 143.3 km NW of Bond Springs on Tanami Rd, 23.03333°S 132.722°E, 573 m, 22 Oct 2001, Cassis, Schuh, Schwartz, Silveira, Wall, *Rhagodia spinescens* R.Br. (Chenopodiaceae), det. NSW staff NSW658306, 2♂ (AMNH\_PBI 00012120, AMNH\_PBI 00012121), 1♀ (AMNH\_PBI 00012122) (AM). **Queensland:** 45 km N of Quilpie, 26.3423°S 144.3078°E, 280 m, 02 Nov 1998, Schuh, Cassis, Silveira, *Muehlenbeckia florulenta* Meissner (Polygonaceae), det. NSW staff NSW427471, 2♂

(AMNH\_PBI 00002517, AMNH\_PBI 00002518), 2♀ (AMNH\_PBI 00002523, AMNH\_PBI 00002524) (AMNH). 82.6 km NW of Quilpie, 26.3479°S 143.6454°E, 190 m, 03 Nov 1998, Schuh, Cassis, Silveira, *Frankenia* sp. (Frankeniaceae), det. NSW staff NSW427347, 1♀ (AMNH\_PBI 00012386) (AM). 9 km WSW of Adavale, 25.91094°S 144.5143°E, 270 m, 02 Nov 1998, Schuh, Cassis, Silveira, *Hakea leucoptera* R. Br. (Proteaceae), det. NSW staff NSW427661, 8♂ (AMNH\_PBI 00006203–AMNH\_PBI 00006210), 8♀ (AMNH\_PBI 00006211–AMNH\_PBI 00006214, AMNH\_PBI 00006280–AMNH\_PBI 00006283) (AM). *Hakea leucoptera* R. Br. (Proteaceae), det. NSW staff NSW427661, 8♂ (AMNH\_PBI 00003054–AMNH\_PBI 00003061), 11♀ (AMNH\_PBI 00002751, AMNH\_PBI 00003062–AMNH\_PBI 00003071) (AMNH). 9.6 km W of Mitchell, 26.49202°S 147.8728°E, 430 m, 31 Oct 1998, Schuh, Cassis, Silveira, 2♂ (AMNH\_PBI 00012382, AMNH\_PBI 00012383), 1♀ (AMNH\_PBI 00012384) (AM). 91 km N of Quilpie, 25.99847°S 144.4098°E, 300 m, 02 Nov 1998, Schuh, Cassis, Silveira, *Amyema quandang* var. *quandang* (Lindley) Tieghem (Loranthaceae), det. NSW staff NSW427341, 1♀ (AMNH\_PBI 00006117) *Hakea leucoptera* (Proteaceae), 13♂ (AMNH\_PBI 00006085–AMNH\_PBI 00006097), 20♀ (AMNH\_PBI 00006098–AMNH\_PBI 00006116, AMNH\_PBI 00012387) *Hakea leucoptera* R. Br. (Proteaceae), det. NSW staff NSW 427661, 1♂ (AMNH\_PBI 00123415), 1♀ (AMNH\_PBI 00123420) (AM). *Hakea leucoptera* R. Br. (Proteaceae) NSW 427661, 13♂ (AMNH\_PBI 00003175–AMNH\_PBI 00003187), 27♀ (AMNH\_PBI 00003188–AMNH\_PBI 00003214) (AMNH). **South Australia:** 11.5 km NE of Wooltana Homestead, 30.33726°S 139.4934°E, 180 m, 06 Nov 1998, Schuh, Cassis, Silveira, 2♂ (AMNH\_PBI 00012388, AMNH\_PBI 00012389), 3♀ (AMNH\_PBI 00012390–AMNH\_PBI 00012392) (AM). 12.5 km E of Martins Well, 31.50001°S 139.2276°E, 162 m, 09 Nov 2001, Cassis, Schuh, Schwartz, *Acacia oswaldii* F.Muell. (Fabaceae), det. NSW staff NSW666370, 3♂ (AMNH\_PBI 00012111–AMNH\_PBI 00012113), 5♀ (AMNH\_PBI 00012114–AMNH\_PBI 00012116) (AM). 15 km S of Bews, 35.48474°S 140.4332°E, 130 m, 08 Nov 1998, Schuh, Cassis, Silveira, Light Trap, 1♂ (AMNH\_PBI 00123407)

(AM). 17 km E of Nepebunna, Gammon Ranges National Park, 30.57312°S 139.1273°E, 480 m, 07 Nov 1998, Schuh, Cassis, Silveira, *Rhagodia parabolica* R. Br. (Chenopodiaceae), 13♂ (AMNH\_PBI 00006284-AMNH\_PBI 00006296), 20♀ (AMNH\_PBI 00006297-AMNH\_PBI 00006316) *Rhagodia parabolica* R. Br. (Chenopodiaceae), 1♂ (AMNH\_PBI 00123416) (AM). 17.6 km S of Innamincka, 27.88068°S 140.6712°E, 130 m, 06 Nov 1998, Schuh, Cassis, Silveira, *Chenopodium auricomum* Lindley (Chenopodiaceae), det. NSW staff, 42♂ (AMNH\_PBI 00006268-AMNH\_PBI 00006279, AMNH\_PBI 00006460-AMNH\_PBI 00006474, AMNH\_PBI 00012039-AMNH\_PBI 00012053), 34♀ (AMNH\_PBI 00006317-AMNH\_PBI 00006350) *Teucrium racemosum* R. Br. (Lamiaceae), det. NSW staff NSW427991, 1♂ (AMNH\_PBI 00012055) *Muehlenbeckia florulenta* Meissner (Polygonaceae), det. NSW staff NSW427467, 3♂ (AMNH\_PBI 00012054, AMNH\_PBI 00012374-AMNH\_PBI 00012375), 7♀ (AMNH\_PBI 00012376-AMNH\_PBI 00012381, AMNH\_PBI 00012393) (AM). 20 km W of Nepabunna, Mt. Serle, 30.55365°S 138.8304°E, 630 m, 07 Nov 1998, Schuh, Cassis, Silveira, 1♀ (AMNH\_PBI 00012385) (AM). 41.5 km NW of Morgan, 33.63335°S 139.9167°E, 150 m, 01 Nov 1995, Schuh, Cassis, and Gross, *Exocarpos aphyllus* R. Br. (Santalaceae), det. B.M. Wiecek 1996 NSW395968, 5♂ (AMNH\_PBI 00002630-AMNH\_PBI 00002634), 6♀ (AMNH\_PBI 00002635-AMNH\_PBI 00002640) (AMNH). 5 km N Yunta toward Arkaroola, 32.53334°S 139.55°E, 250 m, 29 Oct 1995, Schuh and Cassis, undetermined sp. (Chenopodiaceae), det. R.T. Schuh NSW395955, 6♂ (AMNH\_PBI 00002465-AMNH\_PBI 00002470), 9♀ (AMNH\_PBI 00002752-AMNH\_PBI 00002760) (AMNH). 5.4 km N of Yunta, 32.53804°S 139.5598°E, 300 m, 08 Nov 1996, Schuh and Cassis, *Acacia* sp. (Fabaceae), 4♂ (AMNH\_PBI 00006183-AMNH\_PBI 00006186), 16♀ (AMNH\_PBI 00006187-AMNH\_PBI 00006202) (AM). 51 km NW of Morgan, 33.58334°S 140°E, 150 m, 01 Nov 1995, Schuh, Cassis, and Gross, Light Trap, 2♂ (AMNH\_PBI 00012403, AMNH\_PBI 00123404), 2♀ (AMNH\_PBI 00123405, AMNH\_PBI 00123406) (AM). *Eremophila mitchellii* Benth. (Myoporaceae), det. J. Everett 1996 NSW395936, 2♂ (AMNH\_PBI 00002713, AMNH\_PBI 00002714), 2♀

(AMNH\_PBI 00002715, AMNH\_PBI 00002716), Light Trap, 4♂ (AMNH\_PBI 00002709-AMNH\_PBI 00002712), 2♀ (AMNH\_PBI 00002641, AMNH\_PBI 00002642) (AMNH). 52 km SW of Yunta, 32.83335°S 139.1°E, 500 m, 30 Oct 1995, Schuh and Cassis, 24♂ (AMNH\_PBI 00006383-AMNH\_PBI 00006406), 15♀ (AMNH\_PBI 00006407-AMNH\_PBI 00006421) (AM). 58♂ (AMNH\_PBI 00002487-AMNH\_PBI 00002499, AMNH\_PBI 00003004-AMNH\_PBI 00003017, AMNH\_PBI 00003072-AMNH\_PBI 00003102), 43♀ (AMNH\_PBI 00002500-AMNH\_PBI 00002516, AMNH\_PBI 00003018-AMNH\_PBI 00003035, AMNH\_PBI 00003103-AMNH\_PBI 00003110) (AMNH). 6 km N of Sturt Hiway, Mt. Mary Road, 34.33334°S 139.4167°E, 250 m, 01 Nov 1995, Schuh, Cassis, and Gross, *Nitaria billardierei* DC. (Nittrariaceae), det. R.G. Coveny 1996 NSW395966, 3♂ (AMNH\_PBI 00006171-AMNH\_PBI 00006173), 5♀ (AMNH\_PBI 00006174-AMNH\_PBI 00006178) (AM). *Nitaria billardierei* DC. (Nittrariaceae), det. R.G. Coveny 1996 NSW395966, 5♂ (AMNH\_PBI 00002619-AMNH\_PBI 00002623), 6♀ (AMNH\_PBI 00002624-AMNH\_PBI 00002629) (AMNH). 66-69 km NW of Morgan, Cane Grass, 33.58334°S 140.0333°E, 150 m, 02 Nov 1995, Schuh, Cassis, and Gross, undetermined sp., 22♂ (AMNH\_PBI 00002658-AMNH\_PBI 00002679), 19♀ (AMNH\_PBI 00002680-AMNH\_PBI 00002698) (AMNH). 75 km NW of Morgan, 5 km N Cane Grass, 33.53334°S 140.05°E, 100 m, 02 Nov 1995, Schuh, Cassis, and Gross, Light Trap, 1♀ (AMNH\_PBI 00123408) (AM). 96 km NW of Morgan, Pine Valley Stn, 33.31667°S 140.2°E, 150 m, 02 Nov 1995, Schuh, Cassis, and Gross, *Exocarpos aphyllus* R. Br. (Santalaceae), det. B.M. Wiecek 1996 NSW395968, 1♀ (AMNH\_PBI 00002702) (AMNH). Brookfield Cons. Pk., 34.22°S 139.27°E, 25 Nov 1992, I. D. Naumann & J. C. Cardale, 2♂ (AMNH\_PBI 00006179, AMNH\_PBI 00006180), 2♀ (AMNH\_PBI 00006181, AMNH\_PBI 00006182) (AM). Nullabor National Park 5 km E of WA border, 31.63716°S 129.0346°E, 50 m, 22 Oct 1996, Schuh and Cassis, *Myoporum insulare*? R. Br. (Myoporaceae), det. Perth staff PERTH05095239, 4♂ (AMNH\_PBI 00012394-AMNH\_PBI 00012397), 2♀ (AMNH\_PBI 00012398, AMNH\_PBI 00012399) (AM). Scorpion

Springs Cons. Park, 35.60421°S 140.8646°E, 125 m, 10 Nov 1998, Schuh, Cassis, Silveira, *Melaleuca brevifolia* Turcz. (Myrtaceae), det. NSW staff NSW427362, 1♀ (AMNH\_PBI 00123400) (AM). Yunta, 32.58°N 139.559°E, 12 Nov 1994, C.N. and A.S. Smithers, 1♀ (AMNH\_PBI 00123409) (AM). **Victoria:** Little Desert National Park, 5–6 km W of McDonald Hiway, 36.61668°S 141.1667°E, 150 m, 03 Nov 1995, Schuh and Cassis, *Acacia acinacea* Lindl. (Fabaceae), det. B.J. Conn 1996 NSW 395983, 1♀ (AMNH\_PBI 00002643) (AMNH). **Western Australia:** 115.4 km E of Norseman, 32.05143°S 122.9675°E, 600 m, 23 Oct 1996, Schuh and Cassis, *Eremophila ionantha* Diels (Myoporaceae), det. Perth staff PERTH05056152, 2♂ (AMNH\_PBI 00002542, AMNH\_PBI 00002543) (AMNH). 13.2 km N of jct of Agana Kalbarri Rd and Brand Hiway, Galena River Bridge, 27.82917°S 114.6873°E, 190 m, 24 Oct 2004, Cassis, Wall, Weirauch, Symonds, *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6988431, 22♂ (AMNH\_PBI 00015693–AMNH\_PBI 00015714), 12♀ (AMNH\_PBI 00015715–AMNH\_PBI 00015726) (AM). 15 km S of Lake Grace, 33.2367°S 118.4692°E, 400 m, 07 Dec 1997, Schuh, Cassis, Brailovsky, Asquith, 1♀ (AMNH\_PBI 00002548) (AMNH). 2 km E of Nungarin on Rt 50, 31.43596°S 118.2627°E, 330 m, 16 Nov 1999, R.T. Schuh, G. Cassis, & R. Silveira, *Rhagodia preissii* Moq. (Chenopodiaceae), det. Perth staff PERTH05670594, 4♂ (AMNH\_PBI 00006450–AMNH\_PBI 00006453), 6♀ (AMNH\_PBI 00006454–AMNH\_PBI 00006459) (AM). 2.5 km W of Brand Hiway (off Red Emperor Rd), Flat Rocks Beach (S of Geraldton), 29.02061°S 114.7863°E, 22 Oct 2004, Cassis, Wall, Weirauch, Symonds, *Rhagodia preissii* subsp. *obovata* (Moq.) Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6989543, 1♂ (AMNH\_PBI 00012088), 1♀ (AMNH\_PBI 00012089) (AM). 22 km S of Watheroo, 30.43161°S 116.0438°E, 400 m, 02 Nov 1996, Schuh and Cassis, *Halosarcia indica* bidens (Nees) Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH05236843, 5♂ (AMNH\_PBI 00012100–AMNH\_PBI 00012104), 2♀ (AMNH\_PBI 00012425, AMNH\_PBI 00012426) (AM). *Scholtzia capitata* Benth. (Myrtaceae), det. Perth staff

PERTH05236622, 8♂ (AMNH\_PBI 00002473–AMNH\_PBI 00002477, AMNH\_PBI 00002601–AMNH\_PBI 00002603), 11♀ (AMNH\_PBI 00002478–AMNH\_PBI 00002484, AMNH\_PBI 00002604–AMNH\_PBI 00002607) *Halosarcia indica* bidens (Nees) Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH05236843, 12♂ (AMNH\_PBI 00002645–AMNH\_PBI 00002656), 1♀ (AMNH\_PBI 00002657) (AMNH). 27.6 km N of Coral Bay Rd on Cardabia-Ningaloo Rd, 22.90198°S 113.8167°E, 25 m, 29 Oct 2004, Cassis, Wall, Weirauch, Tatarnic, Symonds, *Salsola tragus* subsp. *tragus* L. (Chenopodiaceae), det. Field ID, 11♂ (AMNH\_PBI 00015622–AMNH\_PBI 00015632), 9♀ (AMNH\_PBI 00015633–AMNH\_PBI 00015641) (AM). 31.3 km N of Ravensthorpe, 33.36519°S 119.8792°E, 500 m, 05 Nov 1996, Schuh and Cassis, 1♂ (AMNH\_PBI 00002540), 1♀ (AMNH\_PBI 00002541) (AMNH). 31.7 km W of Agnew toward Sandstone, 27.96227°S 120.4277°E, 800 m, 26 Oct 1996, Schuh and Cassis, *Rhagodia preissii* Moq. (Chenopodiaceae), det. Perth staff PERTH05054753, 1♀ (AMNH\_PBI 00002526) (AMNH). 39 km NE of King River, 34.78937°S 118.2759°E, 200 m, 06 Nov 1996, Schuh and Cassis, *Agonis parviceps* Schauer (Myrtaceae), det. Perth staff PERTH05236827, 1♂ (AMNH\_PBI 00002522) (AMNH). 40 km N of Ravensthorpe, 33.31524°S 119.8151°E, 500 m, 05 Nov 1996, Schuh and Cassis, *Maireana oppositifolia* (F. Muell.) Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH-05236630, 7♂ (AMNH\_PBI 00006129–AMNH\_PBI 00006135), 18♀ (AMNH\_PBI 00006136–AMNH\_PBI 00006153) (AM). 42.5 km N of Kojonup, 33.46391°S 117.0579°E, 400 m, 04 Nov 1996, Schuh and Cassis, *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH-05236835, 2♂ (AMNH\_PBI 00012421, AMNH\_PBI 00012422), 13♀ (AMNH\_PBI 00015773–AMNH\_PBI 00015785) (AM). *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH05236835, 6♂ (AMNH\_PBI 00002581–AMNH\_PBI 00002586), 6♀ (AMNH\_PBI 00002587–AMNH\_PBI 00002592) (AMNH). *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH 05236835, 4♂ (AMNH\_PBI 00191191–AMNH\_PBI 00191194),

4♀ (AMNH\_PBI 00191195–AMNH\_PBI 00191198) (USNM). 43 km N of Norseman, 31.85648°S 121.6414°E, 300 m, 19 Nov 1999, R.T. Schuh, G. Cassis, & R. Silveira, *Olearia muelleri* (Sond.) Benth. (Asteraceae), det. Perth staff PERTH05670691, 8♂ (AMNH\_PBI 00002748, AMNH\_PBI 00003036–AMNH\_PBI 00003042), 13♀ (AMNH\_PBI 00002749–AMNH\_PBI 00002750, AMNH\_PBI 00003043–AMNH\_PBI 00003053) (AMNH). *Atriplex bunburyana* F. Muell. (Chenopodiaceae), det. Perth staff PERTH05671205, 2♂ (AMNH\_PBI 00114226, AMNH\_PBI 00114227) (CNC). 46.5 km W of Yalgoo, 28.41302°S 116.2151°E, 600 m, 27 Oct 1996, Schuh and Cassis, *Eremophila forrestii* F. Muell. (Myoporaceae), det. Perth staff PERTH05671205, 1♀ (AMNH\_PBI 00012424) (AM). 51 km N of Leeman on Indian Ocean Rd, Beekeepers Nature Reserve, 29.5116°S 114.997°E, 06 Nov 2004, Cassis, Weirauch, Tatarnic, Symmonds, *Rhagodia preissii* Moq. (Chenopodiaceae), det. Field ID, 1♂ (AMNH\_PBI 00012096), 1♀ (AMNH\_PBI 00012097) (AM). *Rhagodia preissii* Moq. (Chenopodiaceae), det. Field ID, 6♂ (AMNH\_PBI 00114217–AMNH\_PBI 00114222), 3♀ (AMNH\_PBI 00114223–AMNH\_PBI 00114225) (CNC). 53.9 km N of Kalgoorlie, 30.28882°S 121.2558°E, 600 m, 24 Oct 1996, Schuh and Cassis, 4♂ (AMNH\_PBI 00002717–AMNH\_PBI 00002720), 1♀ (AMNH\_PBI 00002721) (AMNH). 56.6 km W of Yalgoo, 28.42397°S 116.1233°E, 600 m, 27 Oct 1996, Schuh and Cassis, *Melaleuca uncinata* R.Br. (Myrtaceae), det. Perth staff PERTH05120640, 2♀ (AMNH\_PBI 00002520, AMNH\_PBI 00002521) *Dicrastylis fulva forma fulva* Harv. (Chloanthaceae), det. Perth staff PERTH05120632, 1♂ (AMNH\_PBI 00002519) (AMNH). 82.2 km W of Agnew toward Sandstone, 28.01057°S 119.9455°E, 650 m, 26 Oct 1996, Schuh and Cassis, *Alyogyne pinoniana* (Gaudich.) Fryxell (Malvaceae), det. Perth staff PERTH-05056101, 1♂ (AMNH\_PBI 00123418), 1♀ (AMNH\_PBI 00123419) (AM). Blowholes Rd NW of North West Coastal Hiway, Beagle Hill Area, 24.49068°S 113.4626°E, 20 m, 27 Oct 2004, Cassis, Wall, Weirauch, Tatarnic, Symonds, *Salsola tragus* subsp. *tragus* L. (Chenopodiaceae), det. Field ID, 1♂ (AMNH\_PBI 00012086), 1♀ (AMNH\_PBI 00012087) *Enchytraea tomentosa* var.

*tomentosa* R.Br. (Chenopodiaceae), det. Perth staff PERTH6988903, 3♂ (AMNH\_PBI 00015490–AMNH\_PBI 00015492) (AM). Brand Hiway 55.9 km S of Dongarra Road, 29.62934°S 115.2187°E, 100 m, 31 Oct 1996, Schuh and Cassis, *Scholtzia drummondii* Benth. (Myrtaceae), det. Perth staff PERTH-05120209, 2♂ (AMNH\_PBI 00002544, AMNH\_PBI 00002545) (AMNH). Burrup Conservation Area, near Karratha, 20.57805°S 116.79472°E, 17 m, 03 Jun 2004, N. Tatarnic and S. Lassau, *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6988431, 3♂ (AMNH\_PBI 00015727–AMNH\_PBI 00015729), 6♀ (AMNH\_PBI 00015730–AMNH\_PBI 00015735) (AM). Cape Arid National Park, Thomas River Campground, 33.8539°S 123.0126°E, 20 m, 24 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Pimelea ferruginea* Labill. (Thymelaeaceae), det. Perth staff 05672341, 1♂ (AMNH\_PBI 00002546) *Atriplex isatidea* Moq. (Chenopodiaceae), det. Perth staff PERTH05672317, 3♂ (AMNH\_PBI 00002740–AMNH\_PBI 00002742), 1♀ (AMNH\_PBI 00002744) (AMNH). Cape Naturaliste National Park, 33.54034°S 115.0123°E, 50 m, 14 Dec 1997, Schuh, Cassis, Brailovsky, *Rhagodia baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH05055318, 3♂ (AMNH\_PBI 00006422–AMNH\_PBI 00006424), 10♀ (AMNH\_PBI 00006422–AMNH\_PBI 00006431) (AM). *Rhagodia baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH05055318, 28♂ (AMNH\_PBI 00003124–AMNH\_PBI 00003151), 23♀ (AMNH\_PBI 00003152–AMNH\_PBI 00003174) (AMNH). Carnarvon, Small Boat Harbour, 24.90044°S 113.6477°E, 26 Oct 2004, Cassis, Wall, Weirauch, Tatarnic, Symonds, *Chenopodium murale* L. (Chenopodiaceae), det. Perth staff PERTH6989047, 1♂ (AMNH\_PBI 00012090), 1♀ (AMNH\_PBI 00012091) (AM). Conspicuous Beach, Walpole-Nornalup National Park, 10 km E of Nornalup, 35.03725°S 116.8443°E, 30 m, 17 Dec 1997, Schuh, Cassis, Brailovsky, *Rhagodia baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH-05095271, 9♂ (AMNH\_PBI 00006154–AMNH\_PBI 00006162), 5♀ (AMNH\_PBI 00006163–AMNH\_PBI 00006167) (AM). 2♀ (AMNH\_PBI 00002725, AMNH\_PBI 00002726) *Pimelea rosea* R.Br. (Thymelaeae-

ceae), det. Perth staff PERTH05099781, 1♀ (AMNH\_PBI 00002727) *Olearia axillaris* (DC.) Benth. (Asteraceae), det. Perth staff PERTH05095328, 2♂ (AMNH\_PBI 00002722, AMNH\_PBI 00002723), 1♀ (AMNH\_PBI 00002724) (AMNH). Cosy Corner Beach East, Torbay Sound, W of Albany, 35.06033°S 117.6446°E, 2 m, 01 Dec 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Allocasuarina humilis* (Otto and F. Dietr.) L.A.S. Johnson (Casuarinaceae), det. Perth staff PERTH05671345, 1♀ (AMNH\_PBI 00002547) (AMNH). Fitzgerald River National Park, East Myles Beach, 33.93293°S 120.0031°E, 30 m, 06 Dec 1997, Schuh, Cassis, Brailovsky, Asquith, *Pimelea ferruginea* Labill. (Thymelaeaceae), det. Perth staff PERTH05056403, 1♀ (AMNH\_PBI 00002525) (AMNH). Fowlers Camp (at end of Fowlers Camp Rd), Shark Bay World Heritage Area, 26.10801°S 113.6167°E, 24 Oct 2004, Cassis, Wall, Weirauch, Symonds, *Chenopodium murale* L. (Chenopodiaceae), det. Perth staff PERTH-6988016, 70♂ (AMNH\_PBI 00015796–AMNH\_PBI 00015865), 35♀ (AMNH\_PBI 00015866–AMNH\_PBI 00015900) *Salsola tragus* subsp. *tragus* L. (Chenopodiaceae), det. Perth staff PERTH6988024, 1♂ (AMNH\_PBI 00012093), 1♀ (AMNH\_PBI 00012094) (AM). Gladstone, 25.95428°S 114.2464°E, 04 Nov 2004, Cassis, Weirauch, Tatarnic, Symonds, *Salsola tragus* subsp. *tragus* L. (Chenopodiaceae), det. Field ID, 17♂ (AMNH\_PBI 00015593–AMNH\_PBI 00015609), 12♀ (AMNH\_PBI 00015610–AMNH\_PBI 00015621) (AM). Greenough River Mouth, 28.86304°S 114.6343°E, 05 Nov 2004, Cassis, Weirauch, Tatarnic, Symonds, *Rhagodia preissii* subsp. *obovata* (Moq.) Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6988377, 7♂ (AMNH\_PBI 00015493–AMNH\_PBI 00015499), 11♀ (AMNH\_PBI 00015500–AMNH\_PBI 00015510), 4 juveniles (AMNH\_PBI 00015511–AMNH\_PBI 00015514) (AM). Hammersley Inlet Campground, 33.95788°S 119.9146°E, 2 m, 28 Nov 1999, R.T. Schuh and G. Cassis, *Suaeda australis* (R.Br.) Moq. (Chenopodiaceae), det. Perth staff PERTH05671213, 7♂ (AMNH\_PBI 00006432–AMNH\_PBI 00006438), 11♀ (AMNH\_PBI 00006439–AMNH\_PBI 00006449) (AM). Leeuwin Naturaliste National Park, Canal Rocks, 33.66513°S 115.0165°E, 50 m, 15 Dec 1997, Schuh, Cassis, Brailovsky, Asquith, 1♂ (AMNH\_PBI 00002485), 1♀ (AMNH\_PBI 00002486) (AMNH). Mosman Park, Perth, 32.0209°S 115.7687°E, 20 m, 29 Nov 1998, G. Cassis, *Rhagodia baccata* (Chenopodiaceae), 4♂ (AMNH\_PBI 00006118–AMNH\_PBI 00006121), 3♀ (AMNH\_PBI 00006122–AMNH\_PBI 00006124) (AM); 05 Nov 1998, G. Cassis, *Rhagodia baccata baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH 05227283, 12♂ (AMNH\_PBI 00015902–AMNH\_PBI 00015913), 10♀ (AMNH\_PBI 00015914–AMNH\_PBI 00015923) (AM). Peak Charles National Park Campground, 32.88335°S 121.1703°E, 300 m, 20 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Rhagodia preissii preissii* Moq. (Chenopodiaceae), det. Perth staff PERTH05670713, 16♂ (AMNH\_PBI 00012056–AMNH\_PBI 00012071), 14♀ (AMNH\_PBI 00012072–AMNH\_PBI 00012085) *Melaleuca uncinata* R.Br. (Myrtaceae), det. Perth staff PERTH 05670721, 1♂ (AMNH\_PBI 00012351) *Rhagodia preissii preissii* Moq. (Chenopodiaceae), det. Perth staff PERTH-05670713, 6♂ (AMNH\_PBI 00015761–AMNH\_PBI 00015766), 4♀ (AMNH\_PBI 00015769–AMNH\_PBI 00015772) (AM). Pilbara Region: Hamersley Station, Railway Rd. just past Cooks Bore to Ridge paddock, 22.30166°S 117.69305°E, 30 Sep 2004–05 Oct 2004, C.V.M. volunteers, 2♂ (AMNH\_PBI 00123410, AMNH\_PBI 00123411), 3♀ (AMNH\_PBI 00123412–AMNH\_PBI 00123414) (AM). Poison Creek Beach, Cape Arid National Park, 33.90416°S 123.3519°E, 2 m, 25 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Atriplex isatidea* Moq. (Chenopodiaceae) PERTH05672317, 17♂ (AMNH\_PBI 00015648–AMNH\_PBI 00015664), 12♀ (AMNH\_PBI 00015665–AMNH\_PBI 00015676) *Atriplex isatidea* subsp. *tragus* Moq. (Chenopodiaceae) PERTH05672317, 17♂ (AMNH\_PBI 00015648–AMNH\_PBI 00015664), 12♀ (AMNH\_PBI 00015665–AMNH\_PBI 00015676) (AM). *Atriplex isatidea* Moq. (Chenopodiaceae), det. Perth staff PERTH05672317, 3♂ (AMNH\_PBI 00002732–AMNH\_PBI 00002734), 4♀ (AMNH\_PBI 00002705–AMNH\_PBI 00002708) (AMNH). Rossiter Bay, Cape Le Grande National Park, 33.96726°S 122.2674°E, 3 m, 23 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Atriplex isatidea* Moq. (Chenopodiaceae), det. Perth staff PERTH05672317, 1♂

(AMNH\_PBI 00002736), 2♀ (AMNH\_PBI 00002737, AMNH\_PBI 00002738) (AMNH). Stokes Inlet, Stokes National Park, 33.8179°S 121.1506°E, 2 m, 26 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Rhagodia baccata baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH05671922, 19♂ (AMNH\_PBI 00015543–AMNH\_PBI 00015561) (AM). *Rhagodia baccata baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH-05671922, 6♂ (AMNH\_PBI 00003111–AMNH\_PBI 00003116), 7♀ (AMNH\_PBI 00003117–AMNH\_PBI 00003123) (AMNH). Useless Loop Rd ca. 20 km W of jct with Shark Bay Rd, 26.56331°S 113.9338°E, 30 m, 25 Oct 2004, Cassis, Wall, Weirauch, Symonds, *Rhagodia latifolia* subsp. *latifolia* (Benth.) Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6987605, 3♂ (AMNH\_PBI 00015515–AMNH\_PBI 00015517), 8♀ (AMNH\_PBI 00015518–AMNH\_PBI 00015525) (AM). Yalgorup National Park, near Martins Tank Campground, 32.84222°S 115.66222°E, 15 m, 14 Dec 1997, Schuh, Cassis, Brailovsky, *Rhagodia* sp. (Chenopodiaceae), det. Perth staff PERTH05054990, 2♂ (AMNH\_PBI 00006125, AMNH\_PBI 00006126), 2♀ (AMNH\_PBI 00006127, AMNH\_PBI 00006128) (AM). *Rhagodia* sp. (Chenopodiaceae), det. Perth staff PERTH05054990, 2♂ (AMNH\_PBI 00002728, AMNH\_PBI 00002729), 1♀ (AMNH\_PBI 00002730) (AMNH).

**OTHER SPECIMENS EXAMINED: AUSTRALIA:** **Australian Capital Territory:** Weise's Orchard, Pialligo, 35.317°S 149.183°E, 18 Oct 1968, D. Morgan, *Malus* sp. (Rosaceae), 5♂ (AMNH\_PBI 00033216–AMNH\_PBI 00033220), 11♀ (AMNH\_PBI 00033221–AMNH\_PBI 00033231) (ANIC). **New South Wales:** 32.2 km SSW of Bourke on banks of Darling River, 35.3°S 147.9°E, 26 Dec 1973, G. F. Gross, Light Trap, 7♂ (AMNH\_PBI 00038546–AMNH\_PBI 00038551, AMNH\_PBI 00038576), 6♀ (AMNH\_PBI 00038552–AMNH\_PBI 00038556, AMNH\_PBI 00038577) (SAMA). 9.5 km E of Balranald on Sturt Hwy, 34.702°S 143.615°E, 20 Oct 1996, Schuh and Cassis, *Nitraria billardierei* DC. (Zygophyllaceae), det. Perth staff PERTH05095247, 6 juveniles (AMNH\_PBI 00006377–AMNH\_PBI 00006382) (AM). Cowra, 33.819°S 148.658°E, 25 Mar 1960, M. I. Nikitin, Light Trap, 1♂ (AMNH\_PBI 00192756), 1♀

(AMNH\_PBI 00192757) (BMNH). Leeton, 34.549°S 146.401°E, 27 Sep 1968, E. L. Jones, 4♀ (AMNH\_PBI 00038584, AMNH\_PBI 00038585) (SAMA). Murwillumbah, 28.33°S 153.383°E, 08 Apr 1919, F. Muir, 1♂ (AMNH\_PBI 00042182) (BPBM). The Rock Nature Reserve, 35.265°S 147.068°E, 14 Sep 1968, Key, Upton and Balderson, 1♀ (AMNH\_PBI 00033243) (ANIC). **Northern Territory:** 22 miles S of Alice Springs, 24.3°S 134.6°E, 15 Feb 1996, J. A. Grant, 3♂ (AMNH\_PBI 00192758–AMNH\_PBI 00192760), 2♀ (AMNH\_PBI 00192761, AMNH\_PBI 00192762) (BMNH). Alice Springs, 23.699°S 133.881°E, 16 Feb 1966, J. A. Grant, 16♂ (AMNH\_PBI 00015737–AMNH\_PBI 00015752), 8♀ (AMNH\_PBI 00015753–AMNH\_PBI 00015760) (AM); 17 Feb 1996, J. A. Grant, 3♂ (AMNH\_PBI 00192763–AMNH\_PBI 00192765) (BMNH); 14 Feb 1996, J. A. Grant, 1♀ (AMNH\_PBI 00192766) (BMNH). **Queensland:** 91 km N of Quilpie, 25.99847°S 144.4098°E, 300 m, 02 Nov 1998, Schuh, Cassis, Silveira, *Hakea leucoptera* R. Br. (Proteaceae), det. NSW staff NSW427661, 1 adult (AMNH\_PBI 00002703) (AMNH). Dalby, 27.17994°S 151.27176°E, F. H. Hobler, 2♂ (AMNH\_PBI 00038578, AMNH\_PBI 00038579) (SAMA). Sandgate, 27.317°S 153.064°E, 09 Nov 1919, F. Muir, 1♀ (AMNH\_PBI 00002746) (AMNH). **South Australia:** 17.6 km S of Innamincka, 27.88068°S 140.6712°E, 130 m, 06 Nov 1998, Schuh, Cassis, Silveira, *Chenopodium auricomum* Lindley (Chenopodiaceae), det. NSW staff, 3 juveniles (AMNH\_PBI 00006351–AMNH\_PBI 00006353) (AM). 5 km SW of Whyalla, 33.05085°S 137.5004°E, 30 m, 21 Oct 1996, Schuh and Cassis, *Exocarpos aphyllus* R.Br. (Santalaceae), det. Perth staff PERTH 05056209, 6♂ (AMNH\_PBI 00012352–AMNH\_PBI 00012357) (AM). Eyre Peninsula: 35 km N of Streaky Bay, 32.583°S 134.128°E, 20 m, 13 Aug 1978, W. C. Gagne, (Chenopodiaceae), 5♂ (AMNH\_PBI 00042173–AMNH\_PBI 00042177), 3♀ (AMNH\_PBI 00042178–AMNH\_PBI 00042180) (BPBM). Lyup, near Berri, 34.283°S 140.603°E, 28 Oct 1968, N. Richardson, *Prunus* sp. (Rosaceae), 8♂ (AMNH\_PBI 00033232–AMNH\_PBI 00033239), 1♀ (AMNH\_PBI 00033240) (ANIC). *Prunus* sp. (Rosaceae), 2♂ (AMNH\_PBI 00038574, AMNH\_PBI 00038575)

(SAMA). Mount Crawford State Forest, 34.7122°S 138.9469°E, 06 May 1968, D. Colless and Z. Liepa, 2♀ (AMNH\_PBI 00033241, AMNH\_PBI 00033242) (ANIC). Mount Loft, 34.974°S 138.709°E, unknown, 3♀ (AMNH\_PBI 00038580, AMNH\_PBI 00038581) (SAMA). Owieandana, Flinders Ranges, 30.446°S 138.948°E, Nov 1924, Hale & Tindale, 4♂ (AMNH\_PBI 00038570, AMNH\_PBI 00038573), 2♀ (AMNH\_PBI 00038571, AMNH\_PBI 00038572) (SAMA). near Victory Well, Everard Pk. Stn, 27.054°S 132.506°E, 03 Nov 1970, G. Gross, Light Trap, 1♂ (AMNH\_PBI 00038568), 1♀ (AMNH\_PBI 00038569) (SAMA); 29 Oct 1970, E.G. Matthews, Light Trap, 2♂ (AMNH\_PBI 00038563, AMNH\_PBI 00038564), 2♀ (AMNH\_PBI 00038565, AMNH\_PBI 00038566) (SAMA). **Victoria:** 8 km W of Kaniva, 36.377°S 141.241°E, 21 Oct 1963, J. Sedlacek, 1♀ (AMNH\_PBI 00042181) (BPBM). **Western Australia:** 11 mi. SW of Cocklebiddy, 32.45°S 126.03°E, 22 Nov 1969, Britton, Taylor and Upton, Light Trap, 1♂ (AMNH\_PBI 00038557), 6♀ (AMNH\_PBI 00038558–AMNH\_PBI 00038562, AMNH\_PBI 00038567) (SAMA). 13.2 km N of jct of Agana Kalbarri Rd and Brand Hiway, Galena River Bridge, 27.82917°S 114.6873°E, 190 m, 24 Oct 2004, Cassis, Wall, Weirauch, Symonds, *Atriplex amnicola* Paul G. Wilson (Chenopodiaceae), det. Perth staff PERTH6988431, 1 juvenile (AMNH\_PBI 00012350) (AM). 27.6 km N of Coral Bay Rd on Cardabia-Ningaloo Rd, 22.90198°S 113.8167°E, 25 m, 29 Oct 2004, Cassis, Wall, Weirauch, Tatarnic, Symonds, *Salsola tragus* subsp. *tragus* L. (Chenopodiaceae), det. Field ID, 6 juveniles (AMNH\_PBI 00015642–AMNH\_PBI 00015647) (AM). 43 km N of Norseman, 31.85648°S 121.6414°E, 300 m, 19 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Olearia muelleri* (Sond.) Benth. (Asteraceae), det. Perth staff PERTH05670691, 1 juvenile (AMNH\_PBI 00002612) (AMNH). Cape Arid National Park, Thomas River Campground, 33.8539°S 123.0126°E, 20 m, 24 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Atriplex isatidea* Moq. (Chenopodiaceae), det. Perth staff PERTH-05672317, 1 juvenile (AMNH\_PBI 00002743) (AMNH). Carnarvon, Small Boat Harbour, 24.90044°S 113.6477°E, 26 Oct 2004, Cassis, Wall, Weirauch, Tatarnic, Symonds, *Cheno-*

*podium murale* L. (Chenopodiaceae), det. Perth staff PERTH6989047, 1 juvenile (AMNH\_PBI 00012092) (AM). Conspicuous Beach, Walpole-Nornalup National Park, 10 km E of Nornalup, 35.03725°S 116.8443°E, 30 m, 17 Dec 1997, Schuh, Cassis, Brailovsky, *Rhagodia baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH05095271, 3 juveniles (AMNH\_PBI 00006168–AMNH\_PBI 00006170) (AM). Fowlers Camp (at end of Fowlers Camp Rd), Shark Bay World Heritage Area, 26.10801°S 113.6167°E, 24 Oct 2004, Cassis, Wall, Weirauch, Symonds, *Chenopodium murale* L. (Chenopodiaceae), det. Perth staff PERTH6988016, 1 juvenile (AMNH\_PBI 00015901) *Salsola tragus* subsp. *tragus* L. (Chenopodiaceae), det. Perth staff PERTH6988024, 1 juvenile (AMNH\_PBI 00012095) (AM). Mosman Park, Perth, 32.0209°S 115.7687°E, 20 m, 05 Nov 1998, G. Cassis, *Rhagodia baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH05227283, 1♀ (AMNH\_PBI 00015924) (AM). Peak Charles National Park Campground, 32.88335°S 121.1703°E, 300 m, 20 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Rhagodia preissii* Moq. (Chenopodiaceae), det. Perth staff PERTH05670713, 1 juvenile (AMNH\_PBI 00012423) (AM). Poison Creek Beach, Cape Arid National Park, 33.90416°S 123.3519°E, 2 m, 25 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Atriplex isatidea* subsp. *tragus* Moq. (Chenopodiaceae) PERTH-05672317, 15 juveniles (AMNH\_PBI 00015677–AMNH\_PBI 00015691) (AM). Rossiter Bay, Cape Le Grande National Park, 33.96726°S 122.2674°E, 3 m, 23 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Atriplex isatidea* Moq. (Chenopodiaceae), det. Perth staff PERTH-05672317, 1 juvenile (AMNH\_PBI 00002739) (AMNH). Stokes Inlet, Stokes National Park, 33.8179°S 121.1506°E, 2 m, 26 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Rhagodia baccata* (Labill.) Moq. (Chenopodiaceae), det. Perth staff PERTH-05671922, 5 juveniles (AMNH\_PBI 00015562–AMNH\_PBI 00015566) (AM). **MEXICO: Baja California Norte:** Bahia de los Angeles, Llave Island, site 5, 28.99869°N 113.52048°W, 20 m, 11 Mar 1998, F. Sanchez Pinero, 1♂ (AMNH\_PBI 00114164) (CNC). Bahia de los Angeles, Los Gemelos West Island, 28.95548°N 113.48144°W, 3 m, 06 Mar

1998, F. Sanchez Pinero, 54♂ (AMNH\_PBI 00114146–AMNH\_PBI 00114163, AMNH\_PBI 00114170–AMNH\_PBI 00114177, AMNH\_PBI 00114181–AMNH\_PBI 00114186, AMNH\_PBI 00114192–AMNH\_PBI 00114206, AMNH\_PBI 00114210–AMNH\_PBI 00114216), 18♀ (AMNH\_PBI 00114143–AMNH\_PBI 00114145, AMNH\_PBI 00114165–AMNH\_PBI 00114168, AMNH\_PBI 00114178–AMNH\_PBI 00114180, AMNH\_PBI 00114187–AMNH\_PBI 00114191, AMNH\_PBI 00114207–AMNH\_PBI 00114209), 1 juvenile (AMNH\_PBI 00114169) (CNC). **Chihuahua:** Ruinas de Paquime, SW of Nuevo Casas Grande, 30.36888°N 107.94777°W, 1492 m, 26 Aug 1998, M. D. Schwartz, *Chenopodium* sp. (Chenopodiaceae), 31♂ (AMNH\_PBI 00114082–AMNH\_PBI 00114085, AMNH\_PBI 00114088–AMNH\_PBI 00114114), 30♀ (AMNH\_PBI 00114086–AMNH\_PBI 00114087, AMNH\_PBI 00114115–AMNH\_PBI 00114142) (CNC). Sonora: 11 km E of Rosario de Tesopaco on road to La Estrella, 27.83247°N 109.2528°W, 479 m, 03 May 2005, Schwartz, Weirauch, Cervantes, 1♂ (AMNH\_PBI 00108489), 1♀ (AMNH\_PBI 00108490) (AMNH). **NEW ZEALAND:** Hawke's Bay: Napier, 39.5043°S 176.8924°E, 06 Apr 1975, W.J. Knight, 16♂ (AMNH\_PBI 00192767–AMNH\_PBI 00192782), 15♀ (AMNH\_PBI 00192783–AMNH\_PBI 00192797) (BMNH). **Marlborough:** Kaikoura Peninsula, 42.4355°S 173.7102°E, 26 Jan 2000–28 Jan 2000, D.K. McAlpine and R. Meier, 2♂ (AMNH\_PBI 00012439, AMNH\_PBI 00012440) (AM). North Auckland: Days Bay, 36.8128°S 175.1656°E, 01 Feb 1951, T. E. Woodward, *Lepidium oleraceum* Forst. (Cruciferae), 4♂ (AMNH\_PBI 00192748–AMNH\_PBI 00192751), 3♀ (AMNH\_PBI 00192752–AMNH\_PBI 00192754) (BMNH). **NORFOLK ISLAND:** none, 29.033°S 167.949°E, Mar 1971, K.L.S. Harley, 1♀ (AMNH\_PBI 00033251) (ANIC); A. M. Lea, 1♀ (AMNH\_PBI 00038582) (SAMA). **USA: Arizona:** *Cochise Co.*: Chiricahua Mountains, 1 mi N of Rustler Park, 31.91472°N 109.26861°W, 2536 m, 18 Aug 2000–21 Aug 2000, J. B. Woolley, 1♂ (AMNH\_PBI 00184915) (TAMU). Rt. 191, ca. 5 km N of Cochise, 32.16°N 109.94166°W, 20 Jun 2005, A. G. Wheeler, Jr., *Atriplex canescens* (Chenopodiaceae), 8♂ (AMNH\_PBI 00191166–AMNH\_PBI 00191173), 6♀ (AMNH\_PBI 00191174–

AMNH\_PBI 00191179) (USNM). ca. 12 km N of Benson, 32.05833°N 110.34166°W, 21 Jun 2005, A. G. Wheeler, Jr., *Atriplex canescens* (Chenopodiaceae), 1♂ (AMNH\_PBI 00191180), 2♀ (AMNH\_PBI 00191189, AMNH\_PBI 00191190) (USNM). *Navajo Co.:* Holbrook, 34.905°N 110.185°W, 19 Jun 2005, A. G. Wheeler, Jr., 1♀ (AMNH\_PBI 00191181) (USNM). **California:** *Los Angeles Co.:* Intercepted from Australia: Los Angeles, 33.94517°N 118.40266°W, 31 m, 06 Oct 1998, Unknown, *Delphinium* sp. (Ranunculaceae), 2♀ (AMNH\_PBI 00133761, AMNH\_PBI 00133762) (USNM). *Orange Co.:* Trampas Canyon at San Juan Creek, 33.51555°N 117.57888°W, 04 Sep 1999, Yanega and Gates, 1♀ (AMNH\_PBI 00119981) (UCR). *Riverside Co.:* Riverside, 33.95333°N 117.39528°W, 25 Mar 2003, K. W. Cooper, 1♀ (AMNH\_PBI 00133750) (USNM); 07 Aug 2001, K. W. Cooper, 1♀ (AMNH\_PBI 00133746) (USNM); 10 Aug 2002, K. W. Cooper, 1♀ (AMNH\_PBI 00133748) (USNM); 24 Aug 1999, K. W. Cooper, 1♀ (AMNH\_PBI 00133749) (USNM); 14 Oct 2002, K. W. Cooper, 1♂ (AMNH\_PBI 00133747) (USNM). *San Bernardino Co.:* Pinyon Hills, 34.39722°N 117.60944°W, 1447 m, 31 Jul 2003, G. R. Ballmer and D. Powell, 1♀ (AMNH\_PBI 00119982) (UCR). *San Diego Co.:* Intercepted from Mexico: San Diego, 32.68787°N 117.13418°W, 2 m, 11 May 2000, unknown, *Atriplex* sp. (Chenopodiaceae), 1♀ (AMNH\_PBI 00133754) (USNM); 13 Aug 1998, unknown, *Limonium sinuatum* (L.) P. Mill. (Plumbaginaceae), 1♂ (AMNH\_PBI 00133755), 1 juvenile (AMNH\_PBI 00133756) (USNM); 20 Mar 1998, unknown, *Limonium sinuatum* (L.) P. Mill. (Plumbaginaceae), 1♂ (AMNH\_PBI 00133757) (USNM); 19 Mar 1998, unknown, *Limonium sinuatum* (L.) P. Mill. (Plumbaginaceae), 1♀ (AMNH\_PBI 00133758) (USNM); 06 Nov 1996, unknown, *Celosia* sp. (Amaranthaceae), 1♂ (AMNH\_PBI 00133759) (USNM); 18 Sep 2001, unknown, *Amaranthus caudatus* (L.) (Amaranthaceae), 1♀ (AMNH\_PBI 00133760) (USNM). **Florida:** *Broward Co.:* Intercepted from Mexico: Fort Lauderdale [IPRL, USDA], 26.08128°N 80.23768°W, 3 m, 21 May 1997, unknown, *Atriplex* sp. (Chenopodiaceae), 3♀ (AMNH\_PBI 00133751–AMNH\_PBI 00133753) (USNM). **Hawaii:**

French Frigate Shoals: Tern Island, 23.87°N 166.28472°W, 25 Sep 2000, G.M. Nishida, 1♂ (AMNH\_PBI 00042602) *Chenopodium murale* (Chenopodiaceae), 2♂ (AMNH\_PBI 00042599, AMNH\_PBI 00042600), 1♀ (AMNH\_PBI 00042601) (BPBM). **New Mexico:** Otero Co.: Alamogordo, 32.88°N 105.96166°W, 22 Jun 2005, A. G. Wheeler, Jr., *Atriplex canescens* (Chenopodiaceae), 3♀ (AMNH\_PBI 00191182–AMNH\_PBI 00191184) (USNM). **Texas:** Brewster Co.: Big Bend National Park, Rio Grande nr Boquillas, 29.19583°N 102.93194°W, 576 m, 28 Apr 2001, D. E. Baumgardner, 4♀ (AMNH\_PBI 00184911–AMNH\_PBI 00184914) (TAMU). Cameron Co.: Jct. FM 509 & US 281, 26.052°N 97.73583°W, 19 Sep 2006, T. J. Henry and R. Coleman, 1♂ (AMNH\_PBI 00191185), 3♀ (AMNH\_PBI 00191186–AMNH\_PBI 00191188) (USNM).

*Coridromius chinensis* Liu and Zhao  
Figures 1C–D, 4, 6, 7N–P, 9A, map 6

*Coridromius chinensis* Liu and Zhao, 1999: 56–58.  
*Coridromius bufo* Miyamoto and Yasunaga, 1999: 33–35 (junior synonym); Chérot et al., 2004: 63–64 (synonymy).

**DIAGNOSIS:** Distinguished by the following combination of characters: bilobed proepisternum; swollen, rounded lobe at apex of scutellum; mottled coloration; AI with dark brown basal annulation; AII with dark brown annulation but lacking dark brown postmedial band; right paramere with sharp, triangular spicule; left paramere scyphelike with gutter open its entire length. *C. chinensis*, *confusus*, *minusculus*, and *punctatus* are all very similar but can be distinguished from one another by minor variations in color pattern and slight differences in male genitalia and scutellar shape. The apex of the scutellum, while swollen in all four species, is much more so in *chinensis* than in the others. *C. confusus* is also easily distinguished from the others, as it is the only species where the left paramere is extremely thin and forms a completely enclosed tube, open only at the apex. *C. minusculus* also has a very thin left paramere, but it is open the entire length. *C. confusus* and *minusculus* also share identical antennal banding, with a pale yellow band dividing the dark brown annulation on AII,

as well as similar hemelytral coloration, with the dark brown embolium marked with two broad yellow markings apically and basally. *C. chinensis* and *punctatus*, on the other hand, have thicker left parameres and lack a subapical dark brown annulation on AII. To separate *chinensis* from *punctatus* one can look at the coloration of AI: in the former it is pale with a dark brown band, while in the latter it is dark brown with yellow base and apex. Additionally, the ventral apical process on the posterior margin of the pygophore is long and thin in *chinensis* but shorter and thicker in *punctatus*.

**REDESCRIPTION: COLORATION** (figs. 1D, 4, 6): Head: Mottled light orange-brown and whitish yellow; frons with faint light orange-brown vittae, medial stripe pale; depressions on vertex and above antennae darkened; clypeus whitish yellow with two irregular dark brown stripes; mandibular and maxillary plates whitish yellow, becoming orange-brown along posterior margins; gena orange-brown; buccula whitish yellow, brown basally; labrum orange-brown, labium light orange-brown, darkened apically. Antenna: AI whitish yellow with dark brown annulation near base; AII light orange-brown with whitish-yellow subapical band followed by dark brown annulation; AIII and AIV dark brown with whitish-yellow bases. Thorax: Pronotal collar whitish-yellow; pronotum whitish yellow with mottled orange-brown punctures, with two small slightly darker orange-brown markings along anterior or behind eyes, seldom darkened around collar region, sometimes with indistinct whitish-yellow medial stripe, sometimes with two indistinct, broad dark brown markings on pronotal disc, margins without distinct piping; mesoscutum orange-brown; scutellum whitish yellow with mottled orange-brown punctures, medial stripe and apex pale; thoracic pleura dark orange-brown, whitish yellow along caudal margins; proepisternal lobes whitish yellow. Hemelytra: Mostly orange-brown, sometimes with anterior portion of clavus slightly to distinctly darker, commissure pale; corium becoming paler at apex, sometimes with faint darker brown patches; embolium with faint whitish-yellow subbasal and apical markings; cuneus light orange-brown, pale at apex; membrane




Fig. 9. Scanning electron micrographs: **A:** *C. chinensis* ♂ genitalia, posterior view. **B–H:** *C. confusus*: **B:** Head, anterior view. **C:** ♀ scutellum and abdomen, dorsal view. Arrow indicates minor swelling and desclerotization of laterotergite I. **D:** Detail of swelling of laterotergite I. **E:** ♀ thorax and abdomen, lateral view. **F:** Detail of metepimeron and peritreme. **G:** ♂ genitalia, dorsal view. **H:** Detail of ♂ genitalia. LP = left paramere, MES = mesepimeron, MET = metepimeron, MN = metanotum, P = peritreme, RP = right paramere, VAP = ventral apical process.

Sue Lindsay, Australian Museum

brown with whitish-yellow veins bounded by irregular darker brown markings. Legs: Mostly whitish yellow with brown markings; coxae whitish yellow; pro- and mesofemur with broad, subapical irregular brown markings, bothria whitish yellow; metafemur laterally with 6–7 irregular diagonal brown stripes basally, becoming blotchy and indistinct apically; irregular whitish-yellow patch anterodorsally; stripes fusing ventrally with large orange-brown patch, lateral depressions near apex orange-brown; all tibiae whitish yellow with irregular orange-brown markings; metatibial spines dark brown. Abdomen: Light orange-brown, posterior margin of sternite II dark orange-brown. SURFACE AND VESTITURE (figs. 4, 6): Irregular swellings on head with deep, somewhat coalescing punctures; pronotum and propleuron deeply and irregularly punctate; scutellum deeply and irregularly rugulopunctate, impunctate along midline; metepimeron and hemelytra impunctate; metafemur with a few punctures distodorsally; head and dorsum clothed in long, white setae. STRUCTURE (figs. 1D, 4, 6): Head: Highly contoured; frons strongly tumescent, bounded by irregular depressions and three minor swellings laterally and posteriorly; foveae adjacent to eyes and near posterior margin of vertex deep; posterior margin of head medially rounded, not carinate; gena swollen; mandibular plate tumescent. Thorax: Pronotum broad and somewhat elongate, posteriorly rounded, steeply declivit, deeply, irregularly punctate, anterolateral margins weakly carinate; submarginal region of humeral angles strongly excavate; posterior weakly medially cleft; collar region weakly developed; proepisternum bilobed; propleuron deeply irregularly punctate; posterior margin of metepimeron somewhat lobate; metanotum prominent and flared; scutellum dorsally flattened, apex a rounded lobe (figs. 1D, 4). Hemelytra: Costal margin evenly rounded over its entire length, anterior portion not expanded. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. MALE GENITALIA (figs. 7N–P, 9A): Right paramere rounded with a long, sharp triangular process (figs. 7O–P, 9A); left paramere scythe-

shaped, slightly twisted along its axis, thick and laterally compressed at base, becoming thin apically, gutter open over entire length (figs. 7N, 9A); margin of pygophore with long mesal suture and with long, thin fingerlike ventral apical process (fig. 9A). FEMALE PARAGENITALIA: No visible external paragenitalia.

HOST: Collected from the flowers of an unidentified *Macaranga* sp. (fig. 1C–D), *Mallotus japonicus* (Euphorbiaceae) and *Pentaspadon motleyi* (Anacardiaceae).

DISTRIBUTION: *C. chinensis* is one of the most widely distributed species of *Coridromius*, ranging from Nepal and India in the west to Brunei in the east (map 6). It is also known from Northern Queensland, Australia, where it is represented by two specimens.

PARATYPES: CHINA: Guangxi Prov.: Yangshu County, 24.7°N 110.4°E, 11 Sep 1964, Liang-Chen Wang, 3♂ (AMNH\_PBI 00178050–AMNH\_PBI 00178051, AMNH\_PBI 00718052), 1♀ (AMNH\_PBI 00718053) (NKMU).

OTHER SPECIMENS EXAMINED: AUSTRALIA: Queensland: 1 km SE of Mt. Cook, 15.3°S 145.16°E, 13 Oct 1980, T. Weir, 2♀ (AMNH\_PBI 00201293, AMNH\_PBI 00201294) (QM). BRUNEI DARUSSALAM: Bukit Sulang, near Lamunin, 4.67777°N 114.72694°E, 20 Aug 1982–10 Sep 1982, N. E. Stork, *Pentaspadon motleyi* (Anacardiaceae), 1♂ (AMNH\_PBI 00178011) (BMNH). CHINA: Yunnan Prov.: Xishuangbanna Tropical Botanical Garden, edge of gallery forest, 21.93222°N 101.245°E, 547 m, 26 May 2006, N. Tatarnic, *Macaranga* sp. (Euphorbiaceae), 13♂ (AMNH\_PBI 00006050–AMNH\_PBI 00006062), 5♀ (AMNH\_PBI 00006063–AMNH\_PBI 00006067) (AMNH). Xishuangbanna Tropical Botanical Garden, gallery forest by river, 21.93333°N 101.24444°E, 553 m, 26 May 2006–28 May 2006, N. Tatarnic, *Macaranga* sp. (Euphorbiaceae), 14♂ (AMNH\_PBI 00006020–AMNH\_PBI 00006033), 16♀ (AMNH\_PBI 00006034–AMNH\_PBI 00006049) (AMNH). INDIA: Karnataka: Kolar Dist.: Chikkaballapura (Chikballapur), 13.28°N 77.44°E, 1926, T.V. Campbell, 2♂ (AMNH\_PBI 00178005, AMNH\_PBI 00178006), 1♀ (AMNH\_PBI 00178007) (BMNH). Mysore, Nandidrug, 12.18°N 76.42°E, 1930, T. V. Campbell, 8 adults (AMNH\_PBI 00178001)

(BMNH); 1926, T. V. Campbell, 10 adults (AMNH\_PBI 00178003) (BMNH). **Meghalaya:** West Garo Hills, Balphakram NP, 25.18333°N 90.85°E, 250–550 m, 22–27 May 1996, E. Jendek and O. Sausa, 1♂ (AMNH\_PBI 00189982) (NHMW). West Garo Hills, Nokrek NP, 25.49333°N 90.325°E, 950–1250 m, 09–17 May 1996, E. Jendek and O. Sausa, 1♂ (AMNH\_PBI 00018971), 1♀ (AMNH\_PBI 00018972) (NHMW). **Uttaranchal:** Kumaon Div.: West Almora, 30.18333°N 80.01666°E, H. G. Crawford, 1♂ (AMNH\_PBI 00178004) (BMNH). Nainital Dist.: Bindukhera, 29.33333°N 79.5°E, 1938 m, 13 Apr 1909, Unknown, 1♂ (AMNH\_PBI 00178002) (BMNH). **JAPAN: Honshu:** Okayama Pref.: Mount Takakura, Sanyo T., Okayama, 34.75694°N 133.85527°E, 21 Jul 1996, T. Yasunaga, 2♂ (AMNH\_PBI 00190001), 5 adults (AMNH\_PBI 00189999, AMNH\_PBI 00190002) (TYCN). **Kyushu:** Miyazaki Pref.: Okawachi, Shiiba Village, 32.01666°N 131.35°E, 21 Jul 1992, T.J. Henry and A.G. Wheeler, 1♂ (AMNH\_PBI 00185358), 2♀ (AMNH\_PBI 00185356, AMNH\_PBI 00185357) *Mallotus japonicus* (Euphorbiaceae), 1♂ (AMNH\_PBI 00185359), 2♀ (AMNH\_PBI 00185360, AMNH\_PBI 00185361) (USNM). Okinawa Pref.: Ryukyu Islands, Ishigaki I., Kobo, Miyako, 24.2°N 124.09°E, 10–20 Nov 1952, G. Bohart, 1♂ (AMNH\_PBI 00042144) (BPBM). Ryukyu Islands, Miyako I., Hirara, 24.8°N 125.2833°E, 28 m, 23 Oct 1952, G. Bohart, 1♀ (AMNH\_PBI 00042143) (BPBM). **LAO PEOPLE'S DEMOCRATIC REPUBLIC:** Luang Namtha Prov.: Luang Nam Tha, ~3 km N of town, 20.97333°N 101.37777°E, 600 m, 03 Jun 2006, N. Tatarnic and D. C. Darling, *Macaranga* sp. (Euphorbiaceae), 5♂ (AMNH\_PBI 00006078–AMNH\_PBI 00006082), 1♀ (AMNH\_PBI 00006083) (AMNH). Luang Namtha, behind airport, 20.94777°N 101.40305°E, 547 m, 03 Jun 2006, N. Tatarnic and D. C. Darling, *Macaranga* sp. (Euphorbiaceae), 2♂ (AMNH\_PBI 00006075, AMNH\_PBI 00006076) (AMNH). Nam Tha N. P., ~30 min from Ban Nalan village, 20.84027°N 101.34694°E, 639 m, 04 Jun 2006, N. Tatarnic and D. C. Darling, *Macaranga* sp. (Euphorbiaceae), 1♂ (AMNH\_PBI 00006077) (AMNH). ca. 10 km E Luang Namtha, 20.95°N 101.4167°E, 600 m, 19 Jun

1996, Schillhammer, 1♂ (AMNH\_PBI 00190013) (NHMW). **Sekong Prov.:** Bolavens Plateau, N slope, ca. 10 km N. Mg Tha Theng, 15.5°N 106.43333°E, 500–700 m, 29–30 May 1996, Schillhammer, 2♂ (AMNH\_PBI 00189980, AMNH\_PBI 00189981) (NHMW). **NEPAL:** Royal Chitwan National Park, Machan Resort, 27.58305°N 84.5°E, 07–09 Sep 2005, T. Yasunaga, M. Takai and T. Shishido, 1♂, 1♀ (AMNH\_PBI 00190000) (TYCN). **TAIWAN:** Nantou Co.: Lien-hua-chih, 3 km W of Yuchih, 24.21667°N 121.5°E, 12–14 Jul 1992, T.J. Henry and A.G. Wheeler, 1♂ (AMNH\_PBI 00185363), 1♀ (AMNH\_PBI 00185364) *Mallotus japonicus* (Euphorbiaceae), 1♂ (AMNH\_PBI 00185362) (USNM). Pihu, 49 km E of Taipei on Hwy 9 to Yilan, 29.245°N 105.6458°E, 380 m, 18 Feb 1972, T.C. Maa, 1♀ (AMNH\_PBI 00042145) (BPBM). **VIETNAM:** Tam Dao NP, 21.4536°N 105.6436°E, 900 m, 17–18 Jun 1999, Y. Nakatani, 3♂ (AMNH\_PBI 00190009–AMNH\_PBI 00190011) (TYCN).

*Coridromius confusus*, new species  
Figures 4, 6, 7Q–S, 9B–H, map 3

**HOLOTYPE:** Male: **INDONESIA: Sulawesi Utara:** Dumoga-Bone National Park, Plot A, Fog 11, 0.613°N 124.089°E, 230 m, 10 Mar 1985, Royal Entomological Society of London: Project Wallace (AMNH\_PBI 00177949) (BMNH).

**DIAGNOSIS:** Distinguished by the following combination of characters: bilobed proepisternum; apex of scutellum rounded and weakly swollen; mottled coloration; AII with dark brown postmedial band; right paramere with sharp, triangular spicule; left paramere thin and scythelike with gutter closed until apex. This species is very similar to *chinensis*, *minusculus*, and *punctatus*, but the capillary-like left paramere, which is closed until its apex is diagnostic (for further discussion, see *chinensis* above).

**DESCRIPTION: COLORATION** (figs. 4, 6): Mottled cream and light brown. Head: Cream to whitish yellow with light brown markings, especially in punctures and foveae; frons with light brown vittae; mandibular and maxillary plates pale medially, becoming brown around edges; buccula pale, dark brown basally; clypeus pale with two sub-

medial longitudinal light brown to reddish brown stripes; labrum red; labium pale, brown apically. Antenna: AI cream with median dark brown annulation; AII light orange-brown with dark brown apical annulation, broken by a cream band near its base, tip of segment bright red; AIII and AIV dark brown with cream bases, tip of AIII bright red. Thorax: Pronotal collar cream to light yellow brown, darker along anterior and posterior margins, posterior margin somewhat reddened; pronotum cream to whitish yellow with light brown punctures and darker brown mottling, two small dark brown marks along anterior margin behind eyes; with or without pale medial stripe; collar region sometimes darkened, disc sometimes with two irregular broad, darkened patches, lateral and posterior margins without distinct pale piping; mesoscutum dark orange-brown; scutellum cream to whitish yellow with mottled light brown punctures; thoracic pleura dark reddish brown, posterior margins not pale; proepisternal lobes cream. Hemelytra: Mostly orange-brown with cream markings; clavus darkened basally; embolium orange-brown, dark brown medially with broad cream markings anteriorly and posteriorly; cuneus orange-brown, generally becoming reddened on outer margin and at apex; membrane light brown with mottled darker brown markings; veins pale, bounded by irregular darker brown markings. Legs: Mostly cream to pale yellow with brown markings; pro- and mesofemora with broad, subapical dark brown bands, bothria distinct and pale; metafemur laterally with eight irregular diagonal dark brown stripes ventrally blended into a broad dark brown marking, several punctures at apex tinged red, lateroapical depressions orange-brown, bothria pale; tibiae cream with irregular dark brown bands; metatibial spines dark brown. Abdomen: Light orange-brown with darker brown markings laterally along dorsal margin. SURFACE AND VESTITURE: (figs. 4, 6, 9B–F) Irregular swellings on head with deep somewhat coalescing punctures (fig. 9B); pronotum and propleuron deeply and irregularly punctate; scutellum deeply and irregularly rugulopunctate, impunctate along midline (fig. 9C–D); metepimeron (fig. 9E–F) and hemelytra impunctate; metafemur irreg-

ularly punctate; head and dorsum clothed in long, white, decumbent setae. STRUCTURE (figs. 4, 6, 9B–D): Head: Highly contoured; frons strongly tumescent medially, bounded by irregular depressions and three minor swellings laterally and posteriorly; foveae adjacent to eyes and near posterior margin of vertex deep; posterior margin of head medially rounded, not carinate; gena swollen; mandibular plate tumescent (figs. 6, 9B). Thorax: Pronotum broad and somewhat elongate, posteriorly rounded, steeply declivous, anterolateral margin weakly carinate, posterior margins of humeral angles strongly excavate; posterior margin not medially cleft, collar region weakly developed; proepisternum bilobed; posterior margin of metepimeron in the form of a rounded, somewhat triangular lobe; metanotum prominent and flared (fig. 9C–D); scutellum somewhat flattened, apex slightly swollen and rounded (fig. 9C). Hemelytra: Costal margin evenly rounded over entire length. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. MALE GENITALIA (figs. 7Q–S, 9G–H): Right paramere rounded with a long, sharp, triangular apophysis (fig. 7R–S); left paramere scythe-shaped, slightly twisted along axis, thick and laterally compressed at base, becoming very thin, gutter completely closed until apex (figs. 7Q, 9G–H); posterior margin of pygophore with long mesal suture and thin fingerlike ventral apical process (fig. 9G–H). FEMALE PARAGENITALIA: Indistinct; first two right laterotergites dorsally membranous and transversely striate (fig. 9C–D), site of copulation unknown.

ETYMOLOGY: The epithet *confusus* (Latin, “indistinct, hard to recognize”) reflects the difficulty in separating this species from *chinensis* and other related species.

REMARKS: This species was collected in large numbers through canopy fogging by the Royal Entomological Society of London’s Project Wallace expedition in 1985.

HOST: No host records.

DISTRIBUTION: Known only from northern Sulawesi and a single specimen from the Solomon Islands (map 3).

PARATYPES: INDONESIA: Sulawesi Utara: Dumoga-Bone National Park, Plot

A, Fog 13, 0.613°N 124.089°E, 230 m, 11 Jul 1985, Royal Entomological Society of London: Project Wallace, 3♂ (AMNH\_PBI 00192728–AMNH\_PBI 00192730), 5♀ (AMNH\_PBI 00178040–AMNH\_PBI 00178042, AMNH\_PBI 00192737–AMNH\_PBI 00192738) (AM). 2♂ (AMNH\_PBI 00192726, AMNH\_PBI 00192727), 2♀ (AMNH\_PBI 00192735, AMNH\_PBI 00192736) (AMNH). 33♂ (AMNH\_PBI 00177976–AMNH\_PBI 00178000, AMNH\_PBI 00178036–AMNH\_PBI 00178039, AMNH\_PBI 00178044–AMNH\_PBI 00178047), 29♀ (AMNH\_PBI 00178012–AMNH\_PBI 00178035, AMNH\_PBI 00178043, AMNH\_PBI 00192731–AMNH\_PBI 00192734) (BMNH). 2♂ (AMNH\_PBI 00042586, AMNH\_PBI 00042587), 3♀ (AMNH\_PBI 00042588–AMNH\_PBI 00042590) (BPBM). Dumoga-Bone National Park, Plot A, Fog11, 0.613°N 124.089°E, 230 m, 10 Mar 1985, Royal Entomological Society of London: Project Wallace, 1♀ (AMNH\_PBI 00177963) (AM). 11♂ (AMNH\_PBI 00177950–AMNH\_PBI 00177960), 14♀ (AMNH\_PBI 00177961–AMNH\_PBI 00177962, AMNH\_PBI 00177964–AMNH\_PBI 00177975) (BMNH). Dumoga-Bone National Park, Sites 10 & 11, Tumpah Transect, 0.613°N 124.089°E, 1040 m, Feb 1985, J. D. Holloway, Light Trap, 1♀ (AMNH\_PBI 00192739) (BMNH).

OTHER SPECIMENS EXAMINED: SOLOMON ISLANDS: Guadalcanal: Honiara, 9.42805°S 160.05472°E, 200 m, Jan 1973, N. L. H. Krauss, 1♂ (AMNH\_PBI 00042142) (BPBM).

*Coridromius crassus*, new species  
Figures 4, 6, 10A–D, map 3

HOLOTYPE: Male: INDONESIA: Sulawesi Utara: Dumoga-Bone National Park, Gungung Ambang Forest Reserve, near Kotamobagu, 0.55°N 123.98333°E, 1200 m, 18 Feb 1985, Royal Entomological Society of London: Project Wallace (AMNH\_PBI 00192740) (BMNH).

DIAGNOSIS: Distinguished by the following characters: broadly flared hemelytra; bilobed proepisternum; thick, sharply angled left paramere; right paramere U-shaped when viewed from above; males with patch of dense conelike setae on the left posterior margins of abdominal segments VII and VIII. *C. crassus* is similar to *declivipennis*,

*nakatanii*, and *pteraulos* but is much wider across the pronotum and hemelytra, and differs in coloration. The left paramere is also much thicker than in the other three species.

DESCRIPTION: COLORATION (figs. 4, 6): Mostly orange-brown and yellow with darker brown markings. Head: Orange-brown, becoming more yellow on frons; vittae on frons dark brown; lateral and medial swellings on vertex yellow, merging with yellow ocular and posterior margins; clypeus yellow with one medial brown spot and two lateral brown stripes; gena, mandibular and maxillary plates yellow to orange-brown; buccula yellow, dark brown basally; labrum brown, labium yellow, brown apically. Antenna: AI dark brown, apices yellow; AII light orange-brown with pale subapical annulation adjacent to dark brown apical annulation; AIII and AIV dark brown, yellow basally. Thorax: Pronotal collar yellow; pronotum mostly orange-brown with faint yellow medial stripe and margins, two dark brown spots on anterior margin behind eyes, two larger brown spots above collar region, sometimes fused together, anterior of humeral angles sometimes with faint dark brown spots, submarginal region of humeral angles sometimes yellowed; mesoscutum orange-brown, dark brown medially; scutellum orange-brown with medial stripe, apex and sides yellow; thoracic pleura orange-brown to dark brown, posterior margins yellowed; anterior lobe of proepisternum yellow. Hemelytra: Orange-brown with dark brown marking halfway down (except on clavus), slightly yellow along claval commissure and lateral margins; fascia above cuneus yellow; cuneus orange-brown; membrane light brown with dark brown veins. Abdomen: Orange-brown with dark brown and yellow markings along dorsolateral margins. Legs: All coxae pale yellow; pro- and mesofemora pale yellow, brown apically, metafemur yellow to orange-brown with 9 diagonal dark brown stripes on outer surface, lateroapical depression dark brown; tibiae yellow, metatibia sometimes brown. SURFACE AND VESTITURE (figs. 4, 6): Minor swelling on vertex finely punctate; pronotum, propleuron, metepimeron, and scutellum finely punctate; hemelytra and cuneus irregularly punctate; head and dorsum clothed in long, white, decum-




Fig. 10. Left and right parameres of *Coridromius* spp.: A–D: *C. crassus*. E–H: *C. declivipennis*. I–K: *C. drepanopenis*.

bent setae; male abdomen with patch of dense, conelike setae on the left posteroventral margins of segments VII and VIII (similar to *pteraulos*, figs. 16E–F). STRUCTURE (figs. 4, 6): Head: Frons roundly tumescent medially, merging with minor swelling on vertex; vertex with two minor swellings adjacent to eyes; posterior margin of head rounded, somewhat carinate. Thorax: Pronotum broad and steep, posteriorly rounded, lateral margins carinate, anterolateral margins depressed, not upturned, submarginal region of humeral angles excavate, posterior margin steeply declivous, collar region weakly present; proepisternum bilobed; posterior margin of metepimeron in the form of a broad, rounded lobe; metanotum prominent and flared; scutellum somewhat flattened. Hemelytra: Costal margin greatly flared medially, with raised L-shaped fascia at apex. Legs: Metatibial spines long and thick. Abdomen: Postero-lateral margin of abdominal sternite II sharply angular. MALE GENITALIA (figs. 10A–D): Right paramere broad and rounded with thumblike lobe, appears U-shaped when viewed dorsally (figs. 10B–D); left paramere thick and elongate, sharply angled basally, then evenly curved and tapering to apex with a slight twist along its axis, curved at tip (fig. 10A); pygophore with long mesal suture and long fingerlike ventral apical process. FEMALE PARAGENITALIA: Abdominal tergite II asymmetrical, right lateral margin weakly tumescent, posterior margin sulcate.

ETYMOLOGY: The name *crassus*, meaning “thick” or “stout” in Latin, reflects the broad body of this insect.

HOST: No host records.

DISTRIBUTION: Known from northern Sulawesi (map 3).

PARATYPES: INDONESIA: Sulawesi Utara: Dumoga-Bone National Park, Danau Mooat, 0.75694°N 124.4625°E, 1300 m, May 1985, Royal Entomological Society of London: Project Wallace, 1♂ (AMNH\_PBI 00177945) (BMNH). Dumoga-Bone National Park, Gunung Ambang Forest Reserve, near Kotamobagu, 0.55°N 123.98333°E, 1200 m, 18 Feb 1985, Royal Entomological Society of London: Project Wallace, 3♂ (AMNH\_PBI 00177946, AMNH\_PBI 00192742–AMNH\_

PBI 00192743), 2♀ (AMNH\_PBI 00177947, AMNH\_PBI 00177948) (BMNH).

*Coridromius declivipennis* Miyamoto and Yasunaga

Figures 4, 6, 10E–H, 11A, map 6

*Coridromius declivipennis* Miyamoto and Yasunaga, 1999: 35–37 (n. sp.).

DIAGNOSIS: Recognized by the following combination of characters: bilobed proepisternum; rounded and cleft right paramere; long left paramere, sharply curved at base; embolium angularly flared near cuneus. Similar to *crassus*, *nakatanii*, and *pteraulos* but with shorter ventral apical process on the pygophore than in *crassus* and *pteraulos*, and different hemelytral shape, and coloration to all three.

REDESCRIPTION: COLORATION (figs. 4, 6): Mostly light brown with orange and brown markings. Head: Yellow-brown to orange-brown; vittae absent; ocular margins and posterior margin of head yellow; three dark brown spots, one immediately above clypeus and one over each antenna; clypeus, gena, mandibular and maxillary plates slightly paler; buccula dark brown basally then becoming yellow; labrum orange-brown, labium yellow, brown apically. Antenna: AI dark brown, base and apex yellowed; AII light orange-brown with pale subapical annulation and dark brown apical annulation; AIII and AIV dark brown, yellow basally. Thorax: Pronotal collar yellow; pronotum mostly orange-brown with faint yellow medial stripe, yellow margins and faint yellowing at humeral angles; mesoscutum light orange-brown; scutellum orange-brown to dark brown, with faint yellow apex and sides; thoracic pleura mostly orange-brown, posterior margins yellowed; lobes of proepisternum yellow, ventral portion of posterior lobe dark brown. Hemelytra: Yellow-brown to dark brown, darker anteriorly, yellow posteriorly; clavus yellow at commissure; corium and embolium yellow from apex of claval commissure to costal fracture; cuneus yellow-brown to orange-brown, sometimes darker apically; membrane brown with dark brown veins. Abdomen: Mostly yellow-brown to orange-brown, sometimes with dark brown markings around spiracles. Legs: All coxae

pale yellow; pro- and mesofemora yellow, brown apically; metafemur yellow basally, becoming brown apically, with diagonal brown striping on outer surface, sometimes dark brown immediately below striping; tibia yellow, sometimes slightly brown basally. SURFACE AND VESTITURE (figs. 4, 6): Head weakly punctate laterally and medially, densely setose; pronotum, propleuron, metepimeron, scutellum, and hemelytra finely punctate, clothed in long, simple, white, decumbent setae; male with short, dense and thick conelike setae on the left posteroventral margins of abdominal segments VII and VIII. STRUCTURE (figs. 4, 6): Head: Frons tumescent medially, merging with punctate minor swelling on vertex; vertex with two minor swellings adjacent to eyes, shallow depressions behind each swelling; posterior margin of head rounded, weakly carinate. Thorax: Pronotum broad and steep, posteriorly rounded, finely punctate; margins carinate, anterolateral margins somewhat depressed, not upturned, submarginal region of humeral angles excavate; collar region weakly defined; proepisternum bilobed; posterior margin of metepimeron truncate, upper corner slightly extended and rounded; metanotum prominent and flared; scutellum somewhat flattened. Hemelytra: Costal margin carinate, distally expanded. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterolateral margin of segment II angular. MALE GENITALIA: (figs. 10E–H, 11A) Right paramere broad and rounded with thumblike lobe, paramere appears U-shaped when viewed dorsally; left paramere thick and elongate, sharply angled basally, then evenly curved and tapering to apex with a slight twist along its axis, curved at tip; pygophore with a long mesal suture; ventral apical process broad and short (figs. 10, 11A). FEMALE PARAGENITALIA: Abdominal tergite II asymmetrical, right lateral margin weakly tumescent, posterior margin sulcate.

**REMARKS:** Based on the modifications of the female dorsum, it is thought that males inseminate females dorsolaterally.

**HOST:** No host records.

**DISTRIBUTION:** Known from Kyushu, Japan (map 6).

**PARATYPES: JAPAN: Kyushu:** Okinawa Pref.: Iriomote Is. Funaura, 24.3333°N

123.81666°E, 10 May 1993, T. Yasunaga, 3♀ (AMNH\_PBI 00189984) (ULB). Ishigaki Is. Mount Banna, 24.36666°N 124.16666°E, 08 May 1993, T. Yasunaga, 3♂ (AMNH\_PBI 00189983) (ULB). 2♀ (AMNH\_PBI 00185366) (USNM). 1♂, 1♀ (AMNH\_PBI 00189996) (AM), 1♂, 3♀ (AMNH\_PBI 00189997) (TYCN).

***Coridromius drepanopenis*, new species**  
Figures 4, 6, 10I–K, map 5

**HOLOTYPE:** Male: **MALAYSIA: Sabah:** Tenompok, 6.01777°N 116.48555°E, 637 m, 10–19 Feb 1959, T.C. Maa (AMNH\_PBI 00041548) (BPBM type# 16726) (BPBM).

**DIAGNOSIS:** Recognized by its single proepisternal lobe, orange-brown coloration, punctate pronotum and extremely long, thin left paramere. The genitalia of *drepanopenis* are very similar to those of *prolixipenis*, but the latter is much darker in coloration and lacks punctures on the pronotum.

**DESCRIPTION: COLORATION** (figs. 4, 6): Head: Mostly light orange-brown, frons with faint, darker brown vittae; raised tubercles adjacent to eyes yellow; ocular margins and posterior margin of vertex with yellow piping; clypeus, mandibular and maxillary plates, and gena light orange-brown; buccula yellow-orange; labrum and labium orange-brown, darker brown apically. Antenna: AII pale orange-brown, with thin dark brown annulations at base and apex; AIII light orange-brown with dark brown apical annulation; AIV dark brown, yellow basally. Thorax: Pronotal collar yellow, anterior margin dark brown; pronotum mostly orange-brown with faint yellow median stripe; margins with thin yellow piping; proepisternum yellow, remaining thoracic pleura orange-brown, posterior margins becoming yellow; mesoscutum orange-brown; scutellum orange-brown with faint dark brown markings on lateral margins and yellow markings laterally and apically. Hemelytra: Orange-brown, slightly lighter anteriorly and at cuneal fracture; clavus slightly darker than remainder of hemelytron; cuneus slightly yellow basally, becoming orange along outer and apical margins; membrane pale brown with darker brown veins. Abdomen: Uniformly yellow-orange; parameres glossy amber. Legs: Yellow-orange,




Fig. 11. Scanning electron micrographs: **A.** *C. declivipennis* ♂ genitalia, posterior view. **B–F:** *C. ephippius*: **B.** ♀ scutellum and abdomen, with hemelytra removed to expose paragenital swelling, dorsal view. **C.** ♀ thorax and abdomen, right lateral view. **D.** Detail of ♀ paragenital structure. **E.** ♂ genitalia, ventrolateral view. **F.** Detail of left paramere, dorsal view. **G–H:** *C. epithema*: **G.** ♀ thorax and abdomen, right lateral view. **H.** ♀ paragenital swelling of right laterotergites I and II, dorsal view. LP = left paramere, MES = mesepimeron, MET = metepimeron, MLS = mesal longitudinal suture, MN = metanotum, P = peritreme, RP = right paramere, VAP = ventral apical process.

Sue Lindsay, Australian Museum

metafemur faintly orange at apex, laterally with diagonal brown stripes. SURFACE AND VESTITURE (figs. 4, 6): Head smooth and glossy with widely spaced punctuation medially on vertex, sparsely covered with short, white setae; pronotum with shallow, even punctuation; scutellum slightly rugulose and impunctate; propleuron, metepimeron, and hemelytra impunctate; head and dorsum with sparse distribution of short, white, decumbent setae. STRUCTURE (figs. 4, 6): Head: Broad; frons weakly convex; vertex with minor swellings medially and adjacent to eyes; posterior margin of head carinate and weakly medially rounded. Thorax: Pronotum broad and convex, margins thinly carinate, anterolateral margins slightly upturned, submarginal region of humeral angles depressed, posterior margin thin, callar region undifferentiated; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared; scutellum swollen. Hemelytra: Costal margin thin, somewhat flared over its entire length and weakly sinuate. Legs: Metatibial spines short and thin. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not angular. MALE GENITALIA (fig. 10I–K): Right paramere small and triangular (fig. 10J); left paramere extremely long, thin, curved, slightly twisted along its axis (fig. 10I), and extending beyond the margin of the pygophore (fig. 10K); posterior margin of pygophore biconvex, with fold on left forming a shallow groove (fig. 10K). FEMALE PARAGENITALIA: Unknown.

ETYMOLOGY: This species name, *drepano-penis*, is derived from Greek *drepanon* (“scimitar, sword”) and Latin *penis* (lit., “tail”) and reflects its extremely long, sword-like left paramere.

HOST: No host information.


DISTRIBUTION: Known only from Sabah (map 5).

*Coridromius ephippius*, new species  
Figures 4, 6, 11B–F, 12A–C, map 3


HOLOTYPE: Male: PHILIPPINES: Albay Province: Mt. Mayon, 16 km NW of Legaspi, 13.2566°N 123.685°E, 1200–1800 m, 12 May 1962, H.M. Torrevillas (AMNH\_PBI 00041446) (BPBM type# 16727) (BPBM).

DIAGNOSIS: Readily distinguished by the female's unique paragenital structure (fig. 11B–D).


DESCRIPTION: COLORATION (figs. 4, 6): Head: glossy, mostly concolorous light brownish yellow, frons with faint brown vittae; with paired dark brown chevron-shaped markings on vertex behind weakly raised tumescences, sometimes also with median brown marking; labium pale, brown apically. Antenna: A1 brownish yellow with dark brown annulations at base and apex; AII brownish yellow with dark brown annulation; AIII dark brown, pale at base; AIV dark brown. Thorax: Pronotal collar yellow, brown along anterior margin; pronotum brownish yellow to brownish orange, sometimes with two irregular dark brown markings immediately behind eyes, posterior margin also sometimes darkened; mesoscutum brown to brownish orange; scutellum brownish yellow to brownish orange with sometimes fused darker brown medial and lateral markings, anterior angles and lateroapical margins yellow to brownish orange; thoracic pleura brownish yellow. Hemelytra: Brownish yellow in most specimens, in others clavus sometimes brown apically, joining faint brown arc at apex of endocorium; membrane faintly brown with darker brown veins. Legs: Mostly uniform brownish yellow, metafemur mostly pale with eight or nine diagonal brown stripes, metatibia pale. Abdomen: Brownish yellow. SURFACE AND VESTITURE (figs. 4, 6): Head finely and shallowly punctate on vertex and adjacent to eyes; pronotum shallowly but distinctly punctate; propleuron, metepimeron, scutellum, and hemelytra impunctate; head and dorsum clothed with sparse distribution of short, white, decumbent setae; laterally projecting tumescence on right abdominal sternite II of female nodulose and setaceous. STRUCTURE (figs. 4, 6): Head: Frons broadly tumescent medially; with paired slightly raised tumescences adjacent to eyes, foveae behind tumescences shallow; posterior margin of vertex almost flat, not carinate. Thorax: Pronotum broad, margins somewhat carinate, anterolateral margins slightly upturned, not explanate; submarginal region of humeral angles weakly excavate; callar region undifferentiated; proepisternum unilobed;



Map 5. Distribution map of *C. drepanopenis*, *falsicoleus*, *nakatanii*, *pteraulos*, *sommelierii*, *testaceous*, *thalame*, and *zetteli*.



Map 6. Distribution map of *C. chinensis* and *declivipennis*. (Not shown: *C. chinensis* is also known from two specimens collected in northern Queensland, Australia.)



Map 7. Distribution map of *C. carinatus*, *confusus*, *crassus*, *ephippius*, and *marmoreus*.

posterior margin of metepimeron truncate; metanotum not prominent and flared. Hemelytra: Costal margin thin, flared over its entire length and straight to weakly sinuate. Legs: Metatibial spines thin and small. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not angular. MALE GENITALIA (figs. 11E–F, 12A–C): Right paramere small and triangular (fig. 12B–C); left paramere thick and sharply angled at base, becoming straighter and thinner near apex (figs. 11E–F, 12A); ventral margin of pygophore biconvex, with distinct U-shaped groove on left side, no mesal suture or ventral apical process. FEMALE PARAGENITALIA (fig. 11B–D): Right laterotergites II and III fused with right sternites II and III to form a single dorsolateral swelling with laterally projecting tumescence (fig.

11B–D), posterior margin of sternite II flared and carinate (fig. 11C–D).


ETYMOLOGY: From the Latin *ephippium*, meaning “pack saddle”, this name refers to the saddlelike shape of the female paragenital structure.

REMARKS: The paragenitalia of *ephippius* immediately distinguish it from all other species. As with so many other species, the exact function of such elaborate external modifications remains to be discovered.

HOST: No host records.

DISTRIBUTION: Philippines, Albay Province (map 3).

PARATYPES: **PHILIPPINES: Albay Province:** Mt. Mayon, 16 km NW of Lagaspi, 13.2566°N 123.685°E, 1200–1800 m, 12 May 1962, H.M. Torrevillas, 2♂ (AMNH\_PBI 00041447, AMNH\_PBI 00041448), 1♀



Map 8. Distribution map of *C. epithema*, *hermosus*, *minusculus*, *neoguineanus*, *nigrus*, *prolixipenis*, and *punctatus*.

(AMNH\_PBI 00041449) (BPBM), 1♀ (AMNH\_PBI 00041555) (AM); 15 May 1962, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041556) (BPBM). Mt. Mayon, 16 km NW of Lagaspi, 13.2566°N 123.685°E, 900–1800 m, 05 May 1962, H. M. Torrevillas, 6♂ (AMNH\_PBI 00041557–AMNH\_PBI 00041562), 2♀ (AMNH\_PBI 00041563, AMNH\_PBI 00041564) (BPBM). Mt. Mayon, 16 km NW of Lagaspi, 13.2566°N 123.685°E, 1900–2000 m, 13 May 1962, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041565) (BPBM).

***Coridromius epithema*, new species**  
Figures 4, 6, 11G–H, map 8

**HOLOTYPE:** Female: **PAPUA NEW GUINEA: Morobe:** Mount Kaindi, 7.35°S 146.68333°E, 2350 m, 11–18 Sep 1972, J.L. Gressitt (AMNH\_PBI 00041549) (BPBM type# 16728) (BPBM).

**DIAGNOSIS:** Recognized by the following combination of characters: costal margin of hemelytron flared anteriorly, then sharply angled inward from the midpoint to the cuneus, extremely long metepimeral lobe, swollen and depressed right laterotergites II and III, and nearly uniform orange-brown coloration. This species appears very similar to *nigrus* and *hermosus*, both of which share similar paragenital structures, head shape,

hemelytral shape, and metepimeral lobes. All three can be distinguished by their coloration, and *ephippius* differs further by having the longest metepimeral lobe, which is rounded apically (fig. 11G) but shorter and triangular (fig. 13E) in the other species.

**DESCRIPTION: COLORATION** (figs. 4, 6): Nearly uniform orange-brown. Head: Orange-brown; mandibular and maxillary plates, buccula, gena, ocular margins, and posterior margin of vertex slightly lighter orange; labium yellow-orange, brown apically. Antenna: AI and AII orange brown, AIII and AIV missing from specimen. Thorax: Pronotal collar light orange-brown; pronotum orange-brown with paler medial stripe and margins; mesoscutum and scutellum orange-brown; thoracic pleura mainly dark orange-brown with mesopleuron and posterior margins somewhat paler. Hemelytra: orange-brown, membrane somewhat lighter. Abdomen: Dark orange-brown. Legs: Coxae yellow-orange; femora and tibiae orange-brown; metafemur without contrasting dark stripes; tarsi somewhat yellowed. **SURFACE AND VESTITURE** (figs. 4, 6): Impunctate; scutellum weakly rugulose; head and dorsum sparsely coated with white, decumbent setae, most setae knocked off specimen. **STRUCTURE** (figs. 4, 6, 11G): Head: Wide; weakly

tumescent medially; lateral tubercles adjacent to eyes small and ill-defined, bordered posteriorly by shallow depressions; posterior margin of vertex nearly flat. Thorax: Pronotum broad and steep, anterolateral margins weakly upturned, posterior and lateral margins carinate, submarginal region of humeral angles shallowly excavated, collar region undifferentiated; proepisternum unilobed; metepimeron extended in a long, winglike lobe (fig. 11G); metanotum not prominent and flared. Hemelytra: Embolium thin, flaring outward from base then abruptly angling medially at midpoint. Legs: Metatibial spines short and thin. MALE GENITALIA: Unknown. FEMALE PARAGENITALIA (fig. 11G–H): Right laterotergites II and III swollen, desclerotized and depressed, posterior margin of right abdominal sternite II strongly carinate and flared outward, small paragenital opening between sternites II and III (fig. 11G–H).

**ETYMOLOGY:** The name *epithema*, meaning “lid” or “cover” in Greek, reflects the extended metepimeral lobe which partially covers the paragenitalia.

**HOST:** Unknown.

**DISTRIBUTION:** Mt. Kaindi, Papua New Guinea (map 8).

#### *Coridromius falsicoleus*, new species

Figures 4, 6, 13A–B, map 5

**HOLOTYPE:** Female: **MALAYSIA: Sabah:** Tenompok, 6.01777°N 116.48555°E, 637 m, 13 Feb 1959, T.C. Maa (AMNH\_PBI 00041551) (BPBM type# 16729) (BPBM).

**DIAGNOSIS:** Recognized by the following: unique dark brown and yellow-orange color pattern; females with prominent paragenital opening on the right side of the abdomen within sternite II. The male of *falsicoleus* is unknown. *C. falsicoleus*, *nakatanii* and *sommelieri* are the only species with a prominent paragenital opening within the second abdominal segment, which in *falsicoleus* and *sommelieri* extends into the abdomen as a sclerotized copulatory tube. In the latter however, this copulatory tube is corkscrewed while in *falsicoleus* it is not.

**DESCRIPTION: COLORATION** (figs. 4, 6): Mostly dark brown with yellow markings. Head: Mostly dark brown, vittae faintly

present on frons; mandibular plates yellow; maxillary plates and gena dark brown; buccula brown basally, pale along ventral margin; labrum light brown; labium light brown, dark brown apically. Antenna: AI dark brown, lighter brown basally; AII dark brown with two yellow bands, one broad band subbasally, and another, thinner band subapically; AIII and AIV dark brown, yellow basally. Thorax: Pronotal collar yellow; pronotum dark brown with median yellow stripe, posterior of humeral angles yellowed; thoracic pleura dark brown, posterior margins yellowed; mesoscutum dark brown; scutellum dark brown with three yellow spots laterally and apically. Hemelytra: Dark brown, claval commissure yellowed, corium with faint yellow U-shaped pattern. Legs: Pro- and mesofemur yellow, becoming dark brown at apex; metafemur brown, yellow basally, with 8–9 diagonal yellow stripes on outer surface; pro- and mesotibia with basal half dark brown, apical half yellow; metatibia dark brown; tarsi yellow. Abdomen: Chestnut brown, upper margins of each sternite dark brown, bordered posteriorly by yellow. SURFACE AND VESTITURE (figs. 4, 6, 13A): Head, pronotum, propleuron, metepimeron (fig. 20A), scutellum, hemelytra, and cuneus finely punctate; head and dorsum clothed in long, white, decumbent setae. STRUCTURE (figs. 4, 6, 13A): Head: Frons weakly tumescent medially, swelling of frons merging with minor medial swelling on vertex; posterior margin of vertex carinate and rounded. Thorax: Pronotum steep, posteriorly rounded, anterolateral margins not upturned, submarginal region of humeral angles strongly excavated, collar region undifferentiated; proepisternum bilobed; posterior margin of metepimeron extended as a short, broad, rounded lobe (fig. 13A); metanotum prominent and flared; scutellum flattened. Hemelytra: Embolium carinate and flared outward posteriorly. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of sternite II sharply angular. MALE GENITALIA: Unknown. FEMALE PARAGENITALIA (fig. 13A–B): Right abdominal sternite II with prominent paragenital opening, partly obscured by metepimeral lobe (fig. 13A–B), forms a thinly




Fig. 12. Left and right parameres of *Coridromius* spp.: A–C: *C. ephippius* (arrow indicates aedeagus). D–F: *C. hermosus*. G–I: *C. lestoni*. J–L: *C. marmoreus*. M–O: *C. minusculus*.

Nikolai Tatarnic

sclerotized invagination into the abdominal cavity.

**ETYMOLOGY:** Derived from Latin (*falsi-*, “false”, and *coleus*, “sheath” and thus “vagina”), the species name *falsicoleus* refers to this species’ prominent paragenital opening.

**HOST:** Unknown.

**DISTRIBUTION:** *C. falsicoleus* has only been collected in Sabah (map 5).

**PARATYPES: MALAYSIA: Sabah:** Tenompok, 6.01777°N 116.48555°E, 637 m, 10–19 Feb 1959, T.C. Maa, 1♀ (AMNH\_PBI 00041553) (BPBM); 13 Feb 1959, T.C. Maa, 1♀ (AMNH\_PBI 00041552) (BPBM); 15 Feb 1959, T.C. Maa, 1♀ (AMNH\_PBI 00041554) (BPBM).

***Coridromius hermosus*, new species**  
Figures 4, 6, 12D–F, 13C–H, map 8

**HOLOTYPE:** Male: **PAPUA NEW GUINEA: West Sepik Province:** Victor Emanuel Range: Bokubet, 5.16666°S 141.66666°E, 2850 m, 04–09 Jul 1976, J.L. Gressitt (AMNH\_PBI 00041507) (BPBM type# 16730) (BPBM).

**DIAGNOSIS:** *C. hermosus* is recognized by the following characters: proepisternum unilobed; frons and vertex glossy black without vittae; vertex nearly flat; antenna entirely black; scutellum orange; metepimeron extended as a long triangular lobe; metafemur less strongly swollen than in other species; saddlelike female paragenital structure on laterotergites II and III. *C. hermosus*, *epithema*, and *nigrus* all share similar paragenitalia, yet can be told from one another by their diagnostic coloration and variations in metepimeral lobe shape.

**DESCRIPTION: COLORATION** (figs. 4, 6): Head: Mostly black above frontoclypeal suture, orange-yellow below; vertex and frons black with yellow piping along posterior margin and adjacent to eyes; vittae absent; mandibular plates orange; maxillary plates, buccula, and gena pale with orange flecks; clypeus orange to orange-brown; labium pale, brown apically. Antenna: Entirely black. Thorax: Pronotal collar yellow, fading to black along margins; pronotum with pale medial stripe and large subanterior lateral black maculations, remainder orange-brown,


margins with distinct yellow piping; mesoscutum black; scutellum orange; mesepisternum orange-brown, lower half black and smooth. Hemelytra: Mostly brown; clavus mostly dark brown; corium mostly pale brown with obscure dark maculations; embolium flecked with orange to almost entirely orange; membrane dull brown, veins darker. Abdomen: Orange-brown, lateral margins orange-chestnut fading to yellow ventrally. Legs: Pro- and mesofemora chestnut apically, remainder pale; metafemur pale with 7 dark diagonal stripes; metatibia pale. SURFACE AND VESTITURE (figs. 4, 6): Impunctate; head and dorsum with sparse distribution of short, pale, white, decumbent setae. STRUCTURE (figs. 4, 6, 13C–E): Head: Broad, weakly tumescent medially; vertex without small tubercles adjacent to eyes; posterior margin of head nearly flat, with paired transverse foveae adjacent to eyes. Thorax: Pronotum broad, anterolateral margins weakly upturned and explanate, submarginal region of humeral angles weakly excavate, collar region undifferentiated; proepisternum unilobed (fig. 13C); posterior margin of metepimeron extended in a long, triangular lobe (fig. 13E); metanotum not flared (fig. 13D). Hemelytra: Costal margin thin and flared over its entire length and weakly sinuate. Legs: Metafemur only somewhat incrassate; metatibial spines short and thin. Abdomen: When viewed laterally, abdominal sternite II not distinctly angular. MALE GENITALIA (figs. 12D–F, 13G–H): Right paramere triangular and elongate (figs. 12E–F, 13G); left paramere sharply curved basally then continuing mostly straight, recurved at apex (figs. 12D, 13G); left ventral margin of pygophore with short mesal suture and broad ventral apical process (fig. 13H). FEMALE PARAGENITALIA (fig. 13E–F): Right laterotergites II and III swollen, laterotergite II transversely striate, depressed and membranous (fig. 13E–F), site of copulation thought to be intersegmental behind sternite II.

**ETYMOLOGY:** This species name is derived from Spanish, meaning “beautiful”.

**HOST:** Unknown.

**DISTRIBUTION:** West Sepik Province, Papua New Guinea (map 8).

**PARATYPES: PAPUA NEW GUINEA: West Sepik Province:** Victor Emanuel Range:



Sue Lindsay, Australian Museum

Fig. 13. Scanning electron micrographs: **A–B:** *C. falsicoleus*: **A.** ♀ thorax and abdomen, with arrow indicating paragenital opening, right lateral view. **B.** Detail of paragenital opening. **C–H:** *C. hermosus*: **C.** Head and pronotum, lateral view. **D.** ♀ scutellum and abdomen, with hemelytra removed to expose swollen right laterotergites II and III (arrow), dorsal view. **E.** ♀ metepimeron, right lateral view. **F.** Detail of right laterotergites II and III. **G.** ♂ genitalia, dorsal view. **H.** ♂ pygophore, ventral view. LP = left paramere, MEP = metepimeron, MES = mesepimeron, MET = metepimeron, MN = metanotum, P = peritreme, PES = proepisternum, RP = right paramere, VAP = ventral apical process.

Bokubet, 5.16666°S 141.66666°E, 2850 m, 04–09 Jul 1976, J.L. Gressitt, 4♂ (AMNH\_PBI 00041508–AMNH\_PBI 00041511), 5♀ (AMNH\_PBI 00041512–AMNH\_PBI 00041516) (BPBM).

*Coridromius lestoni*, new species

Figures 4, 6, 12G–I, map 2

HOLOTYPE: Male: GHANA: Ashanti: Tafo, 6.7333°N 1.6167°W, 278 m, 28 Jul 1967, D. Leston (AMNH\_PBI 00178008) (BMNH).

DIAGNOSIS: Recognized by its small size, near-uniform dark brown coloration, uniquely flared hemelytra, and the distinctive groove running along the lateral abdominal tergites.

DESCRIPTION: COLORATION (figs. 4, 6): Mostly dark brown with yellow markings. Head: Mostly light to dark orange-brown, frons with faint, dark brown vittae; inner ocular margins and posterior margin of head with yellow piping; depressions above and below lateral tumescences dark brown; clypeus yellow to orange-brown, darker apically, with faint brown markings basally; mandibular plates yellow; maxillary plate and buccula dark brown with faint yellowing on lower margins; gena yellow; labrum red; labium orange-brown, becoming darker at apex. Antenna: A1 yellow with broad subbasal dark brown annulation; AII light yellow-brown with yellow subapical annulation and dark brown annulation; AIII and AIV missing. Thorax: Pronotal collar dark brown; pronotum dark brown, only faintly paler along midline, lateral and posterior margins yellow; thoracic pleura dark orange-brown and dark brown; proepisternum with yellow lobes; posterior margin of propleuron yellow; posterior margin of metepimeron faintly yellow; mesoscutum dark brown; scutellum dark brown with yellow apex. Hemelytra: Dark brown, basally and laterally slightly paler; fascia at apex of embolium yellow; cuneus dark brown; membrane faintly brown with darker brown veins. Legs: Procoxae yellow; meso- and metacoxa dark brown, trochanters pale; pro- and mesofemora dark brown with yellow apices; metafemur light orange-brown to dark brown, with contrasting yellow and dark brown

diagonal stripes on outer surface, three bothria colored yellow; pro- and mesotibiae yellow with two or three irregular dark brown markings basally, apically tinted orange; metatibia yellow with irregular dark brown banding; tarsi yellow. Abdomen: Orange-brown, posterior margins of sternites black, dorsolateral margins with black and yellow markings. SURFACE AND VESTITURE (figs. 4, 6): Head, hemelytra, and metepimeron impunctate; pronotum and scutellum with deep, widely spaced punctuation; propleuron faintly punctate; head and dorsum clothed in long, white, decumbent setae. STRUCTURE (figs. 4, 6): Head: Frons weakly tumescent medially, swelling of frons extending to vertex; lateral swellings on vertex adjacent to eyes broad with well-defined depressions above and below; posterior margin of vertex carinate and rounded. Thorax: Pronotum broad, posteriorly rounded and declivit, anterolateral margins not upturned, depressed adjacent to callar region, submarginal region of humeral angles strongly excavate, callar region weakly defined; proepisternum bilobed; posterior margin of metepimeron truncate, apical angle forming a small, rounded lobe; metanotum prominent and flared; scutellum flattened. Hemelytra: Embolium carinate and sharply flared outward posteriorly, with a distinct raised fascia at apex of R+M vein; costal fracture deep. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally posterior margin of sternite II not sharply angular; dorsolateral margins of laterotergites with shallow, sclerotized groove running lengthwise along the abdomen. MALE GENITALIA (fig. 12G–I): Right paramere small, triangular, and curved (fig. 12H–I); left paramere scythelike, apex broadly acute (fig. 12G); posterior margin of pygophore with long mesal suture and long, fingerlike ventral apical process. FEMALE PARAGENITALIA: No visible external paragenitalia.

ETYMOLOGY: Named after Dennis Leston, who collected both the holotype male and the paratype female.

REMARKS: The shape of the hemelytra and the longitudinal grooves along the dorsolateral margins of the abdomen are unique to this species.

HOST: Unknown.

**DISTRIBUTION:** This species is known only from Ghana (map 2).

**PARATYPE: GHANA: Ashanti:** Burso, 6.69°N 1.63°W, 08 Jul 1969, D. Leston, Pyrethrum knockdown, 1♀ (AMNH\_PBI 00178009) (BMNH).

*Coridromius marmoreus*, new species  
Figures 5, 6, 12J–L, 14A–C, map 3

**HOLOTYPE:** Male: **PHILIPPINES: Rizal:** Mount Montalban, Wa-wa Dam, 14.73016°N 121.185°E, 150–200 m, 02 Mar 1965, H.M. Torrevillas (AMNH\_PBI 00041573) (BPBM type# 16731) (BPBM).

**DIAGNOSIS:** Easily distinguished from other species by its unique mottled coloration and thin, platelike pronotal margins. This species is superficially similar to *chinensis*, *confusus*, *minusculus*, and *punctatus*, however, none of these other species has spots along the pronotal or hemelytral margins. Additionally, the right paramere of *marmoreus* lacks a spined apophysis, the posterior margin of the pygophore lacks a mesal suture, the anterior lobe of the proepisternum is much shorter than the posterior lobe and the propleuron lacks punctuation.

**DESCRIPTION: COLORATION** (figs. 5, 6): Head: With weak vittae on frons, pale with mottled brown maculations; clypeus with two submedial longitudinal brown stripes; mandibular plate yellow to orange above, brown below; maxillary plate yellow to orange, margins fading to brown; buccula yellow, somewhat brown to orange basally; labrum brown; labium orange-brown, darker brown apically. Antenna: AI cream with brown basal annulation; AII light orange-brown with slightly darker brown apical annulation; AIII and AIV brown. Thorax: Pronotal collar yellow with brown anterior and posterior margins; pronotum mostly yellow with brown punctures, humeral angles and collar region lightly brown, two small dark brown markings on anterior margin immediately behind eyes, two to four more transverse brown markings midway up pronotum, and two thin dark brown markings on posterior margin, lateral margins with three or four dark brown spots; mesoscutum brown; scutellum yellow-brown, mottled, with deep brown punctures, two dark brown

marks on lateral margins; thoracic pleura mainly light orange-brown with yellow posterior margins; proepisternum with apices of both projections yellow, the remainder light brown; mesepisternum pale above, abruptly becoming dark brown at venter. Legs: Mostly yellow, pro- and mesofemora with a few small brown markings and an irregular brown band near apex; metafemur yellow at base with two or three diagonal brown stripes blending into irregular brown mottling midway along femur, bothria dark brown; tibiae pale with irregular brown banding, metatibial spines dark brown. Hemelytra: Mottled yellow and brown with patterned darker and paler markings; clavus brown at mid-point, with a dark brown spot at the tip and irregular dark brown marks throughout; endocorium and embolium with irregular brown markings; lateral margins of embolium with irregular dark brown spots; cuneus with irregular, dark brown line following inner and posterior margin, lateral margin pale; membrane tan with veins brown. Abdomen: Reddish brown to yellowish brown, with brown maculations on each segment below spiracles. **SURFACE AND VESTITURE** (figs. 5, 6, 14A–B): Head irregularly punctate, especially in shallow depressions adjacent to eyes; pronotum, scutellum, hemelytra, and metafemur irregularly punctate; propleuron impunctate (fig. 14A); metepimeron impunctate (fig. 14B); head and dorsum sparsely covered in short, pale, decumbent setae. **STRUCTURE** (figs. 5, 6, 14A–B): Head: Highly contoured; frons medially tumescent and laterally depressed, with minor swellings adjacent to eyes; depressions adjacent to eyes shallow; posterior margin of head broadly rounded medially, not carinate. Thorax: Pronotum broad, collar region weakly defined, anterolateral margins weakly explanate, lateral and posterior margins thin and platelike, submarginal region of humeral angles weakly excavated, posterior margin weakly upturned; proepisternum bilobed, with anterior projection much shorter than posterior (fig. 14A); posterior margin of metepimeron with small lobe projecting from upper corner (fig. 14B); metanotum prominent and flared. Hemelytra: Costal margin evenly rounded, thin, and platelike. Legs: Metatibial spines long and thick. Abdomen:

When viewed laterally, posterior margin of abdominal sternite II not angular. MALE GENITALIA (figs. 12J–L, 14C): Right paramere rectangular (fig. 12K–L); left paramere scythe-shaped, apex very slender (figs. 12J, 14C); posterior margin of pygophore without groove, fold, or mesal suture, ventral apical process absent (fig. 14C). FEMALE PARAGENITALIA: No visible external paragenitalia.

**ETYMOLOGY:** This species is named from the Latin *marmoreus*, referring to its mottled coloration.

**HOST:** Unknown.

**DISTRIBUTION:** Collected in large numbers in Luzon, Philippines, with single specimens also found in Papua New Guinea and north Queensland, Australia (map 3).

**PARATYPES:** **PHILIPPINES:** Rizal: Luzon, Montalban Dam, 14.73016°N 121.185°E, 16 Mar 1960, T. C. Maa, 1♀ (AMNH\_PBI 00041624) (AM). Mount Montalban, Wa-wa Dam, 14.73016°N 121.185°E, 175 m, 02 Mar 1965, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041621) (AM). 3♂ (AMNH\_PBI 00041577–AMNH\_PBI 00041579) (AMNH). 3♂ (AMNH\_PBI 00041574–AMNH\_PBI 00041576), 1♀ (AMNH\_PBI 00041580) (BPBM); 03 Mar 1965, H. M. Torrevillas, 3♂ (AMNH\_PBI 00041581–AMNH\_PBI 00041583), 1♀ (AMNH\_PBI 00041584) (BPBM); 06 Mar 1965, H. M. Torrevillas, 2♂ (AMNH\_PBI 00041588, AMNH\_PBI 00041589), 1♀ (AMNH\_PBI 00041590) (AMNH). 3♂ (AMNH\_PBI 00041585–AMNH\_PBI 00041587) (BPBM); 08 Mar 1965, H. M. Torrevillas, 2♀ (AMNH\_PBI 00041592, AMNH\_PBI 00041593) (AMNH). 1♂ (AMNH\_PBI 00041591) (BPBM); 09 Mar 1965, H. M. Torrevillas, 2♀ (AMNH\_PBI 00041594, AMNH\_PBI 00041595) (AMNH). 1♀ (AMNH\_PBI 00041596) (BPBM); 14 Mar 1965, H. M. Torrevillas, 2♀ (AMNH\_PBI 00041597, AMNH\_PBI 00041598) (BPBM); 16 Mar 1965, H. M. Torrevillas, 1♀ (AMNH\_PBI 00041599) (BPBM); 17 Mar 1965, H. M. Torrevillas, 6♂ (AMNH\_PBI 00041600–AMNH\_PBI 00041605), 1♀ (AMNH\_PBI 00041606) (BPBM); 18 Mar 1965, H. M. Torrevillas, 1♀ (AMNH\_PBI 00041623) (AM). 1♂ (AMNH\_PBI 00041607) (BPBM); 19 Mar 1965, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041608), 2♀ (AMNH\_PBI 00041609, AMNH\_PBI 00041610) (BPBM); 22 Mar 1965, H. M. Torrevillas, 1♀ (AMNH\_PBI 00041611) (BPBM); 23 Mar 1965, H. M. Torrevillas, 3♀ (AMNH\_PBI

00041612–AMNH\_PBI 00041614) (BPBM); 24 Mar 1965, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041615), 2♀ (AMNH\_PBI 00041616, AMNH\_PBI 00041617) (BPBM); 25 Feb 1965, H. M. Torrevillas, 1♀ (AMNH\_PBI 00041619) (AM). 1♂ (AMNH\_PBI 00041618) (BPBM); 08 Mar 1965–12 Mar 1965, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041620) (AM); 04 Mar 1965, H. M. Torrevillas, 1♂ (AMNH\_PBI 00041622) (AM).

**OTHER SPECIMENS EXAMINED:** **AUSTRALIA:** **Queensland:** 1 km SE of Mt. Cook, 15.3°S 145.16°E, 13 Oct 1980, T. Weir, 1♂ (AMNH\_PBI 00201295) (QM). Packers Creek, via Portland Roads, 16.48333°S 145.46666°E, 06 Dec 1985, G. Monteith and D. Cook, Pyrethrum knockdown, 1♀ (QM). **PAPUA NEW GUINEA:** **Madang Prov.:** Baiteta, 5.017°S 145.75°E, 01 Jul 1996, O. Missa, Light Trap, 1♂ (ULB).

*Coridromius minusculus* Carvalho  
Figures 5, 6, 12M–O, map 8

*Coridromius minusculus* Carvalho, 1987: 62–63.

**DIAGNOSIS:** Distinguished by the following combination of characters: small size; bilobed proepisternum; apex of scutellum somewhat swollen and rounded; mottled coloration; AI pale with basal dark brown annulation; AII with dark brown postmedial annulation; right paramere with sharp, triangular spicule; left paramere thin and scythe-like with gutter open its entire length. The female of *minusculus* is not known.

**REDESCRIPTION:** **COLORATION** figs. 5, 6: Head: Mottled light orange-brown and whitish yellow; with faint light brown vittae; depressions on vertex dark brown; clypeus whitish yellow with two irregular dark brown stripes; mandibular and maxillary plates whitish yellow, becoming dark brown along posterior margins; gena yellow; buccula whitish yellow, dark brown basally; labrum dark orange-brown, labium light orange-brown, darkened apically. Antenna: AI whitish yellow with dark brown annulation near base, AII light orange-brown with whitish subapical band followed by dark brown apical annulation, AIII and AIV dark brown with yellow bases. Thorax: Pronotal collar whitish yellow; pronotum whitish yellow with mottled orange-brown punc-




Fig. 14. Scanning electron micrographs: **A–C:** *C. marmoreus*: **A.** Head and pronotum, lateral view. **B.** Thorax, lateral view. **C.** ♂ genitalia, posterior view. **D.** *C. monotocopsis* ♂ genitalia, posteroventral view. **E–F:** *C. nakatanii*: **E.** ♂ genitalia, dorsal view. **F.** ♂ genitalia, posterior view. **G–H:** *C. neoguineanus*: **G.** ♂ genitalia, posteroventral view. **H.** ♂ genitalia, posterior view. LP = left paramere, MES = mesepimeron, MET = metepimeron P = peritreme, PES: proepisternum, RP = right paramere, VAP = ventral apical process.

Sue Lindsay, Australian Museum

tures, darkest around callar region, with two small dark brown markings along anterior behind eyes; with indistinct pale medial stripe; margins without pale piping; mesoscutum orange-brown; scutellum whitish yellow with mottled orange-brown punctures, apex pale; thoracic pleura dark orange-brown, whitish yellow along caudal margins; proepisternal lobes whitish yellow. Hemelytra: Mostly orange-brown, anterior portion of clavus slightly darker, becoming paler at commissure; corium with faint darker brown patches, becoming paler at apex; embolium with broad whitish-yellow subbasal and apical spots; cuneus light orange-brown, pale at apex; membrane brown, with pale veins bounded by irregular darker brown markings. Legs: Mostly whitish yellow with brown markings; coxae whitish yellow; pro- and mesofemora with broad, subapical irregular brown markings, bothria whitish yellow; metafemur laterally with four or five irregular diagonal brown stripes basally, becoming blotchy and indistinct apically, irregular whitish-yellow patch anterodorsally, ventrally with dark brown patch below stripes, apical lateral depressions orange-brown, bothria whitish yellow; all tibiae whitish yellow with irregular dark brown markings; metatibial spines dark brown. Abdomen: Light orange-brown, posterior margin of sternite II dark orange-brown. SURFACE AND VESTITURE (figs. 5, 6): Irregular depressions on head with deep punctures, somewhat coalescing; pronotum and propleuron deeply, irregularly punctate; scutellum deeply, irregularly rugulopunctate, impunctate along midline; metepimeron and hemelytra impunctate; clothed in long white setae. STRUCTURE (figs. 5, 6): Head: Highly contoured, frons strongly tumescent, bounded by irregular depressions and three minor swellings laterally and posteriorly; foveae adjacent to eyes deep, posterior margin of vertex medially rounded, not carinate; gena swollen; mandibular plate tumescent. Thorax: Pronotum broad and somewhat elongate, posteriorly rounded, steeply declivit, anterolateral margins weakly carinate; submarginal region of humeral angles strongly excavate; posterior of disc steeply declivit; posterior margin weakly medially cleft; callar region weakly devel-

oped; proepisternum bilobed; posterior margin of metepimeron truncate; metanotum prominent and flared; scutellum somewhat flattened, apex somewhat rounded. Hemelytra: Costal margin evenly rounded over its entire length. Legs: Metatibial spines long and thick. Abdomen: Posterolateral margin of abdominal sternite II not sharply angular. MALE GENITALIA (fig. 12M–O): Right paramere rounded with a long, spined apophysis (fig. 12N–O); left paramere scythe-shaped, slightly twisted along its axis, laterally compressed at base, thin apically, gutter open over entire length (fig. 12M); margin of pygophore with long mesal suture, ventral apical process present and long. FEMALE PARAGENITALIA: Female unknown.

REMARKS: Similar to *chinensis*, *confusus*, and *punctatus* (see remarks for *chinensis* above). Although the name refers to this species' diminutive size, given the amount of size variation we have seen in other species, we cannot be certain that *minusculus* is truly considerably smaller than other species.

HOST: One specimen has been collected from the family Cyperaceae, but we suspect that this represents a sitting record, as no other *Coridromius* species appear to feed on monocots (but see table 2 for another record on Cyperaceae).

DISTRIBUTION: Found in New Guinea (map 8).

HOLOTYPE: Male: INDONESIA: Papua: Boden, 11 km SE Oberfaren, 1.96666°S 138.73333°E, 1000 m, 7–17 Jul 1959, T.C. Maa, (AMNH\_PBI 00042152) (BPBM).

OTHER SPECIMENS EXAMINED: PAPUA NEW GUINEA: Morobe: Bulolo Gorge, 7.19999°S 146.64999°E, 1100 m, 15 Nov 1981, J.L. Gressitt, (Cyperaceae), 1♂ (AMNH\_PBI 00042147) (BPBM).

*Coridromius monotocopsis*, new species  
Figures 1E–F, 5, 6, 14D, 15A–C, map 4

HOLOTYPE: Male: AUSTRALIA: New South Wales: Munmorah State Rec. Area, 33.21916°S 151.57083°E, 11 Dec 1996, L. Wilkie, *Monotoca elliptica* (Epacridaceae), (AMNH\_PBI 00012127) (AM).

DIAGNOSIS: Recognized by the following combination of characters: proepisternum unilobed, metanotum not prominently flared;

posterior margin of pygophore biconvex, with fold on left side forming a shallow U-shaped groove, without small process on right margin of groove; light orange-brown coloration, males generally with dark brown coloring on side of abdomen. *C. monotocopsis* is similar to both Australian species *chenopoderis* and *pilbarensis*, and also resembles *bicolor* and the type species *variegatus*. Of the Australian species, *monotocopsis* is of more or less intermediate size between the other two, broader than *chenopoderis*, and typically uniformly light orange-brown in coloration. Males can be told apart by their relatively longer left paramere and much shallower U-shaped fold on the posterior margin of the pygophore, as well as by the distinctive dark brown band on the sides of the abdomen of most specimens.

**DESCRIPTION: COLORATION** (figs. 5, 6): Mostly orange-brown with darker brown markings, females usually paler than males. Head: Mostly orange-brown, becoming paler ventrally; dark brown vittae on frons; lateral margins bordering eyes and posterior margin of vertex yellow, lateral tumescences on vertex yellow, depressions behind tumescences dark brown; clypeus lighter orange-brown with faint median and lateral brown markings; maxillary plate and buccula yellow; labrum light orange-brown; labium yellow, brown apically. Antenna: AI yellow with thin light orange-brown bands apically and basally; AII orange-brown, not brown apically; AIII and AIV orange-brown, darker than AII. Thorax: Pronotal collar yellow-brown to orange-brown; pronotum orange-brown with faint yellow medial stripe and margins, in males collar region with irregular dark brown to black markings; mesoscutum orange-brown; scutellum orange-brown with white markings at basal angles and on either side of apex, in males sometimes with a broad, dark brown medial marking that sometimes extends over the entire scutellum; thoracic pleura orange-brown with yellow margins, in males sometimes becoming dark brown. Hemelytra: Yellow-brown to orange-brown, sometimes with faint reddish-orange markings at apex of embolium and on cuneus; in males inner margins of clavus and claval commissure dark brown, sometimes fusing laterally with dark brown chevron across

corium; cuneus becoming yellow; membrane faintly brown with orange to brown veins. Legs: Coxae yellow; pro- and mesofemora yellow with faint, irregular subapical brown band; hind femur yellow with 8–9 dark brown diagonal stripes on outer surface, with broad, brown marking ventrally, darker in males; tibiae yellow to orange-yellow; tarsi brown apically. Abdomen: Yellow in females, sometimes slightly reddened along ovipositor; in males yellow with lateral dark brown triangular band tapering toward apex, sometimes with bright reddish-orange tinting on and around genital capsule. **SURFACE AND VESTITURE** (figs. 5, 6): Head glossy with shallow punctures and very short, decumbent setae; pronotum with shallow punctuation; proepisternum, metepimeron and hemelytra impunctate; scutellum weakly rugulose; dorsum sparsely covered in short, simple, white, decumbent setae. **STRUCTURE** (figs. 5, 6): Head: Broad; frons weakly tumescent medially; with paired tubercles adjacent to eyes, bordered posteriorly by shallow depressions; posterior margin of head slightly rounded medially but otherwise flat. Thorax: Pronotum broad and rounded, margins thinly carinate, anterolateral margins weakly upturned, posterior margin not medially cleft, submarginal region of humeral angles weakly excavate; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared. Hemelytra: Lateral margin of embolium thin and evenly rounded, nearly straight. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. **MALE GENITALIA** (figs. 14D, 15A–C): Right paramere broad, short, triangular (fig. 15B–C); left paramere sickle-shaped, strongly attenuate (figs. 14D, 15A); posterior margin of pygophore with shallow folded groove to left of midline (fig. 14D). **FEMALE PARAGENITALIA:** No visible external paragenitalia; site of copulation unknown.

**ETYMOLOGY:** Named for its host plant, *Monotoca elliptica*.

**HOST:** Found almost exclusively on *Monotoca elliptica* (Epacridaceae).

**DISTRIBUTION:** Found primarily in eastern Australia, but also known from four speci-

mens collected on Norfolk Island, where we believe it has been introduced (map 4).

**PARATYPES: AUSTRALIA:** New South Wales: Booti Booti NP, 32.27972°S 152.52444°E, 09 Oct 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 4♂ (AMNH\_PBI 00012228, AMNH\_PBI 00012228–AMNH\_PBI 00012229, AMNH\_PBI 00012229), 3♀ (AMNH\_PBI 00012230–AMNH\_PBI 00012232) (AM); 02 Apr 1998, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012233) (AM); 08 Oct 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♀ (AMNH\_PBI 00012234) (AM); 12 Nov 1996, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 3♂ (AMNH\_PBI 00012235–AMNH\_PBI 00012237) (AM); 30 May 1997, L. Wilkie, 1♂ (AMNH\_PBI 00012238) (AM). Callala Beach, 35.015°S 150.701°E, 18 May 1997, G. Cassis, *Monotoca* sp. (Epacridaceae), 11♂ (AMNH\_PBI 00012256–AMNH\_PBI 00012266), 5♀ (AMNH\_PBI 00012267–AMNH\_PBI 00012271) (AM). Munmorah State Rec. Area, 33.21916°S 151.57083°E, 11 Dec 1996, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 32♂ (AMNH\_PBI 00012123–AMNH\_PBI 00012126, AMNH\_PBI 00012128–AMNH\_PBI 00012150, AMNH\_PBI 00012197–AMNH\_PBI 00012201), 46♀ (AMNH\_PBI 00012151–AMNH\_PBI 00012196) (AM); 18 Nov 1996, L. Wilkie, *Chrysanthemoides monilifera* (Asteraceae), 1♂ (AMNH\_PBI 00012209) *Monotoca elliptica* (Epacridaceae), 6♂ (AMNH\_PBI 00012202–AMNH\_PBI 00012207), 1♀ (AMNH\_PBI 00012208) (AM); 12 Oct 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 4♀ (AMNH\_PBI 00012210–AMNH\_PBI 00012213) (AM); 11 Oct 1997, L. Wilkie, 1♀ (AMNH\_PBI 00012214) (AM); 29 May 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 2♂ (AMNH\_PBI 00012215, AMNH\_PBI 00012216) (AM); 04 Apr 1998, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012217) (AM); 14 May 1998, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 2♂ (AMNH\_PBI 00012218, AMNH\_PBI 00012219) (AM); 15 Dec 1996, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012220) (AM); 23 May 1998, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012221) (AM); 05 Apr 1998, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012222) (AM); 01 May 1997, L. Wilkie, 1♀ (AMNH\_PBI 00012227) *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012223) (AM); 31 May 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012224), 1♀ (AMNH\_PBI 00012225) (AM); 27 Nov 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♀ (AMNH\_PBI 00012226) (AM). Myall Lakes NP, 32.57916°S 152.29083°E, 09 Oct 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 5♂ (AMNH\_PBI 00012240–AMNH\_PBI 00012244), 8♀ (AMNH\_PBI 00012245–AMNH\_PBI 00012252) *Acacia longifolia* (Mimosaceae), 1♂ (AMNH\_PBI 00012239) (AM); 29 Apr 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♂ (AMNH\_PBI 00012253) (AM); 10 Oct 1997, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♀ (AMNH\_PBI 00012254) (AM); 24 May 1998, L. Wilkie, *Monotoca elliptica* (Epacridaceae), 1♀ (AMNH\_PBI 00012255) (AM). Nulliga Beach, 37.06°S 149.53°E, 31 May 1989, G. Cassis, 2♂ (AMNH\_PBI 00012427, AMNH\_PBI 00012428), 1♀ (AMNH\_PBI 00015736) (AM). Royal National Park, Wattomolla Beach car park, 34.13334°S 151.1134°E, 29 m, 14 Nov 2001, Cassis, Schuh, Schwartz, Silveira, *Monotoca elliptica* (Sm.) R.Br. (Ericaceae), det. NSW staff NSW666424, 108♂ (AMNH\_PBI 00015256–AMNH\_PBI 00015333, AMNH\_PBI 00015436–AMNH\_PBI 00015465), 126♀ (AMNH\_PBI 00015334–AMNH\_PBI 00015435, AMNH\_PBI 00015466–AMNH\_PBI 00015489) (AM); 12 Apr 2004, G. Cassis, P. Tinerella, N. Tatarnic, N. Velez, K. McLachlan, *Monotoca elliptica* (Sm.) R.Br. (Ericaceae), 5♀ (AMNH\_PBI 00114228–AMNH\_PBI 00114232) (CNC). Queensland: East Woodmillar, 25.68333°S 151.6°E, 250 m, 17 Dec 1998, Monteith and Gough, 1♀ (AMNH\_PBI 00178068) (QM). Victoria: Little Desert National Park, 5–6 km W of McDonald Hiway, 36.61668°S 141.1667°E, 150 m, 03 Nov 1995, Schuh and Cassis, *Melaleuca wilsonii* F. Muell. (Myrtaceae), det. P.G. Wilson 1996 NSW395985, 10♂ (AMNH\_PBI 00015567–AMNH\_PBI 00015576), 16♀ (AMNH\_PBI 00015577–AMNH\_PBI 00015592) (AM). *Brachyloma daphnoides* (Smith) Benth. (Epacridaceae), det. E.A. Brown 1996 NSW395984, 3♂ (AMNH\_PBI 00002593–AMNH\_PBI 00002595), 5♀ (AMNH\_PBI 00002596–AMNH\_PBI 00002600) (AMNH).




Fig. 15. Left and right parameres of *Coridromius* spp.: A–C: *C. monotocopsis*. D–F: *C. nakatanii*. G–I: *C. neoguineanus*. J–L: *C. pilbarensis*. M–P: *C. prolixipenis*.

OTHER SPECIMENS EXAMINED: NORFOLK ISLAND: 29.033°S 167.949°E, A.M. Lea, 4 adults (AMNH\_PBI 00038583) (SAMA).

*Coridromius nakatanii* Chérot, Konstantinov and Yasunaga

Figures 5, 6, 14E–F, 15D–F, map 5

*Coridromius nakatanii* Chérot et al., 2004: 60–63.

DIAGNOSIS: Easily recognized by the strongly posteriorly flared embolium and long, thin left paramere, closed except at apex. *C. nakatanii* is superficially similar to *crassus*, *declivipennis*, and *pteraulos*, but all four can be told apart by the shape of the hemelytra and their coloration. Additionally, the left paramere of *nakatanii* is thinner, more twisted apically and enclosed until its apex, unlike in the other three species, and the female possesses a prominent paragenital opening within abdominal sternite II.

REDESCRIPTION: COLORATION (figs. 5, 6): Mostly brownish yellow to dark brown with yellow and dark brown markings. Head: Brownish yellow; very faint vittae on frons; lateral swellings adjacent to eyes yellow, merging with yellow ocular and posterior margins; clypeus yellow, brown apically, with one medial light brown spot and two lateral light brown stripes; gena, mandibular and maxillary plates yellow; buccula yellow, dark brown basally; labrum brown; labium yellow, brown apically. Antenna: AI dark brown, yellow at base and apex; AII light orange-brown, slightly darker at base, with dark brown medial band separated from broad, dark brown apical annulation by a thin yellow subapical annulation; AIII and AIV dark brown, yellow basally. Thorax: Pronotal collar yellow, darkened along anterior and posterior margins; pronotum mostly brownish yellow to nearly uniform dark brown, with yellow midline; paler specimens with two small dark brown spots on anterior margin behind eyes, two larger brown patches above callar region and the posterior margin somewhat darkened, darker specimens with faint yellowing submarginally at humeral angles, lateral and posterior margins yellow; mesoscutum in pale specimens orange-brown, becoming dark brown medially and laterally, in darker specimens uniform dark brown; scutellum brownish yellow

to dark brown with yellow midline, apex and sides yellow, basally with dark brown markings medially and laterally; thoracic pleura brownish yellow with some faint darker brown markings, sometimes nearly uniform dark brown, posterior margins of pleura yellowed; proepisternal lobes yellow. Hemelytra: Brownish yellow with punctures darkened, in dark specimens nearly uniform dark brown; clavus darker anteriorly, claval commissure slightly pale; broad, pale specimens with dark brown patch extending across corium and embolium, broken transversely by yellow medial fracture, all specimens with yellow C-shaped pattern near costal fracture; veins yellow above costal fracture; cuneus light to dark brown; membrane pale brown, slightly brown subapically, with dark brown veins enclosing cell. Abdomen: Brownish yellow. Legs: All coxae pale yellow; pro- and mesofemora pale yellow; metafemur yellow with 8 diagonal dark brown stripes on outer surface, dark brown patch mediolaterally, lateroapical depressions dark brown; tibiae yellow, metatibia slightly brown. SURFACE AND VESTITURE (figs. 5, 6): Head, pronotum, propleuron, metepimeron, scutellum and hemelytra finely punctate; head and dorsum clothed in long, white, decumbent setae. STRUCTURE: Head: Frons weakly tumescent medially, merging with minor median swelling on vertex; vertex with two minor swellings adjacent to eyes; posterior margin of head weakly medially rounded, somewhat carinate. Thorax: Pronotum broad and steep, lateral margins carinate, anterolateral margins not upturned, submarginal region of humeral angles excavate, posterior steeply declivent, callar region poorly defined; proepisternum bilobed; posterior margin of metepimeron in the form of a short, broadly rounded lobe; metanotum prominent and flared; scutellum flat dorsally. Hemelytra: Costal margin thinly carinate, greatly flared posteriorly. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of sternite II distinctly angular. MALE GENITALIA (figs. 14E–F, 15D–F): Right paramere broad and rounded with thumblike lobe, paramere appears U-shaped from above (figs. 14E–F, 15E–F); left paramere thick, elongate and sharply angled

basally, then evenly curved and tapering to apex with a slight twist along its axis, curved at tip, gutter tightly closed until apex (figs. 14E–F, 15D); pygophore with a long mesal suture and long, triangular ventral apical process (fig. 14E–F). FEMALE PARAGENITALIA: Female with prominent paragenital opening within abdominal segment II, opening directed caudally.

**REMARKS:** The possession of a prominent paragenital opening in abdominal sternite II is shared with *falsicoleus* and *sommelieri*, however in the latter two species this opening leads to a sclerotized copulatory tube projecting into the abdomen, whereas in *nakatanii* the paragenital opening is angled caudally.

HOST: Unknown.

DISTRIBUTION: Collected in Lao PDR, Vietnam, and India (map 5).

**HOLOTYPE:** Male: **LAO PEOPLE'S DEMOCRATIC REPUBLIC:** Sekong Prov.: Bolavens Plateau, N slope, ca. 10 km N. Mg Tha Theng, 15.5°N 106.43333°E, 500–700 m, 29–30 May 1996, Schillhammer (AMNH\_PBI 00189978) (NHMW).

**PARATYPES:** **INDIA:** Meghalaya: West Garo Hills, Nokrek NP, 25.49333°N 90.325°E, 950–1250 m, 09–17 May 1996, E. Jendek and O. Sausa, 1♀ (AMNH\_PBI 00189979) (NHMW). **VIETNAM:** Blao (Balao), 11.53333°N 107.8°E, 500 m, 14–21 Oct 1960, C. M. Yoshimoto, 1♀ (AMNH\_PBI 00042605) (BPBM). Tam Dao NP, 21.4536°N 105.6436°E, 900 m, 17–18 Jun 1999, Y. Nakatani, 5♂ (AMNH\_PBI 00190003–AMNH\_PBI 00190007), 1♀ (AMNH\_PBI 00190008) (TYCN).

*Coridromius neoguineanus* Carvalho  
Figures 5, 6, 14G–H, 15G–I, map 8

*Coridromius neoguineanus* Carvalho, 1987: 63 (n. sp.).

**DIAGNOSIS:** Recognized by the following combination of characters: body light tan with dark brown spots; right paramere club-shaped; posterior margin of the pygophore with a U-shaped groove to right of centre. The coloration and right paramere shape of this species immediately distinguish it from all others.

**REDESCRIPTION: COLORATION** (figs. 5, 6): Light tan with yellow and dark brown

markings. Head: Mostly light tan; frons without vittae; posterior margin of vertex and inner margins of eyes yellow; dark brown spot in middle of vertex; apex of clypeus dark brown, embrownment sometimes extending the length of clypeus, occasionally partway up frons; mandibular and maxillary plates, gena, and buccula light yellow; labrum dark brown; labium light yellow, brown apically. Antenna: AII yellow; AIII yellow to light brown, with dark brown annulation; AIV dark brown. Thorax: Pronotal collar yellow; pronotum light tan with yellow medial stripe and posterior margin, two dark brown spots behind eyes and two more on disc; mesoscutum light tan; scutellum light tan with two dark brown spots, anterolateral corners and apex yellow; thoracic pleura light tan, posterior margins yellow; mesepisternum and metepimeron with darker brown markings. Hemelytra: Light tan, clavus generally darker; with transverse dark brown band arcing across hemelytra through apex of claval commissure, sometimes broken into separate dark brown markings; membrane pale, veins brown. Legs: Coxae yellow; pro- and mesofemora yellow with light orange-brown subapical marking; tibiae yellow; metafemur yellow with faint brown subapical band and 7–8 transverse stripes on outer side. Abdomen: Yellow with light orange-brown and dark brown markings. SURFACE AND VESTITURE (figs. 5, 6): Head and pronotum densely and finely punctate; scutellum weakly rugulose; propleuron, metepimeron, and hemelytra impunctate; head and dorsum sparsely clothed in short, simple, white setae. STRUCTURE (figs. 5, 6): Head: Frons weakly tumescent medially, nearly flat; posterior margin of head rounded, not carinate. Thorax: Pronotum broad, anterolateral angles weakly upturned and carinate, submarginal region of humeral angles weakly excavate, posterior margin weakly upturned, collar region undifferentiated; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared. Hemelytra: Costal margin thin and platelike, slightly rounded and flared over its entire length; cuneus impunctate. Legs: Metatibial spines short and thin. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not sharply angular.

**MALE GENITALIA** (figs. 14G–H, 15G–I): Right paramere boot-shaped (fig. 15H–I); left paramere stout and sickle-shaped (figs. 14G–H, 15G); posterior margin of pygo-phore simple, with single distinct groove slightly to right of center (fig. 14G–H). **FEMALE PARAGENITALIA:** No visible external paragenitalia.

**REMARKS:** While the left paramere is similar in shape to that of many other species, the distinctive spotted coloration and unique boot-shaped right paramere set this species apart from others.

**HOST:** Unknown.

**DISTRIBUTION:** Known from Papua New Guinea and Western Papua, Indonesia (map 8).

**HOLOTYPE:** Male: **PAPUA NEW GUINEA:** **Western Highlands:** Minj, 5.85°S 144.66666°E, 1700 m, 1959, T.C. Maa (AMNH\_PBI 00042151) (BPBM).

**PARATYPES:** **INDONESIA:** **Papua:** Guega, W. of Swart Valley, 3.6°S 138.41666°E, 1200 m, 15 Nov 1958, J.L. Gressitt, 1♀ (AMNH\_PBI 00041463) (BPBM).

**OTHER SPECIMENS EXAMINED:** **INDONESIA:** **Papua:** Kulima, 4.41666°S 138.98333°E, 1400 m, 19–22 Feb 1960, T.C. Maa, 6 adults, sex undetermined (AMNH\_PBI 00185365) (USNM). **PAPUA NEW GUINEA:** Morobe: Wau, 7.3333°S 146.71667°E, 1200 m, 23 Jan 1978, W.C. Gagne, 1♀ (AMNH\_PBI 00041464), 5♂ (AMNH\_PBI 00041465–AMNH\_PBI 00041469) (BPBM).

*Coridromius nigrus* Carvalho  
Figures 5, 6, map 8

*Coridromius nigrus* Carvalho, 1987: 64–65.

**DIAGNOSIS:** Recognized by its nearly uniform black coloration, long, triangular metepimeral lobe and distinctive paragenitalia. This species appears very similar to *epithema* and *hermosus*, both of which share similar paragenital structures, head shape and hemelytral shape, but differ in coloration.

**REDESCRIPTION: COLORATION** (figs. 5, 6): Nearly uniform reddish black. Head: Reddish black; mandibular plates and apex of clypeus tinted red; maxillary plate yellow; buccula and gena orange-brown; ocular

margins and posterior margin of vertex yellow; labium yellow-orange, apex missing from specimen. Antenna: Grey-brown, AIII and AIV missing. Thorax: Pronotal collar black; pronotum reddish black, becoming more reddened laterally, posterior and lateral margins yellow-orange; mesoscutum and scutellum black with reddish tinting in basal and apical corners; thoracic pleura mainly reddish black, yellow-orange along posterior margins. Hemelytra: Reddish black, lighter laterally and apically; membrane dark grey. Abdomen: Reddish black. Legs: Coxae orange-brown; trochanters pale; pro- and mesofemora and tibiae reddish black; hind femur reddish black with faint black diagonal stripes; tarsi somewhat yellowed. **SURFACE AND VESTITURE** (figs. 5, 6): Impunctate, scutellum rugulose; head and dorsum with white, decumbent setae, most setae knocked off specimen. **STRUCTURE** (figs. 5, 6): Head: Wide; frons weakly tumescent medially; lateral tubercles adjacent to eyes small and ill-defined, bordered posteriorly by shallow depressions; posterior margin of vertex nearly flat. Thorax: Pronotum very broad, anterolateral angles weakly upturned, posterior and lateral margins carinate, submarginal region of humeral angles excavate, callar region undifferentiated; proepisternum unilobed; posterior margin of metepimeron extended in the form of a triangular lobe; metanotum not prominently flared. Hemelytra: Costal margin thin, flared out from base then abruptly angling medially at midpoint. Legs: Metatibial spines short and thin. Abdomen: Postrolateral margin of sternite II not angular. **MALE GENITALIA:** Male unknown. **FEMALE PARAGENITALIA:** Right laterotergites II and III swollen, desclerotized and medially depressed, posterior margin of right abdominal sternite II strongly carinate and flared outward (as in *epithema* and *hermosus*, figs. 11G–H, 13D–F).

**HOST:** Unknown.

**DISTRIBUTION:** Known from Wau, Morobe District, Papua New Guinea (map 8).

**HOLOTYPE:** Female: **PAPUA NEW GUINEA:** **Morobe:** Edie Creek, Wau, 7.31666°S 146.68333°E, 1700 m, 02 Apr 1966, J.L. Gressitt (AMNH\_PBI 00042150) (BPBM).

*Coridromius pilbarensis*, new species  
Figures 5, 6, 15J–L, 16A–B, map 4

**HOLOTYPE:** Male: **AUSTRALIA: Western Australia:** Pilbara Region: Base of Mt. Nameless, adjacent to gate, 22.72694°S 117.74916°E, 05 Jun 2004, M. Bulbert, N. Tatarnic and S. Lassau (AMNH\_PBI 00012358) (AM).

**DIAGNOSIS:** Recognized by the following combination of characters: unique orange, brown, and yellow coloration; proepisternum unilobed; metanotum not prominently flared; posterior margin of pygophore biconvex, with prominent fold on left side forming U-shaped groove, with small apophysis on right margin of groove. Of the Australian species *chenopoderis*, *monotocopsis* and *pilbarensis*, the latter is the largest and generally most consistent in its coloration. Males are readily told by the small ventral apical process adjacent to the deep folded groove on the posterior margin of the pygophore, while females typically have a faintly greenish abdomen.

**DESCRIPTION: COLORATION** (figs. 5, 6): Mostly orange with brown markings above, yellow beneath. Head: Mostly orange-yellow to yellow with dark brown vittae on frons; lateral margins bordering eyes and posterior margin of vertex yellow, lateral tumescences on vertex yellow, depressions behind tumescences dark brown, sometimes with an additional dark brown marking in between; clypeus with irregular median and lateral brown markings; labrum light brown; labium yellow, brown apically. Antenna: AI yellow with narrow light brown bands apically and basally; AII yellow to yellow-orange, slightly brown apically; AIII and AIV orange-brown. Thorax: Pronotal collar yellow with brown to black anterior and posterior margins; pronotum orange with faint yellow midline and margins; collar region with irregular dark brown to black markings; disk sometimes slightly darkened, sometimes with broad dark brown to black markings; mesoscutum orange, black anteriorly; scutellum orange with dark brown markings anteromedially and laterally, apex yellowed laterally; thoracic pleura yellow-orange with yellow margins; prosternum dark brown to black ventrally. Hemelytra: Yellow-orange to orange, lighter

apically; inner margins of clavus and claval commissure dark brown, fusing laterally with dark brown chevron across corium, sometimes with a faintly darkened spot; cuneus becoming yellow; membrane faintly brown with dark brown veins. Legs: Coxae yellow; pro- and mesofemora yellow to orange-yellow with faint, irregular subapical brown band; hind femur yellow with 8–9 dark brown diagonal stripes on outer surface, somewhat orange anteriorly below stripes; tibiae yellow to orange-yellow; tarsi brown apically. Abdomen: In females yellow or yellow-green; in males yellow with lateral brown and orange markings, with bright reddish-orange tinting on and around genital capsule. **SURFACE AND VESTITURE** (figs. 5, 6): Head glossy and shallowly punctate; pronotum shallowly punctate; propleuron, metepimeron, scutellum and hemelytra impunctate; head and dorsum sparsely covered in short, white, decumbent setae. **STRUCTURE** (figs. 5, 6): Head: Frons weakly tumescent medially; paired tubercles adjacent to eyes bordered posteriorly by shallow depressions; posterior margin of head slightly rounded medially but otherwise flat. Thorax: Pronotum broad and rounded, lateral margins thinly carinate, anterolateral angles weakly upturned, posterior margin medially cleft, submarginal region of humeral angles weakly excavate; proepisternum unilobed; posterior margin of metepimeron truncate; metanotum not prominent and flared. Hemelytra: Costal margin thin and weakly rounded, nearly straight. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. **MALE GENITALIA** (figs. 15J–L, 16A–B): Right paramere broad, short, triangular (fig. 15K–L); left paramere sickle-shaped, strongly attenuate (figs. 15J, 16A); posterior margin of pygophore with distinct fold on left forming a U-shaped groove (fig. 16A), the upper right corner forming a small process (fig. 16A–B). **FEMALE PARAGENITALIA:** No visible external paragenitalia; site of copulation unknown.

**ETYMOLOGY:** Named after the Pilbara Region, where the holotype was collected.


**HOST:** Most specimens have been collected from *Melaleuca* spp. (Myrtaceae), but *pilbar-*

*ensis* has also been collected in large numbers on *Dodonaea viscosa angustissima* (Sapindaceae) (see table 2).

**DISTRIBUTION:** Found on both the west and east coasts of Australia (map 4).

**PARATYPES: AUSTRALIA: New South Wales:** 27 km W of Retreat (30 km E Manilla), 30.66668°S 150.8333°E, 350 m, 24 Oct 1995, Schuh and Cassis, *Schinus areira* L. (Anacardiaceae), det. P.G. Wilson 1996 NSW395924, 1♂ (AMNH\_PBI 00012432), 1♀ (AMNH\_PBI 00012433) (AM). Palm Beach, Sydney, 33.598°S 151.326°E, 16 Oct 1988, Richard Bejsak, 1♀ (AMNH\_PBI 00012441) (AM). **Western Australia:** 11 km N of Coolgardie-Esperance Hiway on Kam-balda Road, 31.25231°S 121.5899°E, 320 m, 18 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Melaleuca sheathiana* W. Fitzg. (Myrtaceae), det. Perth staff PERTH-05671396, 2♂ (AMNH\_PBI 00012328, AMNH\_PBI 00012329), 1♀ (AMNH\_PBI 00012330) (AM). 17 mi N of Paynes Find, 29.01°S 117.68°E, 400 m, 10 Mar 1962, E. S. Ross and D. Q. Cavagnaro, 1♂ (AMNH\_PBI 00002644) (AMNH). 20.6 km S of Norseman-Lake King Road on Lake King-Cascades Road, 33.16284°S 120.2813°E, 400 m, 22 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Gastrolobium spinosum* Benth. (Papilionaceae), det. Perth staff PERTH-05671736, 1♀ (AMNH\_PBI 00012331) (AM). 3 km S of Kojonup, Sampson Road, 33.87088°S 117.1648°E, 310 m, 08 Dec 1997, Schuh, Cassis, Brailovsky, Asquith, *Melaleuca rhamphophylla* Schauer (Myrtaceae), det. Perth staff PERTH-05879183, 1♀ (AMNH\_PBI 00012429) (AM). 31.3 km N of Ravensthorpe, 33.36519°S 119.8792°E, 500 m, 05 Nov 1996, Schuh and Cassis, *Santalum lanceolatum* R.Br. (Santalaceae), det. Perth staff PERTH05236665, 3♂ (AMNH\_PBI 00012341-AMNH\_PBI 00012343), 1♀ (AMNH\_PBI 00012344) (AM). 1♂ (AMNH\_PBI 00002527), 11♀ (AMNH\_PBI 00002528-AMNH\_PBI 00002538) (AMNH). 33.3 km S of Norseman, 32.46461°S 121.6778°E, 300 m, 19 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Santalum acuminatum* R. Br. (Santalaceae), det. Perth 05671094, 8♂ (AMNH\_PBI 00012272-AMNH\_PBI 00012279), 7♀ (AMNH\_PBI 00012280-AMNH\_PBI 00012286) (AM). 4.3 km N of Peak Charles National Park, 32.81408°S 121.2114°E, 200 m, 20 Nov 1999, R.T. Schuh

and G. Cassis, *Dodonaea viscosa angustissima* (Sapindaceae), det. Perth staff PERTH-05670705, 1♂ (AMNH\_PBI 00012430) (AM). 43 km N of Norseman, 31.85648°S 121.6414°E, 300 m, 19 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Santalum acuminatum* (R.Br.) A.DC. (Santalaceae), det. Perth staff PERTH-05671094, 6♂ (AMNH\_PBI 00012333-AMNH\_PBI 00012338), 2♀ (AMNH\_PBI 00012339, AMNH\_PBI 00012340) (AM). *Atriplex bunburyana* F.Muell. (Chenopodiaceae), det. Perth staff PERTH05671205, 7♂ (AMNH\_PBI 00002565-AMNH\_PBI 00002571), 9♀ (AMNH\_PBI 00002572-AMNH\_PBI 00002580) (AMNH). 56.6 km W of Yalgo, 28.42397°S 116.1233°E, 600 m, 27 Oct 1996, Schuh and Cassis, *Melaleuca uncinata* R.Br. (Myrtaceae), det. Perth staff PERTH05120640, 1♀ (AMNH\_PBI 00012345) (AM). Augusta, 34.34212°S 115.1661°E, 30 m, 04 Dec 1999, R.T. Schuh and G. Cassis, 1♂ (AMNH\_PBI 00012297) (AM). Cosy Corner Beach East, Torbay Sound, W of Albany, 35.06033°S 117.6446°E, 2 m, 01 Dec 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Santalum acuminatum* R. Br. (Santalaceae), det. Perth staff PERTH05677094, 11♂ (AMNH\_PBI 00002549-AMNH\_PBI 00002559), 5♀ (AMNH\_PBI 00002560-AMNH\_PBI 00002564) (AMNH). Hamersley Inlet Campground, sea level, 33.95786°S 119.91456°E, 28 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Scaevola crassifolia* Labill. (Goodeniaceae), det. Perth staff PERTH05672295, 1♀ (AMNH\_PBI 00012332) (AM). Lake Magenta Road, 4 km N of South Coast Hiway, 33.77872°S 119.2887°E, 400 m, 07 Dec 1997, Schuh, Cassis, Brailovsky, Asquith, *Melaleuca hamulosa* Turcz. (Myrtaceae), det. Perth staff PERTH05054567, 5♀ (AMNH\_PBI 00012324-AMNH\_PBI 00012327, AMNH\_PBI 00012431) (AM). *Melaleuca hamulosa* Turcz. (Myrtaceae), det. Perth staff PERTH-05054567, 1♀ (AMNH\_PBI 00002745) (AMNH). Mt. Lindesay, 23 km NW of Denmark, 34.85058°S 117.2767°E, 100 m, 17 Dec 1997, Schuh, Cassis, Brailovsky, *Agonis linearifolia?* (DC.)Sweet (Myrtaceae), det. Perth staff PERTH05095360, 1♂ (AMNH\_PBI 00002539) (AMNH). Peak Charles National Park Camp-ground, 32.88335°S 121.1703°E, 300 m, 20 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Melaleuca uncinata* R.Br. (Myrtaceae), det. Perth staff PERTH05670721, 4♂ (AMNH\_



Sue Lindsay, Australian Museum

Fig. 16. Scanning electron micrographs: A–B: *C. pilbarensis*: A. ♂ genitalia, posterior view. B. ♂ detail of ventral apical process of pygophore. C–H: *C. pteraulos*: C. ♀ scutellum and abdomen, wings removed, dorsal view. D. Detail of ♀ dorsum, showing flaring of metanotum. E. ♂ genitalia and abdomen, posterolateral view. F. Detail of conelike setae on ♂ abdominal sternites VII and VIII. G. ♀ thorax and abdomen, right lateral view. H. ♀ anterior of right embolium, lateral view. Arrow indicates half-tube-shaped margin. LP = left paramere, MES = mesepimeron, MET = metepimeron, MLS = mesal longitudinal suture, MN = metanotum, P = peritreme, RP = right paramere, VAP = ventral apical process.

PBI 00012287-AMNH\_PBI 00012290), 6♀ (AMNH\_PBI 00012291-AMNH\_PBI 00012296) (AM). *Dodonaea viscosa angustissima* (DC.) J.G. West (Sapindaceae), det. Perth staff PERTH05670705, 8♂ (AMNH\_PBI 00012298-AMNH\_PBI 00012305), 18♀ (AMNH\_PBI 00012306-AMNH\_PBI 00012323) (AM). Pilbara Region: Base of Mt Nameless, adjacent to gate, 22.72694°S 117.74916°E, 05 Jun 2004, M. Bulbert, N. Tatarnic and S. Lassau, 4♂ (AMNH\_PBI 00012360-AMNH\_PBI 00012363), 10♀ (AMNH\_PBI 00012364-AMNH\_PBI 00012373) (AM). Pilbara Region: Karijiji NP, Dales Gorge, Fortescue falls, 22.47694°S 118.54944°E, 07 Jun 2004, M. Bulbert, 1♂ (AMNH\_PBI 00012359) (AM).

**OTHER SPECIMENS EXAMINED: AUSTRALIA:** **Australian Capital Territory:** Canberra, 35.2833°S 149.2167°E, 605 m, 1973, J.C.M. Carvalho, 1♀ (AMNH\_PBI 00192745) (BMNH). **New South Wales:** Bogan River, J. Armstrong, 1♂ (AMNH\_PBI 00192747), 1♀ (AMNH\_PBI 00192746) (BMNH). Cowra, 33.819°S 148.658°E, 25 Mar 1960, M. I. Nikitin, Light Trap, 1♂ (AMNH\_PBI 00192755), 1♀ (AMNH\_PBI 00192744) (BMNH). Saltwater Reserve, 15 km SE of Taree, littoral rainforest, 31.915°S 152.451°E, 10 Dec 1991, G. Williams, *Euroschinus falcata* (Anacardiaceae), 1♀ (AMNH\_PBI 00012435) (AM); 03 Dec 1991, G. Williams, *Euroschinus falcata* (Anacardiaceae), 1♂ (AMNH\_PBI 00001236), 1♀ (AMNH\_PBI 00001237) (AM); 26 Nov 1991, G. Williams, *Euroschinus falcata* (Anacardiaceae), 1♀ (AMNH\_PBI 00001238) (AM). **Western Australia:** Peak Charles National Park Campground, 32.88335°S 121.1703°E, 300 m, 20 Nov 1999, R.T. Schuh, G. Cassis, and R. Silveira, *Melaleuca uncinata* R.Br. (Myrtaceae), det. Perth staff PERTH05670721, 10 juveniles (AMNH\_PBI 00015533-AMNH\_PBI 00015542) (AM).

*Coridromius prolixipenis*, new species  
Figures 5, 6, 15M-P, map 8

**HOLOTYPE:** Male: **PAPUA NEW GUINEA: East New Britain:** Baining Mountains, Raunsepna, 4.433°S 151.783°E, 800–1000 m, 17 Nov 1979, W.C. Gagne (AMNH\_PBI 00041550) *Homalanthus* sp. (Euphorbiaceae) (BPBM type# 16732) (BPBM).

**DIAGNOSIS:** Relatively large species distinguished by the following characters: unilobed proepisternum; dark brown coloration; impunctate pronotum; extremely long and thin left paramere. *C. prolixipenis* is similar to *hermosus* in structure and facial coloration, but the male genital capsule and left paramere are clearly distinct. The overall dark coloration resembles that of *nigrus* and *thalame*, but in *nigrus* the head and legs are all black, while *thalame* lacks the dark apical annulation on AII. Only *drepanopenis* has a left paramere as long and thin as *prolixipenis*, but the latter lacks pronotal punctuation and is much darker in coloration.

**DESCRIPTION: COLORATION** (figs. 5, 6): Head: Mostly dark brown above frontoclypeal suture, yellow below; frons without vittae; ocular margins and posterior margin of vertex yellow, with faint brown mark medially; gena, maxillary and mandibular plates yellow; clypeus and buccula light brown; labrum brown; labium light yellow-brown, becoming darker apically. Antenna: AI dark brown; AII yellow-brown, with darker brown annulation; AIII and AIV dark brown. Thorax: Pronotal collar brown, becoming more orange posteriorly; pronotum dark brown, posterior margin faintly orange; thoracic pleura dark brown; mesoscutum and scutellum dark brown. Hemelytra: Dark brown. Legs: Procoxa yellow, forelegs missing from specimen; mesocoxa and trochanter dark brown, basal 2/3 of mesofemora yellow, apical third dark brown, tibia dark brown; metafemur dark brown, without diagonal stripes on outer margin; all tarsi yellow. Abdomen: Uniformly dark brown; left paramere yellow. **SURFACE AND VESTITURE** (figs. 5, 6): Head smooth, glossy and impunctate; pronotum, propleuron, metepimeron, scutellum, and hemelytra impunctate; head and dorsum sparsely clothed in short, white, decumbent setae, most setae knocked off holotype specimen. **STRUCTURE** (figs. 5, 6): Head: Wide; frons only weakly tumescent medially; posterior margin of head nearly flat. Thorax: Pronotum broad, weakly convex, anterior and anterolateral margins slightly upturned, submarginal region of humeral angles weakly excavate, posterior margin thinly carinate, weakly cleft medially; proepisternum uni-

lobed; posterior margin of metepimeron extended in the form of a long, triangular lobe; metanotum not prominent and flared. Hemelytra: Costal margin thin and flared over its entire length and weakly sinuate. Legs: Metatibial spines short and thin. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. MALE GENITALIA (fig. 15M–P): Right paramere triangular (fig. 15O–P); left paramere long, very thin, and curved, slightly twisted along its axis (fig. 15M), extending over the edge of the pygophore (fig. 15N); fold on left of pygophore forming a shallow groove (fig. 15N). FEMALE PARAGENITALIA: Female unknown.

ETYMOLOGY: Derived from Latin *prolix* ("long") and *penis*, the name *prolixipenis* refers to the extremely long left paramere of the male.

HOST: Collected on an unidentified species of *Homalanthus* (Euphorbiaceae).

DISTRIBUTION: Found in the Baining Mountains, East New Britain, Papua New Guinea (map 8).

*Coridromius pteraulos*, new species

Figures 3, 5, 6, 16C–H, 17A–D, map 5

HOLOTYPE: Male: PAPUA NEW GUINEA: Morobe: Wau, 7.3333°S 146.71667°E, 1300 m, 25 Oct 1982–06 Nov 1982, W.C. Gagne (AMNH\_PBI 00041517) (BPBM type# 16733) (BPBM).

DIAGNOSIS: Recognized by the following combination of characters: left posterior margins of abdominal sternites VII and VIII of male with a patch of dense, conelike setae; anterolateral margin of female hemelytra curled to form a semienclosed tube. This species is very similar to *declivipennis*, except that it possesses a thin, fingerlike ventral apical process on the pygophore (rather than the short and thick process of the latter; see fig. 11A), the hemelytra are less flared posteriorly, and the female possesses modified hemelytra. This hemelytral modification probably acts as a guide for the male's left paramere during copulation.

DESCRIPTION: COLORATION (figs. 5, 6): Mostly brown with orange and brown markings of variable intensity. Head: Yellow-brown to orange-brown; vittae on frons

dark brown; depressions behind paired tu- mescences dark brown; ocular margins and posterior margin of head yellow; clypeus with one medial and two lateral brown stripes; gena, mandibular and maxillary plates yellow to pale orange-brown; buccula brown basally, becoming yellow; labrum brown; labium yellow, brown apically. Antenna: AI dark brown, apex yellow; AII light orange-brown with pale subapical annulation and broad brown apical annulation; AIII and AIV dark brown, yellow basally. Thorax: Pronotal collar yellow; pronotum mostly orange- brown with faint yellow medial stripe, yellow margins and faint yellowing at humeral angles; mesoscutum orange-brown; scutellum darker brown with faint yellow medial stripe, apex and basal corners; thoracic pleura orange to orange-brown, posterior margins yellowed. Hemelytra: Yellow-brown to brown, darker anteriorly, yellow to cream colored along claval commissure, lateral margins, and posteriorly; cuneus yellow- brown to brown; membrane brown with dark brown veins. Abdomen: Yellow-brown to orange-brown with a row of dark brown markings running from front to back, dor- solateral margins with alternate brown and cream markings; posterior margin of segment II dark brown. Legs: All coxae pale cream- colored; pro- and mesofemora cream-col- ored, brown apically; metafemur cream to yellow or orange-brown with 9 diagonal brown stripes on outer surface, sometimes dark brown immediately below striping; tibiae yellow, sometimes slightly brown. SURFACE AND VESTITURE (figs. 5, 6, 16E–F): Head with minor swelling on vertex finely punctate; pronotum, propleuron, and metepimeron finely punctate; scutellum, hemelytra, and cuneus irregularly punctate; clothed in long, white, decumbent setae; left posterior margins of abdominal sternites VII and VIII of male with a patch of dense, conelike setae (fig. 16E–F). STRUCTURE (figs. 3, 5, 6, 16C–H, 17A–D): Head: Frons roundly tumescent medially, merging with minor swelling on vertex; vertex with two minor swellings adjacent to eyes; posterior margin of head rounded, weakly carinate. Thorax: Pronotum broad and steep, posteri- orly rounded, lateral margins carinate, anterolateral margins somewhat depressed, not

upturned, submarginal region of humeral angles excavate, collar region not developed; proepisternum bilobed (fig. 3C); posterior margin of metepimeron truncate (fig. 16G); metanotum prominent and flared (fig. 16C–D); scutellum somewhat flattened (fig. 16C–D). Hemelytra: Costal margin straight and carinate, females with anterior costal margin (both right and left sides) strongly carinate and recurved, forming a semicircular tunnel, open posteriorly (figs. 3C, 16H). Legs: Metatibial spines long and thick (fig. 3A). Abdomen: Posterolateral margin of abdominal sternite II sharply angular (figs. 3B, 16G). MALE GENITALIA (figs. 16E, 17A–D): Right paramere broad and rounded with thumblike lobe (fig. 17B–C), paramere appears U-shaped from above (fig. 17D); left paramere thick and elongate, sharply angled basally, then evenly curved and tapering to apex with a slight twist along its axis, curved at tip (fig. 17A); pygophore with a long mesal suture and long, thin, fingerlike ventral apical process (fig. 16E). FEMALE PARAGENITALIA: Abdominal tergite II swollen, posterolateral margins sulcate; right laterotergite II slightly desclerotized and swollen (fig. 16C–D).

**ETYMOLOGY:** The epithet *pteraulos* (Greek, “tube wing”) refers to the modified hemelytra of the female.

**REMARKS:** Based on the modifications of hemelytra and the anterolateral margin of the abdomen of females, it is believed that the site of insemination in this species is dorso-lateral.

**HOST:** Unknown.

**DISTRIBUTION:** Several specimens have been collected in Wau, Papua New Guinea, with a single specimen also collected in Sabah (map 5).

**PARATYPES: PAPUA NEW GUINEA:** **Morobe:** Wau, 7.3333°S 146.71667°E, 1200 m, 13 Dec 1976, W.C. Gagne, Light Trap, 2♂ (AMNH\_PBI 00041523, AMNH\_PBI 00041524) (BPBM); 02 Dec 1976, W.C. Gagne, Light Trap, 1♂ (AMNH\_PBI 00041519), 1♀ (AMNH\_PBI 00041520) (BPBM); 10 Dec 1976, W.C. Gagne, Light Trap, 2♀ (AMNH\_PBI 00041521, AMNH\_PBI 00041522) (BPBM); 15 Aug 1972, G.G.E. Scudder, Light Trap, 1♂ (AMNH\_PBI 00041525) (BPBM); 13 Sep 1972, G.G.E.

Scudder, Light Trap, 1♂ (AMNH\_PBI 00041526) (BPBM); 12 Sep 1972, G.G.E. Scudder, Light Trap, 1♂ (AMNH\_PBI 00041527) (BPBM); 08–14 Dec 1976, G.F. Hevel and R.E. Dietz IV, Light Trap, 1♂ (AMNH\_PBI 00185367) (USNM). Wau, 7.3333°S 146.71667°E, 1300 m, 25 Oct 1982–06 Nov 1982, W.C. Gagne, 1♂ (AMNH\_PBI 00041518) (BPBM).

**OTHER SPECIMENS EXAMINED: MALAYSIA: Sabah:** Tenompok, 6.01777°N 116.48555°E, 637 m, 10–19 Feb 1959, T.C. Maa, 1♂ (BPBM).

*Coridromius punctatus* Carvalho  
Figures 5, 6, 17E–G, 18A–E, map 8

*Coridromius punctatus* Carvalho, 1987: 65–67 (n. sp.).

**DIAGNOSIS:** Distinguished by the following combination of characters: bilobed proepisternum; somewhat rounded lobe at apex of scutellum; mottled coloration; AII dark brown with pale base and apex; AII without dark brown postmedial band; right paramere with sharp, triangular spicule; left paramere scyphelike with gutter open over its entire length. *C. punctatus* is very similar to *chinensis*, *confusus*, and *minusculus* (see diagnosis for *chinensis* above). The apex of the scutellum is less swollen than in *chinensis*, and *punctatus* is the only species with AII mostly dark brown and a short and thick ventral apical process on the pygophore.

**REDESCRIPTION: COLORATION** (figs. 5, 6): Mottled pale yellow, cream, light orange-brown and dark brown. Head: Mottled pale yellow with light orange-brown markings; faint light brown vittae sometimes present; depressions on vertex dark brown; mandibular plate and clypeus cream to yellow; clypeus with two irregular dark brown stripes, brown apically; maxillary plate orange-brown; buccula and gena cream to yellow; labrum orange-brown, labium light orange-brown, brown apically. Antenna: AI dark brown with yellow base and apex; AII light orange-brown, apically with faint cream annulation followed by dark brown annulation; AIII and AIV dark brown with yellow bases. Thorax: Pronotal collar yellow to light orange-brown; pronotum cream with mottled orange-brown punctures, darker around collar region, with two small dark brown

markings along anterior margin behind eyes; cream-colored medial line, often flanked with two irregular broad, dark brown patches; mesoscutum orange-brown; scutellum cream with mottled orange-brown punctures, apex cream, sometimes with pale midline. Hemelytra: Mostly orange-brown, sometimes with slightly darker brown patches; embolium sometimes with faint to more pronounced broad cream to yellow spots subbasally and anterior to costal fracture, sometimes with red tinge at costal fracture; endocorium with darker brown and cream patterning at apex of claval commissure; cuneus orange-brown, sometimes tinted red; membrane brown, with pale veins bounded by irregular darker brown markings. Legs: Mostly cream to yellow with brown markings; coxae cream; pro- and mesofemora with broad, subapical irregular brown markings; metafemur laterally with eight irregular diagonal brown stripes, ventrally with dark brown patch below stripes, depressions near apex orange-brown, bothria cream; pro- and mesotibiae pale with irregular dark brown markings; metatibia cream to yellow with irregular dark brown markings, sometimes fused; metatibial spines dark brown. Abdomen: Light orange-brown, sometimes with darker brown markings, posterior margin of sternite II dark brown. SURFACE AND VESTITURE (figs. 5, 6, 18A–C): Irregular depressions on head with deep punctures, somewhat coalescing (fig. 18B); pronotum and propleuron (fig. 18A) deeply, irregularly punctate; scutellum deeply, irregularly rugulopunctate, less so along midline; metepimeron (fig. 18C) and hemelytra impunctate; head and dorsum clothed in long white setae. STRUCTURE: (figs. 5, 6, 18A–C): Head (fig. 18A–B): Highly contoured; frons strongly tumescent, bounded by irregular depressions and three minor swellings laterally and posteriorly; foveae adjacent to eyes and near posterior margin of vertex deep; posterior margin of head medially rounded, not carinate; gena swollen; mandibular plate tumescent (fig. 18A). Thorax: Pronotum broad and somewhat elongate, posteriorly rounded, steeply declivous, deeply, irregularly punctate, anterolateral margins weakly carinate, submarginal region of humeral angles strongly excavate; posterior margin weakly cleft me-

dially; collar region weakly developed; proepisternum bilobed (fig. 18A); propleuron deeply, irregularly punctate (fig. 18A); posterior margin of metepimeron either truncate (fig. 18C) or with a small rounded lobe; metanotum prominent and flared; scutellum dorsally flattened, apex a weakly rounded lobe. Hemelytra: Costal margin evenly rounded over its entire length, anterior portion not expanded. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. MALE GENITALIA (figs. 17E–G, 18D–E): Right paramere rounded with a long, sharp, triangular process (figs. 17F–G, 18D–E); left paramere scythe-shaped, slightly twisted along its axis, thick and laterally compressed at base, becoming thin apically, gutter open over entire length (figs. 17E, 18E); posterior margin of pygophore with long mesal suture, short fingerlike ventral apical process present (fig. 18D). FEMALE PARAGENITALIA: No visible external paragenitalia.

HOST: One specimen has been collected from *Rubus* sp. (Rosaceae).

DISTRIBUTION: Found in Papua New Guinea (map 8).

HOLOTYPE: PAPUA NEW GUINEA: Morobe: Mount Kaindi, 7.35°S 146.68333°E, 2300 m, 31 Dec 1964, J. Sedlacek, 1♀ (AMNH\_PBI 00042153) (BPBM).

OTHER SPECIMENS EXAMINED: PAPUA NEW GUINEA: East New Britain: Baining Mountains, Raunsepna, 4.666°S 152°E, 800–1000 m, 17 Nov 1979, W.C. Gagne, 1♂ (AMNH\_PBI 00041455) (BPBM). Morobe: Huon Peninsula: Finisterre Range, 5.8333°S 146.1166°E, 2550 m, 01–21 Oct 1958, W.W. Brandt, 1♂ (AMNH\_PBI 00041462) (BPBM). Mount Kaindi, 7.35°S 146.6833°E, 2360 m, 03 Dec 1977, W.C. Gagne, Light Trap, 1♂ (AMNH\_PBI 00041450), 3♀ (AMNH\_PBI 00041451–AMNH\_PBI 00041453) (BPBM); 05 Dec 1976, W.C. Gagne, *Rubus* sp. (Rosaceae), 1♀ (AMNH\_PBI 00041456) (BPBM); 24 Feb 1979, W.C. Gagne, Light Trap, 1♀ (AMNH\_PBI 00041461) (BPBM). Mount Kaindi, 7.35°S 146.68333°E, 2350 m, 14 Apr 1971, Tawi, 1♀ (AMNH\_PBI 00041454) (BPBM); 23–28 Jul 1970, T. Tigner, 1♂ (AMNH\_PBI 00041459) (BPBM); 27 Nov 1976, W.C. Gagne, 1♀ (AMNH\_PBI 00041460) (BPBM).




Fig. 17. Left and right parameres of *Coridromius* spp.: A–D: *C. pteraulos* (arrow indicates aedeagus). E–G: *C. punctatus*. H–K: *C. sommelieri*, L–O: *C. tahitiensis*. P–R: *C. variegatus*. S–U: *C. zetteli*.

Mount Kaindi, top, 7.35°S 146.68333°E, 24 Apr 1970, A.N. Gillogly & S. Keenan, Light Trap, 1♂ (AMNH\_PBI 00041458), 1♀ (AMNH\_PBI 00041457) (BPBM).

*Coridromius ruwenzorii*, new species  
Figures 5, 6, 18F–H, map 2

HOLOTYPE: Female: **UGANDA:** Ruwenzori Range, Misigo, 0.24719°N 29.82239°E, 2606 m, 02–03 Jul 1952, D.S. Fletcher (AMNH\_PBI 00178010) (BMNH).

DIAGNOSIS: Identified by the following combination of characters: smooth and rounded head; punctate hemelytra; brownish orange coloration with two large dark brown spots on scutellum; unique female paragenitalia. The male of *ruwenzorii* is unknown. Known only from a single female specimen, this species is nearly identical to *bulbopella*, however, the paragenitalia lack the well-developed cup-and-bulb structure of the latter, appearing instead as a much-reduced version thereof, with an additional desclerotized swelling of the first right dorsal laterotergite.

DESCRIPTION: COLORATION (figs. 5, 6) Head: Concolorous orange-brown; vertex without vittae; apex of clypeus slightly brown; maxillary plate brown. Antenna: Mostly yellow; AI with broad brown apical annulation; AII with brown apical annulation; AIII and AIV brown. Thorax: Pronotal collar yellow; pronotum orange-brown with very faint yellow medial stripe, posterior tinged orange; mesoscutum tinted orange; scutellum orange-brown with two large brown markings; thoracic pleura brownish yellow. Hemelytra: Mostly orange-brown, tinted orange at base; claval commissure faintly brown; membrane pale yellow-brown with darker brown veins. Abdomen: Yellow-brown, more orange along dorsal margin, with faint red markings laterally. Legs: Yellow-brown, paler basally, pro- and mesofemora with brown subapical band; metafemur with brown transverse stripes, apically tinted orange. SURFACE AND VESTITURE (figs. 5, 6, 18F–H): Head smooth with a few shallow punctures on vertex, vertex coarsely punctate; pronotum, scutellum (fig. 18F–G), and hemelytra with large, evenly spaced punctures; propleuron, met-

epimeron, and cuneus impunctate; clothed in short, white, decumbent setae. STRUCTURE (figs. 5, 6, 18F–H) Head: Frons with broad medial tumescence contiguous with raised lateral tubercles adjacent to eyes; posterior margin of head distinctly rounded. Thorax: Pronotum broad and evenly rounded, margins thinly carinate, anterolateral margins adjacent to collar region depressed, submarginal region of humeral angles weakly excavated, posterior margin weakly upturned, collar region weakly developed (fig. 18F); proepisternum bilobed; posterior margin of metepimeron broadly rounded (fig. 18H); metanotum prominent and flared; apex of scutellum not a swollen lobe. Hemelytra: Costal margin weakly flared posteriorly, carinate; cuneus small. Legs: Metatibial spines long and thick. Abdomen: Posterolateral margin of abdominal sternite II rounded, not sharply angular (fig. 18H). MALE GENITALIA: Unknown. FEMALE PARAGENITALIA (fig. 18F–G): Inner margins of right dorsal laterotergites II and III weakly lobate, laterotergite I swollen and desclerotized, posterior margin of right abdominal sternite II carinate and slightly flared, indicating probable sight of copulation.

ETYMOLOGY: Named for the Ruwenzori range in Uganda, from which the only specimen has been collected.

HOST: Unknown.

DISTRIBUTION: Found in Uganda (map 2).


*Coridromius schuhi*

*Coridromius schuhi* Linnauvori, 1994: 15–17 (n. sp.).

DISCUSSION: We have not seen any specimens of this species, as it appears that they have been lost. The original description and limited illustrations are not sufficient to distinguish this species from most others within the genus. Until such a time as the specimens are recovered, little can be said about the relationship of this species to others in the genus.

*Coridromius sommelieri*, new species  
Figures 5, 6, 17H–K, 19A–F, 20B, map 5

HOLOTYPE: Female: **MALAYSIA: Sabah:** Tenompok, 6.01777°N 116.48555°E, 637 m,



Sue Lindsay, Australian Museum

Fig. 18. Scanning electron micrographs: **A–E:** *C. punctatus*: **A.** Head and thorax, lateral view. **B.** Head, anterior view. **C.** Thorax and abdomen, lateral view. **D.** ♂ pygophore, posteroventral view. **E.** ♂ genitalia, posterior view. **F–H:** *C. ruwenzori*: **F.** ♀ right wing and hemelytron removed, dorsal view. **G.** ♀ thorax and abdomen, arrow indicates female paragenital structure on laterotergites II and III, dorsolateral view. **H.** lateral view. LP = left paramere, MES = mesepimeron, MET = metepimeron P = peritreme, PES = proepisternum, RP = right paramere, VAP = ventral apical process.

15 Feb 1959, T.C. Maa (AMNH\_PBI 00041567) (BPBM type# 16734) (BPBM).

**DIAGNOSIS:** This species is recognized by the following combination of characters: bilobed proepisternum; embolium with yellow U-shaped fascia near cuneal fracture; outer margin of embolium straight and carinate; male left paramere tightly corkscrewed with apex perpendicular to axis; female with corkscrewed paragenital opening on the right side of the abdomen in sternite II. This species is very similar to *zetteli*, but the ventral apical process on the pygophore is much more pronounced in *sommelieri* and the apex of the left paramere is perpendicular to its axis, rather than almost parallel as in *zetteli*. The female of *zetteli* is unknown.

**DESCRIPTION: COLORATION** (figs 5, 6): Yellow-brown with yellow and dark brown markings. Head: Light orange-brown; frons sometimes with pale brown vittae, faintly yellow along midline; ocular margins, raised lateral tubercles and posterior margin of head yellow; clypeus dark orange-brown, darkest at apex; maxillary plates tinted orange; labrum dark brown; labium light orange-brown, dark brown at apex. Antenna: AI dark brown; AII light yellow-brown with dark brown subapical band separated from dark brown apical annulation by narrow cream-colored band, base somewhat darkened; AIII and AIV dark brown, pale at base. Thorax: Pronotal collar yellow to brownish yellow; pronotum light orange-brown with two large dark brown markings reaching from anterior margin almost to posterior margin, midline pale, lateral and posterior margins yellow; mesoscutum orange-brown; scutellum light to dark brown with yellow midline, apex and sides; thoracic pleura mostly yellow-brown to orange-brown with posterior margins of each sclerite yellowed; mesopleuron dark brown ventrally. Hemelytra: Predominantly light orange-brown, with darker brown markings on corium and embolium above cuneal fracture; costal margins yellow; U-shaped fascia on lateroapical margin of embolium yellow to orange; cuneus dark brown; membrane light brown with dark brown veins. Legs: Pro- and mesofemora and tibiae light yellow-brown, sometimes tinted orange; metafemur yellowed basally, with 8 or 9 diagonal brown

stripes merging with dark brown patch below, apex brown; metatibia orange-brown to dark brown. Abdomen: Orange-brown with alternating cream and dark brown markings along dorsolateral margin. **SURFACE AND VESTITURE** (figs. 5, 6, 19A–B): Head, pronotum, propleuron, metepimeron (fig. 19A–B), scutellum, and hemelytra finely shallowly punctate; head and dorsum clothed in long, silvery, decumbent setae. **STRUCTURE** (figs. 5, 6, 19A–B): Head: Frons broadly tumescent medially; lateral tubercles adjacent to eyes small, bordered posteriorly by shallow foveae; posterior margin of head carinate and medially rounded. Thorax: Pronotum broad, margins slightly carinate, anterolateral margins slightly turned upward, submarginal region of humeral angles excavated, collar region undifferentiated; proepisternum bilobed; posterior margin of metepimeron truncate (fig. 19A–B); metanotum prominent and flared; scutellum flattened. Hemelytra: Costal margin straight and carinate; embolium with U-shaped fascia on lateroapical angle. Legs: Metatibial spines long and thick. Abdomen: Posterolateral margin of abdominal sternite II angular (fig. 19A). **MALE GENITALIA** (figs. 17H–K, 19D–F): Right paramere broad and rounded with thumblike lobe, appearing U-shaped when viewed dorsally (figs. 17I–K, 19E); left paramere straight and tightly coiled along its axis, apex thin and angled perpendicular to axis (figs. 17H, 19E–F); posterior margin of pygophore with broad, triangular ventral apical process and long mesal suture (fig. 19D). **FEMALE PARAGENITALIA** (figs. 19B–C, 20B): On right side of body, corkscrewed copulatory canal located intrasegmentally in sternite II (figs. 19B–C, 20B).

**ETYMOLOGY:** The name *sommelieri* refers to the shape of the male left paramere, reminiscent of a sommelier's corkscrew.

**REMARKS:** *C. sommelieri* is one of the most unusual species of *Coridromius* due to the corkscrewed male left paramere and reciprocal corkscrewed paragenital opening on the female. When dissected, this female copulatory tube is shown to be thinly sclerotized, at the apex of which one finds the puncture wound caused by the male genitalia (fig. 20B). Specimens of *sommelieri*

were initially thought to belong to *zetteli* (e.g., Tatarnic et al., 2006), but closer examination of the male genitalia has revealed that they are in fact separate species. This species also appears similar to *testaceous*, except females of the latter lack any visible external paragenitalia. The male of *testaceous* is unknown.

HOST: Unknown.

DISTRIBUTION: Collected from Tenompok, Sabah (map 5).

PARATYPES: MALAYSIA: Sabah: Tenompok, 6.01777°N 116.48555°E, 637 m, 10–19 Feb 1959, T. C. Maa, 1♂ (AMNH\_PBI 00041571), 1♀ (AMNH\_PBI 00041570) (BPBM); 13 Feb 1959, T.C. Maa, 1♀ (AMNH\_PBI 00041572) (BPBM); 15 Feb 1959, T.C. Maa, 2♀ (AMNH\_PBI 00041568, AMNH\_PBI 00041569) (BPBM).

#### *Coridromius tahitiensis*, new species

Figures 5, 6, 17L–O, 19G–H, 20C, map 1

HOLOTYPE: Male: FRENCH POLYNESIA: Society Islands: Tahiti Island: road to Mt. Marau, 17.61666°S 149.39472°W, 840–850 m, 29 Aug 1977, W.C. Gagne and S.L. Montgomery, Light Trap (AMNH\_PBI 00041530) (BPBM type# 16735) (BPBM).

DIAGNOSIS: This species can be recognized by the following characters: distinct dark brown Y-shaped pattern along inner margins of hemelytra; metafemoral stripes faint or absent; females with a large paragenital opening on the right side of the abdomen between sternites II and III.

DESCRIPTION: COLORATION (figs. 5, 6): Head: Without vittae; pale yellow-brown to brown; labium pale, brown apically. Antenna: AI dark brown; AII pale yellow-brown with dark brown apical annulation; AIII and AIV uniformly dark brown. Thorax: Pronotal collar uniformly yellow; pronotum pale yellowish brown to brown, with a pale medial stripe, darker specimens with a pair of small, dark brown maculations anteriorly, occasionally with broad dark brown rectilinear maculations posteriorly, margins with distinct yellow piping; mesoscutum anteriorly black, caudally brown; scutellum yellow in males, brown with darker brown and yellow markings in females; thoracic pleura mostly brown with margins of each sclerite yellowed.

Hemelytra: Pale yellowish brown to darker brown with distinct dark brown “Y” along inner margins, sometimes with dark brown markings on corium and posterior margin of embolium; cuneus yellow-brown; membrane yellow-brown, veins darker brown anteriorly. Abdomen: Pale yellow, sometimes with brown, orange or red markings; females often with broad, brown markings anterolaterally. Legs: Pro- and mesofemora uniformly pale yellow; metafemur either pale yellow with brown apex (most males) or pale yellow to brown with 7 diagonal brown bands coalescing ventrally into a large brown patch, becoming reddened posterior to striping (females), brown at apex. SURFACE AND VESTITURE (figs. 5, 6, 19G): Head shallowly punctate; pronotum finely punctate; propleuron, metepimeron (fig. 19G), scutellum and hemelytra impunctate; head and dorsum sparsely clothed in short, pale, decumbent setae. STRUCTURE (figs. 5, 6, 19G): Head: Frons weakly tumescent medially; vertex with slightly raised tubercles adjacent to eyes, bordered posteriorly by shallow depressions; posterior margin of head carinate and almost flat. Thorax: Pronotum weakly rounded, finely punctate, broad, anterolateral margins weakly upturned and carinate, submarginal region of humeral angles weakly excavate, posterior margin weakly upturned; proepisternum unilobed; posterior margin of metepimeron truncate (fig. 19G); metanotum not prominently flared. Hemelytra: Costal margin flared over its entire length and weakly sinuate, more so in males. Legs: Metafemur only weakly incrassate; metatibial spines short and thin. Abdomen: In lateral view posterior margin of abdominal sternite II not distinctly angular. MALE GENITALIA (figs. 17L–O, 19H): Right paramere triangular (fig. 17M–O); left paramere thick and scythe-shaped (figs. 17L, 19H); pygophore with a deep, rounded groove on left side of ventral margin, no mesal suture (fig. 19H). FEMALE PARAGENITALIA (fig. 19G, 20C): Right posterolateral margin of abdominal segment II flared outward, distinct paragenital opening between segments II and III (fig. 19G) opening into a thinly sclerotized chamber within the abdominal cavity (fig. 20C).




Fig. 19. Scanning electron micrographs: A–F: *C. sommelieri*: A. ♀ thorax and abdomen, left lateral view. B. ♀ thorax and abdomen, arrow indicates paragenital opening, right lateral view. C. ♀ detail of paragenital opening. D. ♂ pygophore, ventral view. E. ♂ genitalia, dorsal view. F. ♂ detail of left paramere. G–H: *C. tahitiensis*: G. ♀ thorax and abdomen, arrow indicates paragenital opening, right lateral view. H. ♂ genitalia, posterior view. LP = left paramere, MES = mesepimeron, MET = metepimeron, MLS = mesal suture, P = peritreme, RP = right paramere, VAP = ventral apical process.

Sue Lindsay, Australian Museum

**ETYMOLOGY:** Named after Tahiti Island, where the holotype was collected.

**REMARKS:** This is the only known oceanic species with prominent external paragenitalia. It is also one of the few species we have seen with clear sexually dimorphic coloration.

**HOST:** Unknown.

**DISTRIBUTION:** Found in French Polynesia (map 1).

**PARATYPES:** **FRENCH POLYNESIA: Society Islands:** Moorea Island: Mt. Tohiea base: Belvedere Trail, 17.556°S 149.826°W, 400 m, 18–19 Sep 1988, S.L. Montgomery, 1♀ (AMNH\_PBI 00041545) (BPBM). Tahiti Island: Fare Rau Ape Aorai, 17.61916°S 149.49027°W, 600–1400 m, 09 Mar 1977, N.L.H. Krauss, 1♂ (AMNH\_PBI 00041536) (BPBM). Fare Vaiufaufa, 17.7825°S 149.25166°W, 850 m, 03 Jul 1977, W.C. Gagne, Light Trap, 3♂ (AMNH\_PBI 00041533–AMNH\_PBI 00041535), 3♀ (AMNH\_PBI 00041539–AMNH\_PBI 00041541) (BPBM). Mt. Marau, 17.61805°S 149.38333°E, 1400–1500 m, 18–21 Aug 1977, W.C. Gagne and S.L. Montgomery, Light Trap, 2♀ (AMNH\_PBI 00041531, AMNH\_PBI 00041532) (BPBM); 14 Sep 1988, S.L. Montgomery and B.H. Gagne, 2♂ (AMNH\_PBI 00041537–AMNH\_PBI 00041538), 3♀ (AMNH\_PBI 00041542–AMNH\_PBI 00041544) (BPBM). Mt. Marau, summit road, fern banks nr. km 7, 17.61888°S 149.55444°E, 1125 m, 04 Sep 2006, D. A. Polhemus, 1♀ (AMNH\_PBI 00042604) (BPBM). Papenoo Valley, 17.51°S 149.42°W, 250 m, 17–18 Sep 1988, S.L. Montgomery, Light Trap, 1♀ (AMNH\_PBI 00041546) (BPBM). Tahiti Nui, Mt. Marau, gulch off road to summit, 17.6°S 149.56416°E, 885 m, 10 Sep 2006, C. Ewing, *Coprosoma* sp. (Rubiaceae), 1♂ (AMNH\_PBI 00042603) (BPBM).

*Coridromius testaceous* Liu and Zhao  
Figures 2D, 5, 6, map 5

*Coridromius testaceous* Liu and Zhao, 1999: 55–58.

**DIAGNOSIS:** This species is recognized by the following combination of characters: bilobed proepisternum; embolium with yellow U-shaped fascia near cuneal fracture, lateral margins of hemelytra straight and carinate; female without prominent paragenital opening on right side of abdomen. Male unknown. *C. testaceous* is very similar in

appearance to *sommelieri* and *zetteli*; however, it lacks the distinct paragenitalia of the former, which we suspect may also be present in the latter.

**REDESCRIPTION:** **COLORATION** (figs. 2D, 5, 6): Varying from light brown to almost uniform dark brown with yellow markings. Head: Light brownish yellow to orange-brown; frons typically yellow along midline, with very faint to distinct dark brown vittae; clypeus yellowed with darker brown markings at base, dark brown apically; mandibular plates yellowed; maxillary plates brownish orange; buccula yellowed; labrum dark brown basally, labium pale, brown apically. Antenna: A1 dark brown, pale at base and apex; AII light orange-brown with dark brown subapical band, separated from dark brown apical annulation by narrow cream band; AIII and AIV dark brown, pale at base. Thorax: Pronotal collar yellow to orange-brown, darker anteriorly; pronotum light orange-brown to dark brown, with median yellow band (faint in some specimens), yellowed at humeral angles in paler specimens, lateral and posterior margins yellow; mesoscutum orange-brown to dark brown; scutellum light orange-brown to dark brown with yellow midline, apex and sides; thoracic pleura mostly orange-brown with posterior margins of each sclerite yellowed; mesopleuron dark brown ventrally. Hemelytra: Orange-brown to nearly uniform dark brown; in paler specimens clavus slightly paler at claval commissure and along posterior margin, immediately above cuneal fracture; embolium yellow along outer margin, with U-shaped yellow fascia at lateroapical margin; cuneus dark brown; membrane pale brown with dark brown veins. Legs: Pro- and mesofemora yellow at base, becoming orange-brown toward apex; metafemur pale basally, with 8 or 9 diagonal dark brown stripes, stripes merging with dark brown ventral patch in darker specimens, subapical lateral depressions dark brown. Abdomen: Orange-brown with alternating cream and dark brown markings along dorsolateral margin. **SURFACE AND VESTITURE** (figs. 2D, 5, 6): Head, pronotum, propleuron, metepimeron, scutellum, and hemelytra finely, shallowly punctate; head and dorsum clothed in long, silvery, decumbent setae.

**STRUCTURE** (figs. 2D, 5, 6): Head: Frons broadly medially tumescent and laterally depressed; lateral tubercles adjacent to eyes small, bordered posteriorly by shallow depressions; posterior margin of head carinate and medially rounded. Thorax: Pronotum broad, lateral and posterior margins slightly carinate, anterolateral margins slightly upturned, submarginal region of humeral angles excavated, collar region undifferentiated; proepisternum bilobed; posterior margin of metepimeron truncate; metanotum prominent and flared; scutellum flattened. Hemelytra: Costal margin straight and carinate; embolium with U-shaped fascia on lateroapical angle. Legs: Metatibial spines long and thick. Abdomen: Posterolateral margin of abdominal sternite II angular and transversely creased. **MALE GENITALIA:** Unknown. **FEMALE PARAGENITALIA:** Without distinct external paragenitalia.

**REMARKS:** *C. testaceus* is known only from female specimens. Superficially it appears identical to *zetteli*, which is known only from male specimens and has an overlapping distribution. Furthermore, both species have even been collected together, along with *chinensis*, on a single unidentified *Macaranga* tree in southern China. We refrain from synonymizing the two species without further data, however, as we suspect the female of *zetteli* should have matching corkscrewed paragenitalia to accommodate the unusual male intromittent organ (as we see in *sommelieri*), whereas female *testaceus* have no visible external paragenitalia whatsoever. Interestingly, some of the females collected alongside male *zetteli* show damage in the form of irregular depressions on right abdominal sternite II, perhaps from copulation attempts by male *zetteli*.

**HOST:** Recorded from the flowers of *Macaranga* sp. (Euphorbiaceae) in southern China.

**DISTRIBUTION:** Ranging from central China south through mainland Southeast Asia, west to Nepal (map 5).

**HOLOTYPE:** Female: **CHINA: Hainan Prov.:** Jianfengling, 18.7°N 108.8°E, 01 Apr 1985, Le-Yi Zheng, Light Trap (AMNH\_PBI 00178049) (NKMU).

**OTHER SPECIMENS EXAMINED:** **CHINA: Yunnan Prov.:** Xishuangbanna Tropical




Fig. 20. Female genitalia. **A.** Illustration of ♀ *C. variegatus*, showing complete desclerotization of posterior wall. **B.** Dissection of *C. sommelieri* ♀, showing invagination of corkscrewed paragenital opening (arrow). Inset: Detail showing puncture at apex of paragenital opening (arrow). **C.** Dissection of *C. tahitiensis* ♀. Arrow indicates lightly sclerotized invagination of copulatory opening. *Nikolai Tatarnic*

Rainforest Ecosystem Station, ~3 km from entrance, 21.96777°N 101.2075°E, 650 m, 30 May 2006, N. Tatarnic, *Macaranga* sp. (Euphorbiaceae), 3♀ (AMNH\_PBI 00006070–AMNH\_PBI 00006072) (AMNH). Xishuangbanna Tropical Botanical Garden, gallery forest by river, 21.93333°N 101.24444°E, 553 m, 26 May 2006–28 May 2006, N. Tatarnic, *Macaranga* sp. (Euphorbiaceae), 2♀ (AMNH\_PBI 00006068, AMNH\_PBI 00006069) (AMNH). **INDONE-**

**SIA: North Sumatra:** Medan, 3.5833°N 98.6667°E, 29 Apr 1922, L. Fulmek, Light Trap, 1 undetermined (AMNH\_PBI 00189970) (NHMW). **LAO PEOPLE'S DEMOCRATIC REPUBLIC: Vientiane Prov.:** Ban Van Eue, 17.96666°N 102.6°E, 800 m, 16 Apr 1965, J.L. Gressitt, (Cyperaceae), 2♀ (AMNH\_PBI 00042148, AMNH\_PBI 00042149) (BPBM). **NEPAL:** Royal Chitwan National Park, Machan Resort, 27.58305°N 84.5°E, 07–09 Sep 2005, T. Yasunaga, M. Takai and T. Shishido, 2♀ (AMNH\_PBI 00189998) (TYCN). **VIETNAM:** Tam Dao NP, 21.4536°N 105.6436°E, 900 m, 17–18 Jun 1999, Y. Nakatani, 1♀ (AMNH\_PBI 00190012) (TYCN).

*Coridromius thalame*, new species

Figures 5, 6, 21A, map 5

**HOLOTYPE:** Female: **MALAYSIA: Sabah:** Ranau, 5.9667°N 116.6833°E, 1176 m, 22–25 Feb 1959, T.C. Maa (AMNH\_PBI 00041547) (BPBM type# 16736) (BPBM).

**DIAGNOSIS:** Recognized by the following combination of characters: large size; dark brown coloration with yellow and orange head, venter, and legs; relatively short metatibial spines; unique external paragenitalia. The paragenitalia of *thalame* is similar to that of *ephippius*, except the upper posterior margin of right abdominal sternite II is much less tuberculate. The male of *thalame* is unknown.

**DESCRIPTION: COLORATION** (figs. 5, 6): Head: Mostly orange; lateral tubercles adjacent to eyes yellowed, foveae behind tubercles brown; ocular margins and posterior margin of vertex with yellow piping; frons with very weak brown vittae. Antenna: AI orange, brown apically; AII orange-brown with darker brown apical annulation; AIII and AIV missing from specimen. Thorax: Pronotal collar yellow; pronotum dark reddish brown with faint, lighter brown midline; margins yellow-brown; mesoscutum dark reddish brown; scutellum dark reddish brown, yellowed on anterior corners and on either side of apex; proepisternal lobe yellow, other thoracic pleura dark brown dorsally, yellow ventrally, posterior margins somewhat lighter. Hemelytra: Uniform amber brown; membrane brown with dark brown veins.

Abdomen: Amber brown, becoming slightly paler ventrally and at apex. Legs: Coxae yellow; pro- and midlegs yellow-orange; posterior femur yellow basally, brown ventrally and subapically, with diagonal brown stripes on outer surface, apex yellow-orange; hind tibia yellow-orange. **STRUCTURE** (figs. 5, 6, 21A): Head: Wide; frons weakly medially convex; small raised tubercles adjacent to eyes, bordered posteriorly by shallow depressions; posterior margin of vertex only weakly medially rounded, mostly flat, not carinate. Thorax: Pronotum broad; anterolateral margins slightly upturned; callar region undifferentiated; proepisternum unilobed; posterior margin of metepimeron truncate (fig. 21A); metanotum not prominently flared. Hemelytra: Lateral margin of hemelytron flared posteriorly. Legs: Metatibial spines short and thin. Abdomen: Posterior margin of abdominal sternite II not sharply angled. **MALE GENITALIA:** Unknown. **FEMALE PARAGENITALIA:** On right side of abdomen, laterotergites II and III twisted and swollen, posterior margin of sternite II flared to expose distinct paragenital opening between segments II and III (fig. 21A).

**ETYMOLOGY:** The epithet *thalame* (from Greek, for “cave” or “den”) reflects the unusual paragenital opening of this species.

**REMARKS:** *C. thalame* is one of six species collected by T. C. Maa in February 1959 in Sabah. As with several other species, *thalame* is known from only a single specimen.

**HOST:** Unknown.

**DISTRIBUTION:** Collected in Ranau, Sabah (map 5).

*Coridromius variegatus* (Montrouzier, 1861)

Figures 5, 6, 17P–R, 21B–C, map 1

*Ocypus variegatus* Montrouzier, 1861: 67, (n. sp.). *Leptomerocoris variegatus*: Walker, 1873: 145 (new combination).

**Coridromius variegatus:** Poppius, 1911: 15; (Van Duzee), 1932: 180; (Woodward), 1954: 231; (Carvalho), 1955: 68; (Carvalho), 1987: 67–69.


**DIAGNOSIS:** Recognized by the following combination of characters: proepisternum unilobed; metanotum not prominently flared; upper posterior margin of metepimeron with small, rounded lobe; posterior margin of pygophore with fold on left side forming a shallow U-shaped groove, without small apophysis on right margin of groove; left paramere short and thick; right paramere triangular and apically elongate. *C. variegatus* is most similar to *bicolor*, *chenopoderis*, *monotocopsis*, and *pilbarensis*, but can be distinguished from these by minor variations in coloration and male genitalia.

**REDESCRIPTION: COLORATION** (figs. 5, 6): Mostly yellow- to orange-brown, varying in intensity. Head: Yellowish brown; with dark brown vittae; lateral margins bordering eyes and posterior margin of vertex yellow; lateral tubercles yellow, depressions brown to black; vertex with median brown to black marking; mandibular and maxillary plates yellow-brown; buccula darker brown; clypeus with irregular median and lateral reddish brown markings; labrum reddish brown; labium pale brown, dark brown at apex. Antenna: AI pale, sometimes with faint subapical brown band; AII without submedial band, apex with dark brown annulation; AIII and AIV dark brown. Thorax: Pronotal collar yellow to orange-brown, anterior margin dark brown; pronotum mostly orange-brown with faint yellow medial stripe, collar region with irregular dark brown spots coalescing, margins of pronotum with yellow piping; mesoscutum orange-brown; scutellum orange-brown, with paired lateral dark brown markings, corners fading to pale yellow, sometimes with a median dark brown stripe; thoracic pleura mostly dark brown with posterior margins of each sclerite yellowed. Hemelytra: Varying from light yellow-brown to dark brown, paler apically; claval commissure dark brown; endocorium with dark brown arched band coalescing with claval commissure and dark brown spot below; embolium sometimes tinted red or orange at cuneal fracture; cuneus generally yellow-brown, sometimes tinted red or orange; membrane brown with dark brown veins. Legs: Pro- and mesofemora pale yellow-brown; metafemur, pale basally, with 8 or 9 diagonal dark brown stripes on lateral

surface, ventrally brown, somewhat chestnut laterally behind stripes, bothria not readily visible; pro- and mesotibiae pale; hind tibia brown; tarsi brown at apex. Abdomen: Yellow- to orange-brown, often dark brown anteriorly, dorsolateral half of sternites 7–8 yellow in some individuals. **SURFACE AND VESTITURE** (figs. 5, 6): Head and pronotum with shallow punctures; propleuron, metepimeron, scutellum, and hemelytra impunctate; head and dorsum sparsely covered in short, pale, decumbent setae. **STRUCTURE** (figs. 5, 6): Head: Frons broadly tumescent medially; vertex with two slightly raised tubercles adjacent to eyes, bordered posteriorly by shallow depressions; posterior margin of head somewhat carinate, slightly rounded medially but otherwise flat. Thorax: Pronotum broad, margins carinate, antero-lateral margins only weakly upturned, submarginal region of humeral angles weakly excavate; proepisternum unilobed; posterior margin of metepimeron truncate with distinct rounded lobe projecting rearward from upper corner (as in *marmoreus*: fig. 14B); metanotum not prominent and flared. Hemelytra: Lateral margin of hemelytron thin and slightly flared over its entire length. Legs: Metatibial spines long and thick. Abdomen: When viewed laterally, posterior margin of abdominal sternite II not distinctly angular. **MALE GENITALIA** (figs. 17P–R, 21B–C): Right paramere small, triangular, and apically elongate (fig. 17Q–R); left paramere short and scythe-shaped, apex broadly acute (figs. 17P, 21B–C); pygophore biconvex with shallow fold on left ventral margin (fig. 21B–C). **FEMALE PARAGENITALIA:** No visible paragenitalia, but right laterotergite II with small melanized scars in some individuals, indicating probable site of copulation.

**REMARKS:** This is the type species of *Coridromius*. Contrary to Carvalho's re-description (Carvalho, 1987: 67–69), there is no spinelike projection on the pygophore. It is most likely that what has been misinterpreted as a process on the pygophore is in fact the apex of the right paramere.

Up until now most *Coridromius* specimens from the Pacific and Australia have been thought to belong to *variegatus*. Instead, what we have thought of as *variegatus* is in effect four distinct species. Described from



Sue Lindsay, Australian Museum

Fig. 21. Scanning electron micrographs: **A.** *C. thalame*: ♀ thorax and abdomen, arrow indicates paragenital opening, right lateral view. **B–C:** *C. variegatus*: **B.** ♂ genitalia, dorsal view. **C.** ♂ genitalia, posterior view. **D.** *C. zetteli*: ♂ genitalia, posterior view. LP = left paramere, MES = mesepimeron, MET = metepimeron, P = peritreme, RP = right paramere.

New Caledonia, *variegatus* is also found in Fiji, and perhaps other nearby islands. In Australia and New Zealand, however, what we find is not *variegatus* but three other species: *chenopoderis*, *monotocopsis*, and *pilbarensis* (see descriptions above). Superficially all four species are strikingly similar, yet closer examination yields several distinguishing, if subtle, features. *C. variegatus* can readily be distinguished from the others by the small but distinct rounded lobe on the upper posterior margin of the metepimeron (as in *marmoreus*; fig. 14B), and males of *variegatus* by the relatively longer apex of the right paramere.

**HOST:** Specimens have been collected from unidentified species of *Phyllanthus* and *Macaranga* (Euphorbiaceae) (table 2).

**DISTRIBUTION:** *C. variegatus* is known from New Caledonia and Fiji (map 1).

**HOLOTYPE: NEW CALEDONIA: Province des Iles Loyauté:** Lifou Island,

20.88333°S 167.2167°E, 1861, P.A. Montrouzier, 1♀ (AMNH\_PBI 00018973) (NHMW).

**OTHER SPECIMENS EXAMINED: FIJI:** **Western Division:** Nadarivatu, Viti Levu Island, 17.566°S 177.966°E, 823 m, 10 Nov 1964, N. McFarland, 1♂ (AMNH\_PBI 00038586) (SAMA). **NEW CALEDONIA: Province Nord:** Ciu, 21.53333°S 165.95°E, 09 Jan 1969, N. L. H. Krauss, 1♀ (AMNH\_PBI 00042158) (BPBM). Hienghene, 20.68333°S 164.93333°E, 100 m, Jan 1971, N. L. H. Krauss, 5♂ (AMNH\_PBI 00042186–AMNH\_PBI 00042190), 4♀ (AMNH\_PBI 00042169–AMNH\_PBI 00042172) (BPBM). Poindimié, 20.93333°S 165.33333°E, 50 m, Jan 1969, N. L. H. Krauss, 2♂ (AMNH\_PBI 00042206, AMNH\_PBI 00042207), 1♀ (AMNH\_PBI 00042211) (AM). 15♂ (AMNH\_PBI 00042191–AMNH\_PBI 00042205), 3♀ (AMNH\_PBI 00042208–AMNH\_PBI 00042210) (BPBM). Vallée d'Amoa, 20.95°S 165.28333°E, 07 Feb 1963, N. L. H. Krauss, 1♀ (AMNH\_PBI 00042165) (BPBM). **Province Sud:**

Anse Vata, 22.31666°S 166.43305°E, 21 Mar 1961, J. Sedlacek, 1♀ (AMNH\_PBI 00042157) (BPBM). Cap Ndoua, site 1, rainforest, 22.38333°S 166.93333°E, 150 m, 28 Nov 2004, C. J. Burwell and G. B. Monteith, 1♂ (AMNH\_PBI 00178063), 1♀ (AMNH\_PBI 00178064) (QM); 21 Dec 2004, C. J. Burwell and G. B. Monteith, 1♀ (AMNH\_PBI 00178065) (QM). Foret Nord, site 2, 22.31666°S 166.91666°E, 200 m, 02 Dec 2004, S. G. Wright, 1♀ (AMNH\_PBI 00178066) (QM). Isle of Pines, Vao, 22.66888°S 167.489°E, 100 m, Jan 1985, N. L. H. Krauss, 1♂ (AMNH\_PBI 00042596), 1♀ (AMNH\_PBI 00042597) (BPBM). Kwe Binyi River, via Goro, 22.25801°S 167.0051°E, 16 m, 22 Apr 2005, Cassis, Wall, Tatarnic, Monteith, 2♂ (AMNH\_PBI 00006243, AMNH\_PBI 00006244), 2♀ (AMNH\_PBI 00006245, AMNH\_PBI 00006246) (AM). Les Bois du Sud campground, 22.17205°S 166.7604°E, 200 m, 23 Apr 2005–25 Apr 2005, Cassis, Wall, Tatarnic, Monteith, *Phyllanthus* sp. (Euphorbiaceae), 8♂ (AMNH\_PBI 00006215–AMNH\_PBI 00006222), 3♀ (AMNH\_PBI 00006231–AMNH\_PBI 00006233) (AM). *Phyllanthus* sp. (Euphorbiaceae), 8♂ (AMNH\_PBI 00006223–AMNH\_PBI 00006230), 6♀ (AMNH\_PBI 00006234–AMNH\_PBI 00006239) (AMNH). Mokoué to Dothio, 21.5667°S 166.1°E, 150 m, 20 Mar 1968–22 Mar 1968, J. L. Gressitt, 1♀ (AMNH\_PBI 00042167) (BPBM). Monts des Koghis, 22.167°S 166.53°E, 550 m, 20 Sep 1979, W. C. Gagne, *Macaranga* sp. (Euphorbiaceae), 3♂ (AMNH\_PBI 00042591–AMNH\_PBI 00042593), 2♀ (AMNH\_PBI 00042594, AMNH\_PBI 00042595) (BPBM). Mount Koghi, 22.1775°S 166.50888°E, 400 m, 12 Nov 1986–14 Nov 1986, R. L. Brown, 1♂ (AMNH\_PBI 00042154) (BPBM). Nouméa, 22.16°S 165.3°E, 5 m, 28 Feb 1960, J. L. Gressitt, 1♀ (AMNH\_PBI 00042159) (BPBM). Nouméa, 22.16°S 165.3°E, 20 Feb 1963, C. M. Yoshimoto, 1♀ (AMNH\_PBI 00042166) (BPBM). Refuge de Farino campground, 21.64869°S 165.7808°E, 269 m, 25 Apr 2005, Cassis, Wall, Tatarnic, Monteith, (Sapindaceae), 8♂ (AMNH\_PBI 00006247–AMNH\_PBI 00006254), 7♀ (AMNH\_PBI 00006240–AMNH\_PBI 00006242, AMNH\_PBI 00006255–AMNH\_PBI 00006258), 9 juveniles (AMNH\_PBI 00006259–AMNH\_PBI 00006267) (AM). Yahoué, 22.19583°S 166.49861°E, 100 m, Dec

1983, N. L. H. Krauss, 1♂ (AMNH\_PBI 00042155) (BPBM). Yate, 22.16666°S 166.95°E, 26 Mar 1968–27 Mar 1968, J.L. Gressitt & T.C. Maa, 1♀ (AMNH\_PBI 00012098) (AM). 1♀ (AMNH\_PBI 00042598) (BPBM). **Province des Iles Loyauté:** Maré Island, La Roche, 21.48333°S 168.03333°E, Mar 1959, N. L. H. Krauss, 1♂ (AMNH\_PBI 00042168) (BPBM).

*Coridromius zetteli* Chérot, Konstantinov and Yasunaga

Figures 5, 6, 17S–U, 21D, map 5

*Coridromius zetteli* Chérot et al., 2004: 58–60.

**DIAGNOSIS:** This species is recognized by the following combination of characters: bilobed proepisternum; embolium with yellow U-shaped fascia near cuneal fracture, lateral margins of hemelytra straight and carinate; male left paramere tightly corkscrewed with acute apex nearly parallel to axis. *C. zetteli* is similar to *sommelierii*, but can be distinguished by the slightly different shape of the left and right parameres and the much shorter ventral apical process on the pygophore. It is also very similar in appearance to *testaceus*; while *zetteli* is currently known only from males, *testaceus* is known only from females.

**REDESCRIPTION: COLORATION** (figs. 5, 6): Varying from light brown to almost uniform dark brown with yellow markings. Head: Light brownish yellow to orange-brown; frons typically yellow along midline, with very faint to distinct darker brown vittae; clypeus yellowed with darker brown markings at base, dark brown apically; mandibular plates yellowed; maxillary plates brownish orange; buccula yellowed; labrum dark brown basally; labium pale, brown apically. Antenna: AII dark brown, pale at base and apex; AII light orange-brown with dark brown subapical band, separated from dark brown apical annulation by narrow cream band; AIII and AIV dark brown, pale at base. Thorax: Pronotal collar yellow to orange-brown, darker anteriorly; pronotum light orange-brown to dark brown, with median yellow band (faint in some specimens), yellowed at humeral angles in paler specimens, lateral and posterior margins yellow; mesoscutum orange-brown to dark

brown; scutellum light orange-brown to dark brown with yellow midline, apex, and sides; thoracic pleura mostly orange-brown with posterior margins of each sclerite yellowed; mesopleuron dark brown ventrally. Hemelytra: Orange-brown to nearly uniform dark brown; in paler specimens clavus slightly paler at claval commissure and along posterior margin, immediately above cuneal fracture; embolium yellow along outer margin, with U-shaped yellow fascia at lateroapical margin; cuneus dark brown; membrane pale brown with dark brown veins. Legs: Pro- and mesofemur yellow at base, becoming orange-brown toward apex; metafemur pale basally, with 8 or 9 diagonal dark brown stripes, stripes merging with dark brown ventral patch in darker specimens, subapical lateral depressions dark brown. Abdomen: Orange-brown with alternating cream and dark brown markings along dorsolateral margin. SURFACE AND VESTITURE (figs. 5, 6): Head, pronotum, propleuron, metepimeron, scutellum, and hemelytra finely, shallowly punctate; head and dorsum clothed in long, silvery, decumbent setae. STRUCTURE (figs. 5, 6): Head: Frons broadly tumescent medially and depressed laterally; lateral tubercles adjacent to eyes small, bordered posteriorly by shallow depressions; posterior margin of head carinate and medially rounded. Thorax: Pronotum broad, lateral and posterior margins slightly carinate, anterolateral margins slightly turned upward, submarginal region of humeral angles excavated, collar region undifferentiated; proepisternum bilobed; posterior margin of metepimeron truncate; lateral lobes of metanotum prominent and flared; scutellum flattened. Hemelytra: Lateral margins of hemelytron straight and carinate; embolium with U-shaped fascia on lateroapical angle. Legs: Metatibial spines long and thick. Abdomen: Posterior margin of abdominal sternite II angular. MALE GENITALIA (figs. 17S–U, 21D): Right paramere broad, somewhat club-shaped, with thumblike lobe, appearing U-shaped when viewed from above (figs. 17Q–R, 21D); left paramere straight and tightly coiled five times along its axis, ending in a tapered point nearly parallel in direction to the paramere body (figs. 17S, 21A). FEMALE PARAGENITALIA: Unknown.

REMARKS: *C. zetteli* is known only from males. Since the corkscrewed left paramere of *zetteli* is nearly identical to that of *sommelierii*, we suspect that the female of *zetteli* should also have a comparable corkscrewed paragenital opening. As discussed above, *zetteli* is also very similar to *testaceus*, which is ironically known only from females. Both species were recently collected together in southern China by the senior author, raising the possibility that they may be conspecific. *C. testaceus*, however, lacks the corkscrewed paragenital opening we would expect if these were in fact the same species.

HOST: Recorded from *Macaranga* sp. (Euphorbiaceae) in southern China.

DISTRIBUTION: Known from southern China and Laos (map 5).

HOLOTYPE: Male: LAO PEOPLE'S DEMOCRATIC REPUBLIC: Sekong Prov.: Bolavens Plateau, N slope, ca. 10 km N. Mg Tha Theng, 15.5°N 106.43333°E, 500 m, 29 May 1996–30 May 1996, Schillhammer, (AMNH\_PBI 00189975) (NHMW).

PARATYPES: LAO PEOPLE'S DEMOCRATIC REPUBLIC: Sekong Prov.: Bolavens Plateau, N slope, ca. 10 km N. Mg Tha Theng, 15.5°N 106.43333°E, 500 m, 29 May 1996–30 May 1996, Schillhammer, 3♂ (AMNH\_PBI 00189974, AMNH\_PBI 00189976–AMNH\_PBI 00189977) (NHMW).

OTHER SPECIMENS EXAMINED: CHINA: YUNNAN PROV.: Xishuangbanna Tropical Rainforest Ecosystem Station, ~3 km from entrance, 21.96777°N 101.2075°E, 650 m, 30 May 2006, N. Tatarnic, *Macaranga* sp. (Euphorbiaceae), 2♂ (AMNH\_PBI 00006073, AMNH\_PBI 00006074) (AMNH).

## DISCUSSION

With this paper we increase the number of described species of *Coridromius* from 11 to 32 (33 if we include *schuhii*). With the exception of a few species, our conclusions are based on the examination of very few specimens collected from a limited number of localities. As a result, we suspect that the number of species of *Coridromius* must be greater than reported herein. For example, throughout Africa only three species of *Coridromius* are known from a total of six localities. In all likelihood there are many more species to be discovered in

Africa, once more sampling is undertaken in this relatively unexplored region. Similarly, while a relatively large number of species (five) have been described from Papua New Guinea, the specimens all result from a small number of collecting events, and again from relatively few localities. No doubt other known areas of high biodiversity, such as Borneo, Sulawesi, and other localities throughout the Old World tropics and subtropics, will ultimately yield more new species.

As the only mirid known to practice traumatic insemination, *Coridromius* arguably has the most unusual biology in this family. We believe that in order to fully understand the nature of this unusual form of mating, we must be able to confidently determine the phylogenetic placement of *Coridromius* within the Miridae, and also reevaluate the relationships of those other Cimicomorpha that practice traumatic insemination (i.e., the Prostemmatinae, Plokophilidae, Anthocoridae, Polycenidae, and Cimicidae). Unfortunately, it is this very form of mating that has canalized many of the morphological characters within *Coridromius* (and within those other traumatically inseminating taxa as well), making taxonomic placement difficult to assess with morphological characters alone. We hope this issue may be alleviated with the use of molecular components in addition to morphological data in future investigations into the systematic placement of these unusual bugs.

#### ACKNOWLEDGMENTS

We wish to thank Sue Lindsay (AM) for providing assistance with the scanning electron micrographs for most taxa, while Michael D. Schwartz kindly provided the scanning electron micrograph of *C. carinatus*. Hannah Finlay (AM) assisted with the dissection of female specimens and kindly provided her rough sketches to be used for further illustrations. Gareth Carter and Michael Elliot (AM) helped in the preparation of the distribution maps. Matthew Bulbert (AM) created the excellent photographs of *C. monotocopsis* on *Monotoca elliptica*, Tomohide Yasunaga (JICA) provided his beautiful image of *testaceus*, and Chris Darling (Royal Ontario Museum) also provided his

photograph of *chinensis* on an unidentified *Macaranga* tree. Chris Darling and Qing-Jun Li (Xishuangbanna Tropical Botanical Garden) were instrumental in assisting with NT's fieldwork in southern China and Laos.

Geoff Monteith (Queensland Museum) was invaluable as guide, companion and collector during fieldwork in New Caledonia. Michael Wall (San Diego Natural History Museum), Celia Symonds (AM), Christiane Weirauch (University of California, Riverside), and Randall Schuh (AMNH) must also be thanked for their outstanding collecting efforts across Australia and New Caledonia.

We wish to thank all those who provided the material necessary to complete this project. A special thank you to Frédéric Chérot and Tomohide Yasunaga, who entrusted us with the only specimens of several important species. Likewise, this work would have been impossible were it not for the generous loan of all *Coridromius* specimens from the Bishop Museum, where most of the material is housed. Similarly, Mick Webb loaned us many specimens from the Natural History Museum, London, Randall Schuh provided us with a nearly infinite series of Australian specimens and Tom Henry from the NMNH loaned us the one and only specimen of *Coridromoides carinatus*.

We also thank Dieter Hochuli for his constant support, advice, and good humor. This manuscript was significantly improved with the reviews of Randall Schuh, Michael Schwartz, and Frédéric Chérot. Lastly, we are grateful to Katrina Menard, who read through this manuscript and provided invaluable editorial suggestions.

This research was conducted during the course of the senior author's Ph.D. research, which was funded by the NSF Planetary Biodiversity Inventories grant DEB-0316495 to R. T. Schuh and G. Cassis.

#### REFERENCES

- Arnqvist, G., and L. Rowe. 2005. Sexual conflict. Princeton, NJ: Princeton University Press, 360 pp.  
Carayon, J. 1959. Insémination par 'spermalege' et cordon conducteur de spermatozoïdes chez *Stricticimex brevispinosus* Usinger (Heteroptera, Cimicidae). Revue de Zoologie et de Botanique Africaines 60: 81–104.

- Carayon, J. 1966. Traumatic insemination and the paragenital system. In R.L. Usinger (editor), Monograph of the Cimicidae (Hemiptera-Heteroptera), 81–166. College Park, MD: Entomological Society of America.
- Carayon, J. 1977. Insémination extragénitale traumatique. In P.P. Grassé (editor), Traité de Zoologie 8(V–A): 351–390. Paris: Masson.
- Carvalho, J.C.M. 1952. On the major classification of the Miridae (Hemiptera). (With keys to subfamilies and tribes and a catalog of the world genera.) Anais da Academia Brasileira de Ciências 24: 31–110.
- Carvalho, J.C.M. 1955. Keys to the genera of Miridae of the world (Hemiptera). Boletim do Museu Paraense Emilio Goeldi Nova Serie Zoologia 11: 1–151.
- Carvalho, J.C.M. 1956. Insects of Micronesia. Heteroptera: Miridae. Insects of Micronesia 7: 1–100.
- Carvalho, J.C.M. 1958. A catalogue of the Miridae of the world. Part III. Arquivos do Museo Nacional Rio de Janeiro 47: 1–161.
- Carvalho, J.C.M. 1987. The genus *Coridromius* Signoret, with descriptions of new species (Hemiptera, Miridae). Revista Brasileira de Entomologia 31: 61–69.
- Cassis, G., and G.F. Gross. 1995. Hemiptera: Heteroptera (Coleorrhyncha to Cimicomorpha). In W.W.K. Houston and G.V. Maynard (editors), Zoological Catalogue of Australia. Vol. 27.3A. Melbourne: CSIRO, Australia, 506 pp.
- Cassis, G., and L. Vanags. 2006. Shield bugs of Australia (Insecta: Heteroptera: Scutelleridae): generic conspectus, new species, host plants, and classification. Denisia 19: 275–398.
- Cassis, G., Wall, M., and Schuh, R.T. 2007. Insect biodiversity and industrializing the taxonomic process: the plant bug case study (Insecta: Heteroptera: Miridae). In T.R. Hodkinson, J. Parnell, and S. Waldren (editors), Taxonomy and systematics of species rich taxa: towards the tree of life, 193–212. Boca Raton: CRC.
- Chérot, F., F. Konstantinov, and T. Yasunaga. 2004. Two new plant bugs of the genus *Coridromius* Signoret, 1862 from the Oriental Region with a new synonymy (Heteroptera, Miridae, Orthotylinae, Halticinae). Belgian Journal of Entomology 6: 57–67.
- Distant, W.L. 1914. Rhynchota from New Caledonia and the surrounding islands. In F. Sarasin and J. Roux (editors), Nova Caledonia zoologie 1: 4: 10: 369–390.
- Samouelle, G. 1819. The entomologist's useful compendium; or an introduction to the knowledge of British insects, .... London: Thomas Boys, 496 pp.
- Linnauvori, R.E. 1994. Orthotylinae of West, Central and North-east Africa (Heteroptera, Miridae). Acta Zoologica Fennica 193: 1–84.
- Liu, G.Q., and R.J. Zhao. 1999A. New species of genus *Coridromius* Signoret from China (Heteroptera: Miridae). Acta Zootaxonomica Sinica 24: 55–58.
- Miyamoto, S., and T. Yasunaga. 1999. Discovery of the genus *Coridromius* Signoret (Heteroptera: Miridae) from Japan, with descriptions of two new species. Biogeography 1: 33–37.
- Montrouzier, P.A. 1861. Essai sur la faune entomologique de la Nouvelle-Calédonie. Annales de la Société Entomologique de France (4)1: 59–74.
- Morrow, E.H., and G. Arnqvist. 2003. Costly traumatic insemination and a female counter-adaptation in bed bugs. Proceedings of the Royal Society of London Biological Sciences Series B 270: 2377–2381.
- Poppius, B. 1911. Beiträge zur Miriden-Fauna Australiens. Översigt af Finska Vetenskaps-Societetens Förfärlingar 53A(3): 1–16.
- Reinhardt, K., R. Naylor, and M.T. Siva-Jothy. 2003. Reducing a cost of traumatic insemination: female bedbugs evolve a unique organ. Proceedings of the Royal Society of London Biological Sciences Series B 270: 2371–2375.
- Schuh, R.T. 1974. The Orthotylinae and Phylinae (Hemiptera: Miridae) of South Africa with a phylogenetic analysis of the ant-mimetic tribes of the two subfamilies for the world. Entomologica Americana 47: 1–332.
- Schuh, R.T. 1995. Plant bugs of the world (Insecta: Heteroptera: Miridae). Systematic catalog, distributions, host list, and bibliography. New York: New York Entomological Society, 1329 pp.
- Signoret, V. 1862. *Coridromius* n. n. f. *Ocypus* Montrouzier. Bulletin de la Société Entomologique de France (4)2: 5.
- Stutt, A.D., and M.T. Siva-Jothy. 2001. Traumatic insemination and sexual conflict in the bed bug *Cimex lectularius*. Proceedings of the National Academy of Sciences of the United States of America 98: 5683–5687.
- Tatarnic, N.J., G. Cassis, and D.F. Hochuli. 2006. Traumatic insemination in the plant bug genus *Coridromius* Signoret (Heteroptera: Miridae). Biology Letters 2: 58–61.
- Van Duzee, E.P. 1932. New Hemiptera Heteroptera from the Marquesas. Bulletin of the Bernice P. Bishop Museum 98: 180.
- Walker, F. 1873. Catalogue of specimens of Hemiptera Heteroptera in the collection of the British Museum. Part VI. London: British Museum, 210 pp.
- Woodward, T.E. 1954. New records and descriptions of Hemiptera Heteroptera from the Three Kings Islands. Records of the Auckland Institute and Museum 4(4): 215–233.