


5896 -- 6242

BIRDS

BIRDS.







Pange  
Sept. 12, 1914.

5896. Sunbird  
1892. An

Stomach  
chysal  
in tree  
were

3 we

Beng

5897. Podiceps  
brown

bill

and

feet

me

con

feet

5898. Podiceps  
brown

gray

at s

feet

rem

of a s

P-

and

rea

rivers

Panga

Sept. 12. 1914.

1160

5896. Sunbird ♂ testes slightly enlarged.  
5892. *Anthreptes aurantiium*

Stomach content: 4 caterpillars & 2

chrysalids. Like no 5896, feeding  
in trees along river bank. No more  
were noticed on the Amurimi, but  
3 were seen on the Lindi, below  
Bengamisa.

5897. Podica. ♂. imm. Iris rather light greenish brown, rim of eyelids dusky brown, bill blackish, but underside of mandible and cutting edges of both jaws whitish, feet orange, dusky brown on back of metatarsi and about joints. Stomach contained remains of insects and a few bits of quartz.

5898. Podica. ♀. ovary slightly enlarged. Iris brownish yellow, rim of eyelids brownish gray; bill dusky brown above, orange red at sides and whitish below (towards base) feet orange-red. Stomach contained remains of insects, including wings of a small dragon-fly.

These two "Sun Grebes", mother and young, were swimming near a low grassy part of the river bank. They did not dive, but made for the bank. The

Panga  
Sept 13.  
58991

Foot

59001

59002

59003

59004

59005

59006

59007

59008

59009

59010

59011

59012

59013

59014

59015

59016

59017

59018

59019

59020

59021

59022

59023

59024

59025

59026

59027

59028

59029

59030

59031

59032

59033

59034

59035

59036

59037

59038

59039

59040

59041

59042

59043

59044

59045

59046

59047

59048

59049

59050

59051

59052

59053

59054

59055

59056

59057

59058

59059

59060

59061

59062

59063

59064

59065

59066

59067

59068

59069

59070

59071

59072

59073

59074

59075

59076

59077

59078

59079

59080

59081

59082

59083

59084

59085

59086

59087

59088

59089

59090

59091

59092

59093

59094

59095

59096

59097

59098

59099

59100

59101

59102

59103

59104

59105

59106

59107

59108

59109

59110

59111

59112

59113

59114

59115

59116

59117

59118

59119

59120

59121

59122

59123

59124

59125

59126

59127

59128

59129

59130

59131

59132

59133

59134

59135

59136

59137

59138

59139

59140

59141

59142

59143

59144

59145

59146

59147

59148

59149

59150

59151

59152

59153

59154

59155

59156

59157

59158

59159

59160

59161

59162

59163

59164

59165

59166

59167

59168

59169

59170

59171

59172

59173

59174

59175

59176

59177

59178

59179

59180

59181

59182

59183

59184

59185

59186

59187

59188

59189

59190

59191

59192

59193

59194

59195

59196

59197

59198

59199

59200

59201

59202

59203

59204

59205

59206

59207

59208

59209

59210

59211

59212

59213

59214

59215

59216

59217

59218

59219

59220

59221

59222

59223

59224

59225

59226

59227

59228

59229

59230

59231

59232

59233

59234

59235

59236

59237

59238

59239

59240

59241

59242

59243

59244

59245

59246

59247

59248

59249

59250

59251

59252

59253

59254

59255

59256

59257

59258

59259

59260

59261

59262

59263

59264

59265

59266

59267

59268

59269

59270

59271

59272

59273

59274

59275

59276

59277

59278

59279

59280

59281

59282

59283

59284

59285

59286

59287

59288

59289

59290

59291

59292

59293

59294

59295

59296

59297

59298

59299

59300

59301

59302

59303

59304

59305

59306

59307

59308

59309

59310

59311

59312

59313

59314

59315

59316

59317

59318

59319

59320

59321

59322

59323

5

Panga

Sept. 13, 1914.

*Estrilda mefoda*

5899 Red-cheeked Waxbill ♂. testes slightly enlarged. Iris brown, shading to light gray on inner rim; bill scarlet; feet rather dark grayish brown. Crop and stomach contained grass-seeds.

*Estrilda mefoda*

5900 Same as preceding. ♀. o. e. Iris brown, slightly reddish exteriorly; bill dull scarlet, feet rather dark grayish brown. Crop & stomach contained grass-seeds.

*Estrilda mefoda*

5901 Same as preceding ♀. imm. Crop and stomach contained grass-seeds.

This species was <sup>first</sup> seen yesterday, some distance above Panga, but is common here, feeding in small flocks on the tall grasses. The

Common  
further  
down  
Aruwimi  
at Stanley  
ville.  
& Boma

finely-barred species <sup>A.</sup>, so common at Avakubi, seems to be absent, nor have we seen the black-bellied

*Estrilda atricapilla*


*Estrilda since* leaving Avakubi.  
(<sup>1</sup> of the black bellied species seen at Stanleyville, first week in Oct. 14) <sup>See no. 5874</sup>

So far as I am aware, Waxbills and "mannikins" make very little noise, only very weak or lisping notes being heard from them, and nothing in the way of a song.

5902 Gray Flycatcher. ♂ testes slightly enlarged. Iris dark brown, bill black, feet dark bluish-gray. Stomach contained 2 caterpillars.

Panga.

Sept. 13. 1914.



remains of insects

5903. mate of preceding ♀. ovary slightly enlarged

Iris dark brown, bill black, feet dark bluish-gray. This pair of flycatchers was found in a tree in the second-growth back of the station at Panga.

5904. Bee-eater (m. galario) ♂ + <sup>australis</sup> w.e. Iris rather dark crimson, bill & feet black.

(The related species with rufous back, was seen at Bombuva Sept. 21.)

Sept. 14. <sup>Pygochelidon</sup> ~~Pygochelidon~~ schlegeli 5905. ♂. testes somewhat enlarged, skull not quite fully ossified. Stomach contents rice.

5906. <sup>Pygochelidon</sup> ~~Pygochelidon~~ schlegeli Same as preceding. ♂. testes somewhat enlarged.

Skull not quite fully ossified. Iris dark brown, rim of eyelids pinkish, bill blackish at base, scarlet at tip and sides; feet light brownish. Crop and stomach contains rice.

5907. <sup>Pygochelidon</sup> ~~Pygochelidon~~ schlegeli Same as preceding. ♀. ovary somewhat enlarged. Iris dark brown, rim of eyelids blackish; bill blackish at base, scarlet at tip and sides, feet rather light dull brownish. Stomach contained rice.

These wasbills are regular inhabitants of rice fields when the grain is ripening or being cut, but instead of clinging to the heads, they feed on the ground.

Panga  
Sept 14.

wide

wall

swif

an

they

the

dry

last

sub

fla

tim

freq

the

will

bus

prep

the

for

up

fla

th

an

Ga

of

Panga

Sept 14. 1914.

evidently on fallen grains, keeping well concealed, and generally darting swiftly and silently, if flushed, into a neighboring thicket. Sometimes they will fly off and alight again in the rice, but are as a rule decidedly shy. The rice-harvest, in this region, lasts from August to January or February, for different fields may be planted at wide intervals. At other times of the year the birds seem to frequent the almost impenetrable thickets about old plantations and villages, but are only very rarely to be seen along paths.

Just at present they are paired, preparatory to nesting, but by February they form small flocks with their young.

Of course many other birds are fond of rice. Here we have noticed especially, engaged in this village, flocks of *Hypothymis* and *Spermestes* (of the latter 2 species, the larger one, and the dark backs) a few *Pyrenestes* (small one) and an occasional pair of partridges (*Francolinus spearmatus*). The waxbills of the genus  *Estrilda* seem never to eat rice, their bills being probably too small.

Panga  
Sept. 14. 1914.

5909



Panga  
Sept. 14. 1914.

Just as the natives of the Uele are obliged to mount guard over their fields of dura-grain or "vundi", so those here need to chase the birds from their rice. A small shelter is often erected for a watcher on some commanding termite-hill, and long strips of bark, as cords, laid over the rice in different directions, being tied to any bushes or small banana trees there happen to be. The jerking of the ends of these cords, several of which may be knotted together where they intersect, serves to frighten away birds feeding in the grain.

5910

5908. *Stiphrornis ruficollis* <sup>var. *anthracaster*</sup> Orange-breasted Warbler ♀. o. e., an egg in oviduct. Stomach contained remains of insects and a caterpillar. Seems not to require virgin forest, for I have several times seen it in badly mutilated woods, in the disagreeable undergrowth that usually springs up under such conditions. Has a low protracted scolding note that I have found very difficult to distinguish from one of the calls of the smaller Turdines.

Sept. 15  
5911.

5912.  
5913.

5914.

5915.

Panga

Sept. 14. 1914.

5909. *Audubonius serimus* A greenish bulbul with yellowish under-parts. ♂ imm. testes slightly enlarged, Iris rather light brown, bill dull light brown, feet dark gray. Stomach contents: berries. It is strange that I never noticed this bulbul at Avakubi, tho several were seen at Panga, and along the Lindi near its mouth.
5910. *Tricholaemus purpuratus* Long-tailed Barbet. ♂. t. n. e. Stomach contents: berries and a snail. Iris carmine. Heard commonly along the Aruwimi, as is also *Tricholaema*, *Buccanodon*, the red-rumped *Barbatula*, etc.
- Sept. 15. 1914. 5911. *Malimbus subnudulus centralis* Black Weaver with scarlet crown & nape. ♂ testes slightly enlarged. Stomach contained 4 caterpillars and remains of insects.
5912. *Hypargos schlegeli* Pytela. ♂. t. e. Crop & stomach contained rice.
5913. " " ♀. t. e. Stomach contained sand & a gram of (rice)
5914. " " ♀. immature. Ovary much enlarged, an egg in oviduct. Stomach contained sand and rice.
5915. Black Titmouse. ♂. t. e. Stomach contained a green caterpillar. A number of these were feeding in tall trees about a rice-field, uttering single short whistles and

Panga

Sept. 15, 1914.

Panga  
Sept 16



5916. Shrike (*Sigmodon*) ♂ t.e. Stomach  
contents: insects mainly orthopterous.

5917. Same as preceding. ♀ imm. S.C.: insects.

Two flocks of about 6 each were met with on the borders of a clearing. They show great solicitude for a wounded member of the party.

Like most of the family, they have harsh notes in addition to their softly whistled call.

5918. Bulbul. ♂ imm. Stom. cont. pieces of a beetle & other soft insects. Iris gray, rim of eyelids yellowish-green, maxilla mainly dusky, mandible dirty greenish-gray, corners of mouth and edges of both jaws yellow; feet pale bluish-gray, claws pale yellowish.

Seems not to be a common species.

Was sitting in a small tree in forest, raising its wings and giving a scolding note.

- Sept. 16 *Exilides melpoda*  
5919. Red-chested Waxbill. ♀. ovaries slightly enlarged; skull not quite completely ossified. Crop contained grass-seeds.

5920. Gray Warbler. ♂. testes slightly enlarged. Iris grayish-brown, maxilla black, mandible dull gray, feet light blue.

Stom

lun

sp. 21

dark

dull

blue

green

Stom

an

5921

cont

gray

slate

light

brown

black

grey

green

yellow

red

orange

purple

pink

white

black

grey

yellow

red

orange

Panga

Sept. 16. 1914.

Stomach contents: insect remains.

Found in old, fairly open, second-growth.

5921. Green-backed Cuckoo. ♂. imm. Iris <sup>*Chrysococcyx intermedius*</sup> dark grayish-brown; bill greenish, dusky above, brighter below, and slightly bluish at sides of base, rim of eyelids greenish; feet blue, claws black.

Stomach contents: remains of insects

and 1 hard seed. Shot from a tall tree in a clearing

Sept 17. *Muscicapa striata* 5922. Woodpecker. ♂. t. n. e. Stomach

contained 4 insect larvae (coleopterous)  
Iris dark crimson, tip of bill whitish  
rest of maxilla black, mandible shading  
posteriorly to pale bluish; feet olive.

Was drumming at intervals,  
like *M. xantholophus*, but not very  
loud. A couple of others were  
also heard nearby. They seemed to  
like large trees overlooking second-  
growth, and gave also an  
excited, reiterated vocal sound,  
not very loud - ~~notes~~ nearby  
so loud, for instance, as that of  
*M. goertae*, which it resembles slightly.

5923. Drongo. ♂. t. n. e. Iris scarlet,  
<sup>*Dicrurus caerulescens*</sup> bill and feet black. S.C.: insect remains.  
Found low down in second growth.

Panga  
Sept 17. 1914.

but usually favors tall trees dominating second-growth or about clearings.

Common between Banalia & Bengamisa has loud, harsh, & semi-metallic voice.

5924. Bulbul (*Ixonotus*) ♀ imm. Iris <sup>guttatus</sup> light grayish-brown; maxilla dark brownish-gray, mandible light gray; feet bluish-gray. Stomach contents: fruit & 4 small caterpillars.

5925. *Pytilia schlegeli*.

5926. Brown-chested Woodpecker. ♀. s.w.e.

Stomach contents: ants and a soft pupa of some larger insect.

5927. Black Weaver (*Melanerpes coronatus*) ♀. ovary slightly enlarged. Stomach contents: insects.

Was engaged, with her scarlet-crowned mate, in completing their nest, attached to the drooping end of a slender, thorny acacia-vine, 30 feet from the ground. This nest was mainly composed of dry spiral tendrils from some vine, and small dry twigs and leaf stems, the "spout" not very long, but as usual with this genus, decidedly wide (7 cm.) above the nest additional material



Panga. Sept 17. 1914.

1170

(including some green moss)  
had been hung to the supporting  
creeper, so that the total length  
of the structure was 51 cm.

Both male & female were at work,  
the former occupied with the entrance,  
the latter with the top.

Sept 18. *Sturnida myopoda* ♂  
5928 Pratincole ♀. o.m.e. Stomach contained  
Glareola nordmanni (ad. in winter plumage)  
several beetles.

5929. Pratincole. ♀. im. Stom. cont. beetles.

Glareola nordmanni

No. 5928 + 9 alighted in the station - the  
first seen this season.

5. [Altho the water of the Aruwimi  
is now very high, covering most of the  
rocks, the smaller Glareola nuchalis  
was common from Avakubi to  
(But apparently not near Banaliza)  
Panga, often perching on dead  
limbs sticking up out of the stream.

I indeed a large flock was once  
seen in a leafy tree standing on  
a shoal in the river, where a  
colony of weavers was also established.

All those I saw had the white collars.

Sept. 19. (Left Panga) *Laniarius leucorhynchus*

5930. Black Bush-Shrike. ♀. ovaries slightly  
enlarged. Iris dark red-brown. Stomach  
contents: insects.

Sept 20 (arr. Bombwa)

5931. Skeleton of Kingfisher (*C. rudis*) ♀.

*Sitara monacha* Sharpe.

5932. Small yellow Weaver with black tips to  
tail.

# Banalia

Sept. 20. 1914



testes somewhat enlarged. Stom. cont. insects.

This is a bird we collected while coming up the Congo, and now met with again for the first time at a village called "Bombwa" on the left bank of the Aruwimi, 25 or 30 miles above Banalia.

5933. Dove (*Chalcopelia*). ♂ t.e. Common hereabouts, calling continually. The notes resemble those of *Tympaniotria*, the voices of the two species are usually easy to distinguish. *Chalcopelia* here is found mainly in the neighborhood of villages, *Tympaniotria* in second-growth, and *Calopelea* in forest, though this is not of course an absolute rule. At Paradjie, for example, *Chalcopelia* frequents the open brush, and *Tympaniotria* the woods. [*Calopelea*, besides being seen at Panga, was heard along the Aruwimi, and between Banalia & Bengamisa.]

Sept. 21. (Bombwa)  
5934. Chestnut Babbler (*Phyllanthus*) ♂ testes

slightly enlarged. Iris dark brownish-red, bill creamy greenish-yellow, feet pale grayish-green. Stomach contents: remains of insects.

Banalia  
Sept. 21. 1914

5935. ♀

5936. ♂

5937. ♂

5938. ♂

5939. ♂

5940. ♂

5941. ♂

5942. ♂

5943. ♂

5944. ♂

5945. ♂

5946. ♂

5947. ♂

5948. ♂

5949. ♂

5950. ♂

5951. ♂

5952. ♂

5953. ♂

5954. ♂

5955. ♂

5956. ♂

5957. ♂

5958. ♂

5959. ♂

5960. ♂

Banalia (Village Bombwa)

Sept 21, 1914

5935. Young of preceding. ♂ imm. Stom cont: insects.

5936. Same. ♂ imm. Iris dark brown, rim of eyelids greenish-yellow, bill greenish-white, deep yellow along cutting edges towards base, feet pale grayish-green. Stom cont. a grasshopper & other insects.

5937. <sup>Andropadus sp.</sup> Greenish Bulbul ♀ c.m.e. S.C. berries.

5938. Short-tailed Warbler (*Sylviella denti*?) ♂ testes somewhat enlarged. Iris light brown, bill dusky, feet pinkish-brown. Stomach contents: insects. Found in undergrowth of some rather open woods.

5939. <sup>Rufa, vulpina</sup> Bathmedonia <sup>Rufa, vulpina</sup> ♀ ovary slightly enlarged

5940. Thrush (*Cossypha*) <sup>cyanoptera</sup> ♂ t.e. S.C. insects.

Very common along Aruwimi

5941. <sup>Rufa, vulpina</sup> Small Weaver ♂. testes somewhat enlarged

5.5932. Stomach contents: insects.

Just behind the "gîte d'âpage"

there were 3 nests of this weaver suspended from the tips of the tall grasses. <sup>8 ft. above the ground</sup> These nests are peculiar in being woven partly of the seed-bearing grass-tips, as well as strips from the leaves.

The nest of no. 5941 contained 2 young, whose feather-sheaths were beginning to burst. The nest

Baralia

Sept. 21. 1914.



down that still remained was grayish-white.

The other nests were empty, but a gray flycatcher (no. 5942) was seen to enter one of them several times, coming out and perching beside the <sup>empty</sup> nest.

5942. Flycatcher. ♂ t.e.  
*Pedicularis hamatus*

5943. Large black-headed Bee-eater. ♀.  
*Nectarogone brevirostris*  
ovary slightly enlarged. Iris bright crimson  
bill black, feet dark greenish-gray.  
Stom. cont.: pieces of a beetle & a wasp.

Flew about village on river-bank;  
had very weak voice. Another seen  
on an island near Stanleyville.

5944. Skeleton of Machaer <sup>Whamphus</sup> andersoni. ♀.  
ovaries slightly enlarged. Stomach  
contents: a small brown bat (*Pipistrellus*)  
remains of another bat, and a few feathers.

Sept. 22. Flying about village at dusk. *Hirundo nigrita*

5945. Dark blue Swallow. ♂. testes somewhat  
enlarged. Very common along Aruvimini.

5946. Weavers. ♂. testes slightly enlarged. Stomach  
contained seeds & a few pieces of insects.

Sept. 23. 5947. Small yellow bellied Sunbird with green back.

♀. ovary slightly enlarged. *Anthrax collaris hypoleucus*  
Her nest, of the usual pear shape, was  
composed of strips of dry grass and other  
vegetable material, lined with plant-  
down, bound together exteriorly with

Bansbie  
Sept. 23. 1

silk,

even

lung

20 fu

petal

wor

inf

and

part

5948. Sta

5949. Sm

ace

res

mor

gre

yellow

in

Sept. 25. 13

5950. La

5951. Ch

Iri

ye

ba

g

ca

g

ba

g

ca

1-167

Banalia

Sept. 23. 1914.

silk, and adorned with a small amount of caterpillars excrement. It hung from a creeper, at a height of 20 feet, directly over a path; in a patch of second-growth. The 2 young were just ready to leave the nest, in fact they did so when disturbed, and had the plumage of the upper parts metallic green.

5948. Starling. <sup>Lamprocorax purpureiceps</sup> ♀. s.m.e. S.C.: a few small berries.

5949. Small Hawk. <sup>Accipiter sharpei</sup> ♂. t.m.e. Iris bright scarlet, eyelids orange, cere & corner of mouth yellow, bill black, with mandible grayish at base; feet bright cadmium yellow, claws black. Stomach contained insect-remains and feathers. Seen pursuing a small bird.

Sept 25. Banalia & Mbanga

5950. Larger Tropic. ♂. t.m.e. S.C.: caterpillars.

5951. Chrysocolaptes flavicularis. ♂. testes slightly enlarged. Iris dull chrome-yellow, edge of eyelids light yellowish-green, bill dull greenish-yellow, base of maxilla blackish, feet dull yellowish-green, claws black, Stomach contained caterpillars and a few insects.

Shot in a tree devastated by caterpillars.

This is a bird of the tall trees, in, the virgin forest as well as those overlooking second-

Banalia  
Sept 25. 1914.



growth or plantations, very difficult to see, and more so to kill. Its voice is not infrequently heard, the common note being a series of short, sweet whistles, loud enough to be heard plainly for several hundred yards. Sometimes two birds call together. Once I heard another note of a loud, but less musical whistles, that reminded me of *Eurystomus gularis*.

It was in last February that I first became acquainted with these notes, and have since heard them <sup>many times</sup> near Penge, Avakubi, & Nkayu, <sup>as well as</sup> at Pangia.

5952. Thrush (*Erythropygia*). ♂ testes slightly enlarged. Stomach contents: insects.

5953. Warbler (*Sylvietta*). ♂ imm.

5954. Hawk ♂ juv. Iris yellowish-green.  
S. 5949. ~~Accipiter sharpei~~ rim of eyelids ochre, cere and corners of mouth chrome-yellow, bill black, feet cadmium yellow, claws black. Stomach contained flesh & feathers of a small bird

Sept. 26. (Mbanga - mchunge)  
5955. Warbler (*Sylvietta*) ♂ imm. S.C. a caterpillar

S. 5958. <sup>dentata</sup> and insects. Shot from a tree in a former village site, now grown up with thickets.

5956. Thrush (*Alethe*). ♀ ovary slightly enlarged

Banalia  
Sept 26

5954. K.  
Hale  
Stom

way  
the  
also  
(Co

ce  
Oa  
in  
su  
ca

5958. Q.  
5959. Pa  
5960. Q.  
5961. S.

F

Banalia

Sept 26. 1914.

5957. King fisher. ♂. testes slightly enlarged  
~~Haleyornis malimbicus prenticii~~.  
Stomach contained bones of a frog. Heard  
very frequently along the Aruwimi, where  
the chestnut-backed Haleyornis seemed  
also common.

[Coming down the Aruwimi we saw  
or heard rather often Ceryle maxima, but  
Ceryle rudis was somewhat less numerous.  
One or two Alcedo were noticed nearly  
every day, but Corythornis was not  
seen once, and Ispidina did not seem  
common.

5958. Fly catcher, ♂. t. n. e. S.C.: caterpillars & insects.  
(Smithornis)  
<sup>Apololemus</sup>

5959. Parisoma, ♀. t. n. e. Stom. cont: caterpillars.  
<sup>plumbeum</sup>  
<sup>cognitus</sup>

5960. Fly catcher (Erythrocercus) ♂ im.

5961. Small Trogan. ♂. t. n. e. S.C.: caterpillars.  
Apolodernas nigra.

The excrement of the 2 trogons here has a  
far more foul odor than usual among birds,  
perhaps due to their diet of caterpillars.

This trogon is a forest bird, and  
was seldom heard on this road.

During the first of the four days between  
Banalia & Beniarnisa, we saw no real  
forest at all, everywhere villages or  
places formerly cultivated. During the  
other three days there were occasional  
stretches of forest, interspersed with

# Banalia

Sept. 26, 1914.

Bengamisa  
Sept. 28, 1914.

numerous clearings and "matangos".

From Bengamisa to Kaparata, too,  
practically no forest has been left.

5962. Cuckoo. ♀ juv. Stomach contents: beetles.  
*Cuculus solitarius*.

Iris dark grayish-brown, rim of eyelids  
dull greenish; maxilla black, mandible  
dull yellowish-green with black tip.  
corners of mouth reddish-orange; feet  
bright yellow. (Very dark colored barred below)

- Sept. 27. (mchungu - mbedu) *Sturnurus*  
5963. Titmouse (*Anthoscopus*) ♀. Ovary slightly

enlarged, skull not completely ossified. Iris  
dark brown, bill dusky, but light gray along  
edges of mandibles and at base; feet bluish.  
Mouth remained slightly open after death as in Wood Hoopoes.  
One of a small flock. These titmice were

never seen at Avakubi.


*Sturnus phoeniceus*

5964. Fly catcher. ♂ t.m.e. Iris dark brown,  
tip of bill black, rest of bill, corners of  
mouth, and rim of eyelids blue, whole  
interior of mouth greenish yellow, feet  
rather light blue.

5965. Hawk. ♂. testes slightly enlarged. Iris  
*Accipiter sharpei* scarlet. Stomach contents: pieces of an insect.

5966. Rail (*Corithruwa*) ♂. testes slightly enlarged.  
Stomach contained a yg hemipter & other insect-  
remains.

[The similar rail (= *S. elegans*)  
with brown spotted  
back, that calls mournfully at night,



Bengamisa  
Sept. 28. 1914.

was heard 2 or 3 times at Banalia.]

Sept. 28. (mbudu - Bengamisa) *Oscus minucius*

5967. Flycatcher ♂. testes slightly enlarged; skull not quite completely ossified.

5968. <sup>Panisoma holopodum</sup> Gray Warbler ♂. testes somewhat enlarged

5.5920

Stomach contents: a small caterpillar and insect remains. Iris dark brown, rim of eyelids pinkish brown, bill black, becoming light gray beneath mandible; feet rather light blue. Shot in woods near a stream, found by following its note, which may be distinguished from that of the species like no. 5959 by having regularly 4 syllables, somewhat run together.

5969. Swallow (<sup>H. lucida</sup> *H. angolensis*) ♂. t.n.e.

5970. " " " ♂ .. .

5971. " " " ♂ .. .

5972. " " " ♂ .. .

5973. <sup>H. lucida</sup> Iris dark brown, bill & feet black ♀. one. skull not quite fully ossified.

This swallow has not been observed since we left Stanleyville in 1909, but would seem to be common at Bengamisa. For fully 30 or 40 were flying about one of the houses during the early afternoon, alighting beneath the roof of the porch. They were not nesting, nor did they roost there at night. Their resemblance to

Bengamisa

Sept 28. 1914.

*H. rustica*, in habits, flight, and voice is extremely close, and they are probably also fond of villages, for no more were noticed till we were very close to Stanleyville.

No. 5980 (Note) A nest of this gray flycatcher containing 2 eggs, collected near Bomile on Sept 11, 1914, was placed 4 ft above the water in a fork of a branch sticking up near the shore. Along that part of the Aruwimi 2 other nests of the same bird were observed in similar exposed positions, and 1 in a hollow of a larger stub.  
measurements of Bomile eggs: 18.0 x 13.3; 18.5 x 13.3

me, but the large wholly black member of the genus was not observed. On the Lindi (which natives here - Bamanya - call the "Luindi") we saw both the species like no 5974, and the small Chatuna with much white on underparts & rump. (*C. sabini*)

5975. Palm Swift. ♀. ovary slightly enlarged.  
*Tachymenoides parvus*  
Stomach contained many winged ants and a small beetle. Practically none noticed during the trip down the Aruwimi.
5976. Green Sunbird. ♂. testes somewhat enlarged,  
*Cyanomitra obscurus*  
shell not quite completely ossified. Stomach contained 2 hard round seeds from berries.
5977. Paroquet. (*A. zenkeri*) ♂. testes slightly enlarged  
iris orange. Crop contained small seeds.  
One of a flock of 5 or 6, that came to feed in a wild fig tree.

## Bengamisa

Sept 28. 1914.

*H. rustica*, in habits, flight, and voice, is extremely close, and they are probably also fond of villages, for no more were noticed till we were very close to Stanleyville.

The extensive white patches in the tail

make recognition easy in life  
coming in flocks to catch winged termites leaving nest.  
Common at Stanleyville  
sitting on rocks  
and roofs near rapids.

Sept. 29 (Bengamisa - Kaparata)

5974. Large stub-tailed Swift. ♂ [im] testes very small. *Chaetura cassini*. Stomach filled with winged ants. Seen occasionally along the Aruwimi, but the large wholly black members of the genus was not observed. On the Lindi (which natives here - Bamanga - call the "Luindi") we saw both the species like no 5974, and the small *Chaetura* with much white on underparts & rump. (*C. sabini*)

5975. Palm Swift. ♀. ovary slightly enlarged.

*Tachymenoides parvus*. Stomach contained many winged ants and a small beetle. Practically none noticed during the trip down the Aruwimi.

5976. Green Sunbird. ♂. testes somewhat enlarged, *Cyanomitra obscurus*. skull not quite completely ossified. Stomach contained 2 hard round seeds from berries.

5977. Parrot. (*A. zenkeri*). ♂. testes slightly enlarged. Iris orange. Crop contained small seeds.

One of a flock of 5 or 6, that came to feed in a wild fig tree.



Bengamisa

Sept 28

5974. 1

Sept 30

5979. 1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

## Bengamisa

1180

Sept. 29. 1914.

5978. Dark gray Cuckoo. ♂. t. n. e. S.C.: caterpillars.

Sept. 30. (Kaparata - Stanleyville.) Hirundo nigrita

5979. Dark blue Swallow. ♀. o. n. e. Common along this part of the Lindi, just now they are almost invariably seen in pairs.

5980. Gray Flycatcher. ♀. o. n. e. Common along this part of R. Lindi.

5981. Small gray Warbler. ♀. ovary somewhat enlarged. Stomach contained 2 small caterpillars and remains of insects.

The song of this little warbler was frequently heard along the Aruwimi, but usually they are more or less hidden by the foliage, and the boat goes by so quickly that one scarcely gets a chance to secure the bird, without a special effort.

Stanleyville

Oct. 25.

5982 Sunbird. ♀. ovary slightly enlarged. <sup>515894</sup> Anthreptes orientalis <sup>(Gmelin)</sup>. Preening itself, in a dry tree near river bank.

Cited as *redula*  
5983 Brownish Grass-Warbler. ♂. testes slightly enlarged. Iris light brown, bill black, slightly grayish below; feet buffy pink, claws dusky.

This is evidently the species common about posts in the Ituri. It is common here, too, but there is also a similar species (collected in 1909). <sup>(in voice as well)</sup> with streaked black, resembling

Stanleyville  
Oct. 25th 1914.

Stanley  
Oct. 25th

the swamp-loving species of the Vele, but here not strictly confined to wet situations. The two species here have somewhat similar notes, mostly dry, trilling efforts, but the one with the streaked back often <sup>changes</sup> ~~finds~~ up with a "chiv-chiv-chiv..." which is distinctive.

5984 <sup>♂</sup> Black-crowned Waxbill, with blackish belly. ♀. ovary slightly enlarged, skull not completely ossified. Iris dark brown, bill black, with a triangular spot of light crimson on each side of mandible near base; feet blackish.

Hitherto I had never noticed this form except at Avahubi; but here at Stanleyville it is rather common, the less numerous than the white-bellied one, with which it may be found in company. Both are practically voiceless, feeding in small companies, picking seeds from the tall grasses, when they display great talents in balancing, or hopping on the ground.

5985 <sup>♂</sup> Yellow- & black Weaver (*Pyromelana*?)  
♂. testes enlarged. Iris dark brown, bill black above, shading to dark greenish-gray

Stanleyville.

Oct. 25. 1914.

beneath mandible; feet brown. Stomach contained small seeds.

5986. Same as preceding. ♂. t. e. S.C.: small seeds.

<sup>Pyromela cana</sup>  
Fairly common in the high grass along roads and in old clearings hereabouts. At present they seem to be breeding, and go in pairs, the male very attentive to his dull-colored mate, and making short flights over the grass, that remind one of the similar performance of "Bishop-birds" (the more rapid). But the wings appear to be kept somewhat above the horizontal, and are often beaten so forcibly that it can be heard <sup>(the faintly)</sup> for a dozen yards. While perching the male seems a pretty bird - much more so than when in the hand — on account of the contrast of its black and yellow plumage, the feathers of the rump being slightly raised so as to show conspicuously.

Nov. 1. 5987. Barbet (*Gymnobucco*) ♂. testes slightly enlarged. Iris scarlet; skin of cheeks, the rest entirely naked dark brownish pink or dull red-brown, bill dusky-brown, feet dusky-brown. Stomach contained berries.

# Stanleyville

Nov. 1. 1914.

*Gymnobucco bonapartei*

5988. Same as preceding ♂ testes slightly enlarged

Iris reddish brown, scarlet above pupil.

Stomach contained berries.

*Gymnobucco bonapartei*  
5989. Same. Testes slightly enlarged. Iris

reddish brown, scarlet above. Stomach

contained pieces of a bug.

Very common at Stanleyville,

frequenting the trees, where they

cut the dead limbs.

Comparatively

I know, these birds

almost be called

5990. Warbler (*Sylviola*)

enlarged.

5991. Slender-billed "We

enlarged. Stomach con-

(I noticed a few testes)

not appear to be com-

mon. 5992. Bishop-Bird. *Pycnonotus*

" " ♂ t

5993 " " ♂ t

5994 " " ♂ t

Found in the es-

tall grass near the

station; but I have

generally distributed

5995. Pyrenestes. ♂ over

Iris dark red, eyelids pale blue above &

nov. 8. *Pyrenestes*. ♂ over

Iris dark red, eyelids pale blue above &

No 5985 has been seen at Bonai in a  
green marsh. At Matadi we saw  
what were probably Colius passer, in  
dull plumage, & Videa also occurring  
in the Bas-Congo, molting into breeding  
plumage towards the middle of January



## Stanleyville

Nov. 1. 1914.

*Gymnobucco bonapartei*

5988. Same as preceding ♂. testes slightly enlarged  
Iris reddish brown, scarlet above pupil.  
Stomach contained berries.

*Gymnobucco bonapartei*

5989. Same. ♀. ovary slightly enlarged. Iris  
reddish brown, scarlet above. Stomach  
contained berries and pieces of a bug.

Very common near Stanleyville,  
particularly frequenting the larger trees, where they  
cut their nests in the dead limbs.

Compared to most of the other barbets  
I know, these brown ones might  
almost be called spuds. ~~that~~

5990. Warbler (*Sylviella*). ♀. ovary somewhat  
enlarged.

*Nigrita fusconota*

5991. Slender-billed "Weaver". ♂. testes slightly  
enlarged. Stomach contained small fruits.  
(Tho noticed a few times at Avalubi, did  
not appear to be common there)

Nov. 5. 5992. Bishop-Bird. *Pogonornis flammiceps*

5992. Bishop-Bird. ♂. testes much enlarged.

5993. " " " ♂ t.e. Crop contained green rice

5994. " " " ♂. t.e.

Found in the extensive areas of  
tall grass near the "lazaret" N of the  
station; but I hardly think it is  
generally distributed here.

- Nov. 8. 5995. Pyrenestes. ♀. ovary somewhat enlarged  
Iris dark red, eyelids pale blue above &

Stanleyville  
Nov. 8 1914.

below, black before & behind eye; bill dusky bluish, somewhat lighter on upper side of maxilla; feet greenish-brown. Crop and stomach contained seeds, rather soft ones; indeed I have never been able to understand, from its food, why this bird should have such a remarkable development of the jaws.

This <sup>example</sup> seems to be of the big-billed kind.

5994.

v. de Villegas showed  
big variety he had

*lucida subalaris*  
*longirostris*, ♀ imm.

feeding everywhere,  
ad trees, as well as  
sts in the rivers. A  
thered one rainy

Dec. 10 - 1914  
Today at Yarouge a ♂ of Pyrocephala  
feathering was seen, in full  
breeding plumage, near Bumba  
we saw again what seemed to be

examples of this species, but all  
in brown plumage. On late  
Dec. & early January, too, in  
the Kas Congo, they are all  
in eclipse plumage. The <sup>Pyrocephala</sup> ~~Pyrocephala~~  
is molting. There seems to be  
no "imperfection" in the latter region (over)

giving its flight  
never observed anywhere  
about Aduku

the Kas Congo, they are all  
in eclipse plumage. The <sup>Pyrocephala</sup> ~~Pyrocephala~~  
is molting. There seems to be  
no "imperfection" in the latter region (over)  
the Kas Congo, they are all  
in eclipse plumage. The <sup>Pyrocephala</sup> ~~Pyrocephala~~  
is molting. There seems to be  
no "imperfection" in the latter region (over)

Stanleyville

Nov. 8 1914.

below, black before & behind eye; bill dusky bluish, somewhat lighter on upper side of maxilla; feet greenish-brown. Crop and stomach contained seeds, rather soft ones; indeed I have never been able to understand, from its food, why this bird should have such a remarkable development of the jaws.

This <sup>example</sup> seems to be of the big-billed kind, but at Panga M. de Villegas showed men of the small-billed variety he had shot.

5996. Swallow (*H. lucida subalaris*) ♀ imm.

Very common here, feeding everywhere, and resting on dead trees, as well as on rocks and posts in the rivers. A large number gathered one rainy morning to catch termites flying up behind our house.

No. Nov. 22. *H. lucida rotundirostris*  
is of course common at Stanleyville  
nowhere.

5997. Rufous-winged Shrike. ♀ imm. Iris dark brown. Bill blackish above, shading to gray beneath mandible; feet light blue. Stomach contained remains of insects, including orthoptera.

Fairly common amid the grass and brush here, giving its flight song occasionally. never observed anywhere about Stanleyville

Stanleyville

Nov. 8. 1914.



5998. *Spermophilus* <sup>griseata</sup> ♀. ovary slightly enlarged.  
skull not quite entirely ossified (transparent spots persisting in parietal region) Iris brown, of a medium tint, rim of eyelids pale-blue (not swollen); bill pearly blue, shading to scarlet along cutting edges and on tip of mandible; feet brownish-green. Stomach contents: seeds.

Shot among grass & bushes near a small brook.

5999. *Piculetus* ♂. t. n.e. Stomach contained 6 white "grubs". Was climbing about a dead limb in a rather small tree out in an open patch of tall grass.

6000. *Cossypha cyanocampter*. ♂. testes somewhat enlarged. Stomach contained a spider, a small caterpillar, and pieces of insects. in dense second growth — there were 3 singing within a circle of 15 yds diameter.

6001. Warbler (*Cisticola*) ♂. im. Iris light grayish-brown, bill dark brown above, shading to buffy-yellow below; interior of mouth mostly yellowish, but base of tongue black above, and a large black

area on roof of throat (palate yellow save for posterior edge), feet pale pinkish-brown.

6002. Flycatcher (*Muscicapa grisola*). ♂.

t.v.e. First seen this season.

A red-backed shrike was also observed today. <sup>See No. 6006.</sup> The bank swallow was first seen on its return from north, Sept 5 (Avahubi). The Swallow, Sept 29, the blue-headed yellow wagtail Oct 14.

As for African birds - a flock of 5 storks (*Abdimia*) was seen circling overhead today. No "merles metalliques" have been seen here during the past month, tho we saw some in Aug 09; very probably they disappear as at Avahubi.

*Pycnonotus afer*

6003. Weaver. ♀ im. Stomach contained grass-seeds. S. 5986.

6004. Warbler (*Parisoma?*) ♂ im. <sup>plumbeous</sup> Stomach contained a small caterpillar and insect remains.

6005. Brownish Grass-Warbler, with streaked back. *Cisticola lugubris*

♀. Ovary slightly enlarged. Iris light brown, bill dark brown along culmen, shading to light gray below; feet pinkish-buff. Common.

Stanleyville.

Nov. 14 1914.

6006. *Erneocorax collaris*. Red-backed Shrike. ♀. o.w.e. Iris brown, bill dark brown above and at tip, shading to light gray on base of mandible; feet dusky greenish.
6007. *Pomatorhinus auricapillus*. Rufous-winged Shrike. ♀. ovary slightly enlarged. Iris dark purplish-brown, with inner rim much lighter & grayish; bill black, feet light blue. An adult male examined today (too badly shot to save) had the eye of exactly the same color. Utters a short "chuck!", repeated at intervals, from concealment in bushes.
6008. *Philemonia indicata*. White-eyed Bulbul. ♂. t. n. e. Iris white, bill black; feet dark bluish-gray. Skull not completely ossified. Was giving a rather weak whistled note of 2 syllables, very different from its ordinary voice.
- Nov. 15. 6009. Weaver. ♂. testis slightly enlarged. Stomach contained grass-seeds and small stones. *Ptyonoprogne labradorica*. Commencing post-nuptial molt — the new feathers, the small, show plainly that the bird in eclipse plumage is streaky brown above and white & buff below, all the black feathers, and even the yellow rump-patch, disappearing entirely.



area on roof of throat (palate yellow save for posterior edge), feet pale pinkish-brown.

6002. Flycatcher (*Muscicapa grisola*). ♂.

t.v.e. First seen this season.

A red-backed shrike was also observed today. <sup>See No. 6006.</sup> The bank swallow was first seen on its return from north, Sept 5 (Avahubi), the Swallow, Sept 29, the blue-headed yellow wagtail Oct 14.

As for African birds - a flock of 5 storks (*Abdimia*) was seen circling overhead today. No "merles metalliques" have been seen here during the past month, tho we saw some in Aug 09; very probably they disappear as at Avahubi.

*Pyromelana afra*

6003. Weaver. ♀ im. Stomach contained grass-seeds. S. 5986.

6004. Warbler (*Parisoma?*) ♂ im. <sup>plumbeum</sup> Stomach contained a small caterpillar and insect-remains.

6005. Brownish Grass-Warbler, with streaked back.

*Cisticola lugubris*  
♀. Ovary slightly enlarged. Iris light brown, bill dark brown along culmen, shading to light gray below; feet pinkish-buff. Common.

Stanleyville.  
Nov. 22. 1914.

no heads, legs or wings.

A nest of this species of sunbird, found today in a rather large tree along a road, at a height of 30 feet, was of typical shape, composed of fine strands of dry down vegetable material & pieces of dry leaves, lined with plant-down, and decorated externally with pieces of bark, lichen & caterpillar excrement. This nest contained 2 young, with eyes still closed, bill pink, gape pale yellow, feather sheaths still very small, and dusky. No down whatever. Both parents were seen in the tree, the female entering the nest.

6015. Sandpiper (Gallinago) ♀. o.n.e.

6016. Large Snipe. ♀. o.n.e. Stomach contents.

s. 6011. Gallinago media.

(16 minutes) several small stones, remains of insects (?), and a small white maggot-like insect larva.

6017. Snipe (G. gallinago) ♀. o.n.e. Stomach contents: insect-remains.

6018. Same as preceding ♀. o.n.e. Stomach contained Gallinago gallinago a few small stones and remains of arthropods.

Nos 6016-18 were put up singly in a small open marsh in a pasture. Nos 6015

& 6016 were quite fat, but 6017-18 decidedly lean.

6019. White bellied Starling (Cinnyris inclusus l. leucogaster) [♂] not seen since leaving.

6020 Weaver Finch (Spermestes) [♂] <sup>nutata</sup> Same as 5998.

6021. Nightjar.

Nos 6019-21, preserved by injection with formalin, and presented by Dr. Grossule, were collected at Stanleyville from 4 to 6 months ago. Nightjars must be scarce, I have not yet

seen one.

Nov. 23. '14.

6022. Brown-cheeked Woodpecker. ♀. ovary slightly enlarged. Stomach contained ants and their larvae.

6023. Crow. ♀. o.v.e. Iris dark brown, bill & feet black. Common about Stanleyville, but do not come to roost every night in the station, as formerly. At sunset, on Oct 16, however, a flock of 35 was noticed; pairs are far more usual. Their note is a rather low, hoarse, and slightly prolonged "kāhk". I have seen one pursued by drongos. So they may sometimes be approached to within 10 yards when on the ground, they are usually shyer. Up to the time of our arrival here, none had been seen since Niagara.

Nov. 24

6024. ~~Circus cyaneus~~. ♂. testes slightly enlarged.  
*Dryotis bairdii* Stomach contained scales of a snake.

Nov. 27.

6025. Reed-Warbler (*Calamocichla*) ♂ testes slightly enlarged, skull not completely ossified. Iris rather light-brown, bill dusky-brown, shading to buffy-gray on base of mandible, feet rather light bluish-gray. (See note under no. 6034.)

6026. Grass-Warbler (*Cisticola*) <sup>fuscus</sup> ♂. testes slightly enlarged. Stomach contained a green

Stanleyville  
Nov. 27. 1914

Sta  
Nov

grasshopper and remains of other insects.

6027. *Paradise Flycatcher* ♂. testes slightly enlarged.  
*Terpsiphone viridis*

6031.

Preserved with formalin, presented by Dr Grossule.  
(Of the median rectrices, the right one is about  
7 inches long and pure black, the left one  
is about 4 inches, black with a little rufous  
at tip; back rufous.)

6032

6028. *Paradise Flycatcher* ♀. ovary slightly enlarged.  
*Terpsiphone viridis*

Preserved with formalin, presented by Dr Grossule.

Nov. 29. *Terpsiphone viridis*

6029. *Paradise Flycatcher* ♂ t. n. e. Iris dark  
brown; bill, eyelids, & feet blue, the first-  
mentioned with a little black at tip;  
interior of mouth light yellowish green.  
In the midst of the moult, with middle  
tail-feathers just sprouting, but lack  
wholly white.

6030. *Paradise Flycatcher* ♂. testes much enlarged.  
*Terpsiphone viridis*

Bill bright blue, black at very tip; iris  
dark brown; eyelids & feet bright blue;  
whole interior of mouth greenish yellow.

Comparing nos 6029 and 6030 while  
fresh, I found the blue of bill, eyelids  
and feet somewhat brighter in the  
latter, and the color of the interior of  
the mouth was likewise brighter.



Stanleyville  
Nov. 29. 1914.

6031. *Cossypha verticalis*, ♂ testes slightly enlarged  
skull not completely ossified. Iris dark  
brown, bill black, feet purplish-gray.  
Stomach contents: remains of a small  
round millipede.
6032. Same as preceding. ♂ testes slightly  
enlarged, skull not quite fully ossified.  
Stomach contained remains of rather  
small beetles.

Three were singing this morning  
in a small dense patch of bushes  
and trees, near a swampy creek,  
but surrounded <sup>on the other side</sup> by tall grass.

*Cossypha cyanocampter* of course  
inhabits the dense second growth so  
extensive about Stanleyville, but  
one example of *C. verticalis* had  
<sup>hylangutia</sup> already been seen several times  
near the cemetery and among the  
coffee trees close to the station. This  
latter species has here been heard  
to mimic *Francolinus squamatus*,  
*Merops persicus*, *Asturina* & *Turdus*.  
Its singing is <sup>far much</sup> louder than, but hardly  
so rich in tone as that of *C. cyanocampter*.



193  
Stanleyville  
Nov 29-1914.

6033. Sunbird ♂. testes slightly enlarged.  
S. 6014.

Skull not quite fully ossified. Stomach contained a spider and a small caterpillar

6034. Warbler (Calanocichla) ♂. testes slightly enlarged.  
S. 6025.

Iris rather light brown, rim of eyelids light greenish; maxilla dusky brown, mandible shading from grayish at tip to buffy at base, gape light yellow, feet rather light gray, interior of mouth yellow.

One or two of these birds, whose voice is exactly the same as those of Faradje & Parava, <sup>originally</sup> haunted some tall grass (not matete) on dry ground, but close to a creek. This peculiar guttural "churr" was heard once at Banalia.

6035. Skeleton of Black crested Eagle (Lophoetus)

♀. right ovary very much enlarged (laying),

Yanonge left one very small. Crop contained 3 rats.  
Dec 10. 14.

6036. Paradise Fly catcher. ♂. t. e. Iris dark brown, bill and eyelids bright blue.

<sup>Serpinkhae viridis</sup> 6044  
6037. Paradise Fly catcher ♂. t. e.

Shot close together. ♂ color of back simply a question of age?

Isangi 6038. Pyrenestes ♂. t. w. e. A rather large flock was going to roost in high grass <sup>outcaves</sup> in rubber plantation. Another male



Sam  
Dec 16

6039. B.

6040. B.

6041. B.

6042. C.

Bamambwa  
Dec 11. 14  
6043.

6044.

6045.

6046.

6047.

Sangi

Dec 10. 1914

was shot & had likewise small beak.

6039. Barbet (*Pricholaema*) ♂ t.m.e. S.C.: fruit.

6040. Long-tailed Sunbird <sup>*aneognathus*</sup> ♂ t.e. skull not fully ossified. *Nectarinia congoensis*. Iris dark brown, bill and feet black. Stomach contained remains of small spiders.

6041. Same. ♀ o.m.e. skull not fully ossified.

*Nectarinia congoensis* Iris dark brown, bill brownish-black, feet black. Stomach contained remains of small spiders.  
Nos 6040 & 1 were feeding at the brilliant red flowers of an ornamental tree known here as the "*Poinciana regia*" "flamboyant"

6042. Coucal. ♀ o.m.e. S.C.: grasshoppers & *Centropus anselli* crickets, larvae of wasps, a caterpillar, snails (with their soft shells torn off), a few slugs. Iris reddish-brown, bill

*Bambu* blackish; feet dusky.

6043. Long-tailed Sunbird. ♂ t.e. S.C. tiny spiders

6044. " <sup>*Nectarinia congoensis*</sup> ♂ .. .. ..

n. Baobabs.

6045. Thrush (*Alectro*) ♂ imm. S.C. insects & no woosname bones of tiny frog.

6046. Small Plover. ♂ t.m.e. Iris very dark

*Charadrius marginatus pallidus*. Brown, bill black, with a little greenish-gray at base of mandible, feet light greenish-gray, slightly buffy on metatarsi

6047. Same, ♂ t.m.e.

Near Lie.  
Dec 12. 1914

- Sitagra* (*meanscephala*?)  
6048 A Yellow Weavers with black head ♂.  
t. m. e. Iris brown; bill black, shading  
to greenish-gray on base of mandible;  
feet brownish-pink. S.C.: small seeds.  
In long grass on edge of an island.
6049. Long tailed Sunbird ♂ t. e. A  
*nestchiria conspicua*. sunbird nest, probably built by this  
species, was hanging from a bush,  
over the water. A ♂ sunbird was  
near it.

nowelle Anvers - Dec 14.

6050. Small Plover, ♀ ovary slightly enlarged.  
*Charadrius m. pallidus*.  
S.C.: remains of insects, including many  
heads of termites.

6051. Weavers (*Spermophilopsa*) ♀ imm. Iris

no. 6051.

Cog  
60

60



Interior of mouth yellow, 3  
black spots on palate, outer ones  
large, posterior pair lacking.  
a black crescent beneath  
tongue, on mandible; no  
mottles on tongue.

*Spermophilopsa* ♀ imm.

Nowelle Anvers, Dec 14. 1914.

Not seen above Cogilhaterville, the from,  
there on along the river they were very



Near Lie.  
Dec 12. 1914

Cogui  
Dec 1

*Sitagra* (*meioscephala*?)

6048 A Yellow Weavers with black head ♂.

t. w. e. Iris brown; bill black, shading to greenish-gray on base of mandible; feet brownish-pink. S.C.: small seeds.

In long grass on edge of an island.

6049 Long-tailed Sunbird ♂ t. e. A

*nestchuria conspicua*

sunbird nest, probably built by this species, was hanging from a bush over the water. A ♂ sunbird was

near it.

nowelle Anvers - Dec 14.

6050 Small Plover, ♀. ovary slightly enlarged.

*Charadrius m. pallidus*

S.C.: remains of insects, including many heads of termites.

6051. Weavers (*Spermophilopsa*) ♀ inv. Iris

brown; rim of eyelids pale bluish;

feet olive.

Coguihatville. Dec 15.

6052. Warbler (*Cisticola*) ♂. testis slightly enlarged.

Found all along the Congo, down to

Boma.

*Tylorhoes erythronotus*

6053. Red-rumped Barbatula. ♂. testis slightly enlarged. S.C.: fruit.

6054 Swallow (*Riparia*) ♀. ovary slightly enlarged

*Riparia riparia*

6055 *Riparia riparia* .. ♀ ..

6056. *Riparia riparia* .. ♀. o.w.e. Iris dark brown; bill black, feet dusky brown.

Not seen above Coguihatville, the from there on along the river they were very

com

6057. W

med

gr

no

Be

w

6058. mi

d

r

Cogui

6059. mi

6060. mi

# Coquilhatville

Dec 15, 1914.

common, down to Boma.

6057. Weaver (*Vidua*) ♂. t.e. <sup>area</sup> skull, as usual, not completely ossified. Crop contained small grass-seeds. (A male with long tail also noticed today. During my stay at Boma, no males in breeding plumage were seen.)

6058. Long-tailed Sunbird ♂. t.e. S.C. remains <sup>*nectarina* *cyanura*</sup> of tiny spiders. A common bird about villages and posts and in trees along river-bank; but I have not heard any note from it.

Coquilhatville Dec 16,

6059. Rose-breasted Bee-eater ♂. t.e.

coquilhatville Dec 16, 1914 (*Wangita*)

A blackish swallow-like bird seen flying about here also, <sup>over village</sup> abt as large as a martin perhaps, seemed to have something red on front of head. Not a *Psalidoprocne*.

*Pseudochelidon erythromelas*

See 6060

A number were flying about, with other swallows, over the river and a native village. not noticed again.

6061. Weaver (*Nigrita*). ♂. t.e.



Inches

# Coquilhatville

Dec 15, 1914.

common, down to Boma.

6057. Weaver (*Vidua*) ♂. t.e. skull, as usual, not completely ossified. Crop contained small grass seeds. (A male with long tail also noticed today. During my stay at Boma, no males in breeding plumage were seen.)


6058. Long-tailed Sunbird ♂. t.e. S.C. remains nectarina cyanura of tiny spiders. A common bird about villages and posts and in trees along river-bank; but I have not heard any note from it.

Coquilhatville. Dec 16.  
6059. Rose-breasted Bee-eater. ♂. t.m.e. Merops malimbicus Iris bright crimson, bill black, feet dusky-brown.

Seen first near Bumba, common in flocks of 30 or more all along river, many seen as far down as Kunzulu, near Kwamouth. The voice is of the usual character, too, and rather loud, but hoarser than usual. ("chick-a-l")

6060. Small brownish Swallow. ♂. t.e. A number were flying about, with other swallows, over the river and a native village. Not noticed again.

6061. Weaver (*Nigrita*). ♂. t.e.



Near Coquillatville

Dec 16. 1914

Lake  
Dec 17.

6062 Small Plover ♂ t. m. e. SC: insects  
*Charadrius marginatus pallidus*.  
6063 " " t. m. e.

6064 " " ♀ o. n. e.  
going in small parties  
a common species, found of river banks,  
villages, etc. Like its relatives, active  
at dusk.

6065 Long-tailed Sunbird. ♂. Testes somewhat  
*nectarinia congrua*. enlarged. S.C. remains of small spiders.

Iribu - Dec 17. *Hirundo puebla*

6066. Striped-breasted Swallow. ♂. Testes slightly enlarged  
lady-birds and  
1 bug. Two

flying about on  
dark blue swallows  
along the river

Kinshassa. I have not seen  
species with a  
frequenting it  
but it is said  
between Bon

6067 Long-tailed  
*nectarinia congrua*

6068. Mate of 6067

Lubolela ♂

6069. Horned Owl

*Bubo lunatus*

yellow, bill black

approaching yellow

inches

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

No. 6072. Iris gray; practically same shade as feathers of back of head; bill yellow; blue streak of face unpaired, orange around side of posterior border, that of throat dirty flesh-color tinged with yellow, feet pale pinkish gray, claws shading to yellow on tip, close to feathers, and rose to dusky toward tip, nails black.

6072. G

g

b

b

w

c

a

t

b

1

1

1

Near Coquillataville

Dec 16. 1914

Culic.  
Dec 17

6062 Small Plover ♂ t. m. e. S.C.: insects  
*Charadrius marginatus pallidus.*  
6063. " " t. m. e. "

6064 " " ♀ o. n. e. going in small parties  
a common species, found of river banks,  
villages, etc. Like its relatives, active  
at dusk.

6065. Long-tailed Sunbird ♂. Testes somewhat  
*nectarinia congoensis.*  
enlarged. S.C. remains of small spiders.

Irebun - Dec 17. *Hirundo puebla*

6066. Stresemann's breasted Swallow ♂. Testes

slightly enlarged, shall not completely  
ossified. S.C. many small bees, some  
lady-birds and other small beetles +  
1 bug.

Two of these swallows were  
flying about over the water. The

dark blue swallow is found all  
along the river as far down as  
Kinshassa. At Boma and Matadi  
I have not seen it; a rehorn-crowned  
species with slender outer rectrices  
frequenting the river banks there;  
but it is said (Drouaise) to occur  
between Boma & Banana.

6067. Long-tailed Sunbird ♂. t. e. S.C. a  
*nectarinia congoensis.* tiny fish  
etc.

6068. Mate of 6067 ♀. ovary somewhat enlarged. S.C. tiny  
Culicoides Dec 17. species.

6069. Horned Owl ♂. t. m. e. Iris chrome-

*Bubo lunatus.* yellow, bill light grayish green,  
approaching yellow distally; cere



Lukolela

Dec 17. 1914.

greenish buff; toes yellowish; claws black with gray bases.

Sitting in tall tree at dusk on

edge of forest

Bolobo. ~~This 18~~ <sup>citroga rufa</sup> 6070. Tiny brown Warbler, ♂ t. sl. e. SC immots


6071. " " " ♂ . . . .

Iris light brown. Bill blackish, greenish gray beneath, basally; feet pale buff.

Everywhere common in lower Congo, flying high in the air, often spirally.

[Today for the first time we began to see large patches of open brush country, and in a short walk at Bolobo at sundown I was surprised to see how the general aspect of the vegetation resembled that of the Nile, while the few birds we saw, <sup>and heard,</sup> greatly strengthened this impression].

6072. Pantalus. ♀ imm. Iris gray, bill <sup>This ibis</sup> yellow; bare skin of face orange red, but orange around posterior border, throat still duller; feet pale pinkish gray, yellow on tibiae, dusky on tips of toes. (Captive specimen, presented by Dr. Girling.) Only



Bolobo

Dec 18. 1914

one example seen from the steamer  
on trip down the Congo, a beautiful  
adult, on Dec 13 th.

Kwamouth Dec 19.

*Trochocercus torquatus*

6073. Bulbul with black breast band. ♂. testes  
slightly enlarged. Iris very dark brown,  
bill & feet black.

6074. Same. ♂ imm. S.C. fruit.

6075. " ♀ imm. " "

Found in open brush country. Have  
a sort of twittering note. A nest  
of this species, found this afternoon  
at Kungulu was placed low  
down (4 ft) in a bush, a frail  
cup of slender grass- & plant stems,  
containing 2 eggs, pinkish white,  
indistinctly speckled with darker  
pinkish, and a faint rufous zone  
about larger end.

6076. Large brown Warbler. ♂. testes slightly  
enlarged. Common at Kwamouth  
& Leopoldville. Has same song as  
in Nelle


*Thamnophilus nigra*

6077. Black Thrush with white shoulder  
patches. ♂. t.w.e.

*Cichladusa lateralis*

6078. Warbler ♂. testes slightly enlarged.

Common, Bolobo. Kwamouth, Boma,  
etc. Song same as in Nelle.



Kwa  
Dec

6079

L

v

x

C

N

s

Kung

6081

6082

6083

# Kwamouth

Dec. 19. 1914.

*Batis helleri congoensis*

6079. Flycatcher. ♀ imm. Common at Leo & Boma. Beats its wings sometimes with a slight noise, utters short weak whistles.

*Lagonosticta conica - Sharpe*

6080. Small reddish "Weaver". ♀. ovary slightly enlarged.

6081. Large slaty Flycatcher. ♀ imm.  
*melanornis ater*


Kunzulu.

6082. Coucal. ♂. t. n. e., Iris scarlet, bill black, feet rather dark blue-gray.  
*Cuculus senegalensis*  
Stomach contents: remains of insects (orthoptera, coleoptera) and a caterpillar.

At Kunzulu the brush was rather tall, with some small trees, especially back from the river, but in general, from Kwamouth to Boma, the bushes were sparser and more stunted, the country far more barren and apparently infertile than in the rule. This tendency is most pronounced at Boma; where many of the hills have practically no brush, only coarse grass, with here or there an occasional bush.

Kinsassa - Dec 20.

6083. Palm Swift. ♂. testes somewhat enlarged.  
Flies into a lighted room at night. Common





Kinshassa  
Dec 20, 1914.

everywhere in lower Congo.

Leopoldville. Dec 21.

6084. Black Titmouse with white on wing. ♂ im.  
*Melaniparus niger insignis*  
S.C.: caterpillars. Rather common here.

6085. Gray Titmouse with pale rufous belly. ♀ im.  
*Melaniparus nigerinus*  
*Iris brownish-gray*, bill black, feet  
light blue. Stomach contained caterpillars.

6086. *Sylvia* ♀ im. Iris light brown. In  
open brush.

6087. *Cisticola natalensis strangei*  
Stout-billed grass-Warbler. ♂ testes  
slightly enlarged. S.C. orthopterous insects  
Common in lower Congo. Habits &  
notes same as in Nelle.

6088. Nightjar. ♂ im. *Caprimulgus natalensis*

6089. Sunbird ♂ t.e. Iris pale brown, bill &

6090. " " ♂ " " foot black.

Common. *mirafra leichori*

6091. Clapper-Lark. ♂. testes small, skull not  
quite completely ossified.

6092. Thrush (*Erythropygia*) ♂. t.w.e.

Leopoldville Dec 22.

6093. Lapwing. ♀ ovary slightly enlarged.  
*Stephanius infinitus*  
Iris yellow, with greenish-gray inner  
rim; bill & feet blackish. S.C.: insects.

6094. Same ♂ im, Iris chrome-yellow, with  
greenish-gray inner rim, bill & feet blackish.  
S.C.: remains of insects (including beetles).

Leopoldville  
Dec 22. 1914.

Walking about in pairs or parties  
of 4 or 5 on cultivated ground.

Have a short complaining call  
when flushed.

*Brachypteryx maximus remissa*

6095. Large brownish Flycatcher. ♂. t. n. e.  
Iris dark brown, bill & feet blackish.

6096. *Estrilda cinnabrinova clarkei*.  
Waxbill. ♂. t. n. e. In flocks, feeding  
on ground, and alighting occasionally  
on grass. Common also at Boma.

6097. Greenish Bullbul. ♂. t. e. *Audropygia*

6098. *Parisoma* <sup>plumbeum</sup>, ♂. t. n. e.

Physville. Dec 23.


6099. Barbet (*Gymnobucco*) <sup>pili</sup> ♂. t. e. s.c. fruit.

6100. " " " ♂. t. e. " "

Iris dark brown, skin of face blackish,  
bill rather light rufous brown, feet  
dusky-brown.

In this region there are often  
large wooded areas, but not the  
tall equatorial forest. A number  
of these barbets were sitting on  
a large dead tree where they  
had cut their nests, exactly like  
their congeners in the Ituri, etc.

Many starlings, like no. 6101, were  
also perching there, and perhaps  
nest in the deserted holes of the barbets.



Thysville

Dec 23. 1914.

6101. Starling, ♂ *Poecoptera lugubris*. testes slightly enlarged.  
6102. " " ♂. testis somewhat enlarged; skull  
not quite completely ossified. Iris cadmium-  
yellow.

6103. Same. ♀ o. e. Iris bright yellow

6104. Bunting. ♀ imm. Crop & stomach contained  
*Trinoglossa tachys* small seed & a tiny grasshopper.

matadi

Dec 25. *Cichladusa ruficanda*

6105. Rufous-<sup>x</sup> gray Thrush. ♂. testes  
slightly enlarged. Iris rather light  
red-brown; bill blackish; feet rather  
light bluish. Stomach contents: remains  
of a spider & insects.

First seen at Kinshassa, a  
common bird in the Bas-Congo.  
Lives in pairs or small family parties,  
is rather arboreal in its habits, but  
may also be seen low down in  
bushes, and does not shun the  
neighborhood of houses. Has some  
rich melodious notes, but they  
are not very long, or else simply  
repeated; and as in some species  
of *Cossypha* they are apt to be  
given louder towards dusk.

matadi

Dec 25. 1914.

6106. Finch (*Serinus*) <sup>moyambicus tando</sup> *t. testis* slightly enlarged. Iris brown, bill dusky brownish paler below, feet brownish. Common at Kivamouth, Leo, & Boma. Resembles its relative of the Nile in general habits as well as in its slightly musical cheeping notes.

6107. Swift (*Cypselus*) with white throat and rump <sup>*affinis*</sup> *t. t. e.* Iris dark brown, bill black, toes dusky brown. Stomach contained remains of winged ants. Common about rocky hills at matadi. Along the river above matadi there is a high, overhanging, cliff, beneath the shelter of which are two large clusters of nests, evidently of these swifts, which frequent the spot in numbers, but too far away for me to be sure. It reminded me very much, however, of the colony at Piaga, near Faradje.

Dec 26.

6108. Pytilia <sup>*melba* (L.)</sup> *A. t. m. e.* Iris light yellowish-brown, bill rose-color, feet grayish brown. Stomach contained heads of termites.
- Pytilia <sup>*melba* (L.)</sup> *A. Same. T. e. m. e.* Iris light yellowish brown. Bill dusky brown above, shading to rose.

Matadi

Dec 26, 1914.



at sides and below; feet grayish-brown.  
Stomach contained remains of insects,  
including termites. Found on the  
barren hills about Matadi. Seen once  
near Boma.

6110. Bunting. ♂. t. n. e. Iris dark brown,  
<sup>anisognathus</sup> bill blackish above, buff below, and  
yellow at base of mandible & gape; feet  
buffy-brown.
6111. Same. ♂. testes slightly enlarged. S.C.: seeds
6112. ♀. o. n. e. "

Very common at Matadi, even hopping  
about like sparrows in streets and  
yards. I have seen a dozen on a  
roof at a time, but usually they are  
in smaller numbers. They are  
also found on the hills, along the  
railway tracks — in short, almost  
everywhere. (Passer griseus has not  
is not common in the Bas-Congo)

*Laniotrochus lanioides*, Sharpe

6113. A small reddish "Weaver". ♂. t. n. e.

Iris brown, maxilla dark gray, mandible  
pale rose with dusky tip, feet dusky.  
Stomach contents: small seeds.

*Cagnotrochus congoensis*.

6114. A "Weaver". ♂ imm. Iris brown, bill blue,  
shading to black at tip. feet dark bluish.  
Crop filled with small seeds.

matadi.

Dec 26, 1914.

*Cichladusa ruficanda*

6115. Thrush ♀ imm. Iris light brown.  
5. 6105. bill dusky brown; feet blue. S.C. insects


6116. Brown Nightjar ♀. ovary slightly  
~~Caprimulgus feridus~~ enlarged. Iris dark brown, rim of eyelids  
light brown, bill brown, shading to  
black at tip, feet dark brown. Stomach  
contained a small beetle.

Put up on the slope of a stony  
hill near matadi; flew a little  
way, lit on a rock, and then began  
to flutter about as tho wounded.  
at the spot where she had taken  
wing crouched her downy young one  
no 6118 was found on the same  
hill, and the day following,  
on the high rocky hill called  
"Loadi" south of matadi, 4  
goatsuckers, presumably of this  
same species, were observed

6107 Young of no 6116. ♀ juv. Iris dark  
~~C. feridus~~ brown, bill brown at base, black at  
tip; feet gray.

6118. Nightjar. ♀ imm. (*C. fossii*)

Dec. 27. Kingfisher. ♂ imm. Iris dark brown,  
~~Haleyia alluvialis orientalis~~ rim of eyelids brownish, bill dirty  
scarlet, blackish near tip, but pale  
dull orange at very point; feet orange



Matadi

Dec 27. 1914.



red. with all scales dusky brown.

S.C.: a mantis, a dragon fly, a grasshopper & a small snail

6120 <sup>Hypphantornis collaris</sup> Large Black & yellow Weaver. ♂. testes somewhat enlarged. Iris orange-red; bill black; feet pinkish brown. Crop and stomach contained palm-nut pulp & insect-remains. Shot from a tree where *Melanopteryx nigerrimus* was nesting. This yellow weaver seems to replace *H. cinnullatus* in the Bas-Congs, and is very common at Boma.

6121 <sup>Ploceus superciliosus (Shell.)</sup> Weaver, ♀. o. n. e. S.C. seeds & insects.

Seen from Kivamouth to Boma. In habits and notes same as in Ndele.

6122 <sup>Zosterops nelli</sup> <sup>l. l. t. melanurus</sup> Zosterops nelli melanurus. S.C.


6128. The note of this dove might be written: kū k-k-k-k-kūkū.

but sometimes the introductory part is doubled: kū kū k-k-k-k-kūkū

6123.

6124.

Matadi  
Dec 27. 1914.



red. with all scales dusky brown.

S.C.: a mantis, a dragon fly, a grasshopper & a small snail

6120 *Hypothymis collaris* <sup>Red</sup> Large Black & yellow Weaver. ♂. testes somewhat enlarged. Iris orange-red; bill black; feet pinkish brown. Crop and stomach contained palm nut pulp & insect-remains. Shot from a tree where *Melanopteryx nigerrimus* was nesting. This yellow weaver seems to replace *H. cucullatus* in the Bas-Congs, and is very common at Boma.

6121 *Ploceus superciliosus* (Shell.) <sup>Red</sup> Weaver. ♀. o.m.e. S.C. seeds & insects.

Seen from Kivamouth to Boma. In habits and notes same as in Uele.

6122 *Pytilia zonata* <sup>mella</sup> ♂. testes slightly enlarged. S.C.: remains of termites (workers, not winged individuals)

6123. Pipit. ♀ imm *Anthus leucophrys* (*angolensis*?)

6124. Gray Partridge (*Sternistes*) ♀. o.m.e. Iris dark brown; bill, feet, and naked skin of face & throat scarlet. Crop contained many small bulbs, O, numbers of ants, and a small snail. Plushed on a grassy hillside.

2. Partridges have been heard calling

matadi

Dec 27. 1914.

at sundown at Bolobo, Kungulu,  
and matadi. The note resembles  
somewhat that of the francolin of  
the Uele, but is perhaps not quite  
so harsh.

Dec 28 *Cagornisticus conica*  
6125 " Weaver". ♂. t.n.e. Crop contained  
tiny seeds.

6126 <sup>himba nubila</sup> Streak-breasted Swallow ♀ s.n.e.  
6127. . . . . ♀ . . .

Common at Matadi. Its old nests  
may be found beneath rocky ledges,  
but it has also learned to nest  
about houses, and nests may be  
seen, for example, beneath the  
eaves of the railway station.  
Also occurs at Boma, where the 2  
other species of brown-rumped  
swallows both are found.

Boma. Dec 31.

6128. Dove. ♂. t.n.e. Iris dark brown,  
eyelids gray; bill black; feet reddish-  
purple. Rather common about  
Boma. Has a low, hoarse, cooing  
note, that seems slightly different  
from that of birds in the Uele.

Other pigeons noticed here at  
Boma are *Vivago calva*, *Chalcophilia*  
*africana* (abundant) and the brown.



Borna

Dec 31. 1914.



collared pigeon (Streptopelia semitorquata)

Impelion rufocinerea

6129. Paradise Flycatcher. ♂ im. Iris

- dark brown, bill blackish brown,  
feet light blue.

Born

6135

In grove of trees, rubber trees &  
others, planted at agricultural  
station "Kalamu".

Dendropicos portoricensis

6130. Woodpecker ♀. o. n. e. Iris dark red

Cicadula ruficeps

6131. Thrush. ♂ im. S.C. insects & palm nut fibers

5. 6. 115. " 6132. "Babbler" (Crateropsg) ♀. o. n. e.

- Iris scarlet on outside, shading to  
bright yellow on inner edge, bill  
black, feet dark gray. S.C.: palm  
nut fibers, insect - remains.

Skulking in parties of 3 to 6  
among bushes & trees in hollows  
near water. Have, <sup>horse</sup> calling, &  
whining notes, but not so loud  
as either of the species of the Ule.

6133

6133. Sunbird ♂. testes somewhat enlarged

Cinnyris managuensis ~~delesserti~~

- Iris dark brown, bill & feet black.

S.C.: a ball of rubber & remains of  
winged termites.

6134

6134. Yellow Finch with black face & throat

Serinus capistratus

- ♂. testes somewhat enlarged. Iris rather

light brown, bill brownish above.

4 whitish below, feet pinkish-brown.

6135

Jan

6136

Boma  
Dec 31, 1914.

Commonly captured and kept in cages by blacks at Boma.

6135. *Cossypha (heuglini?)* ♂ t. m. e.  
skull almost completely ossified. Iris dark brown, bill black, feet violet-brown. Stomach filled with black-winged termites.


Common here; resembles in habits & song *C. verticalis*, and is apt to sing loudly at sunset. It certainly imitates other birds, this example having mimicked a bee-eater (*m. percicus*) very well. Probably the same species heard at Bolobs, Leo, Thysville & Matadi.

6136. Skeleton of *Scopus umbretta*.

S.C. 4 small fish & a shrimp  
Jan. 1. an old deserted nest ~~of~~ <sup>one</sup> new unfinshed  
6137. Waxbill ♂ t. c. Iris dark brown,  
<sup>one</sup> Estilda ~~actioid~~ <sup>angolensis</sup> Kehl.  
bill scarlet, feet dark brown, Crop contained small grass seeds.

6138. <sup>1</sup> Same. ♀ o. e. Iris brown, bill scarlet  
feet dark brown.

Feeding in small parties in tall grass in a marsh, frequented also by the common red-cheeked waxbill.



Borna.

Jan 2. 1915.

*Cyornis trivirgatus cyanocapilla*

6139. Sunbird. ♂. testes slightly enlarged  
Iris dark brown, bill & feet black.

6140. Chat. ♂ t-n-e. Iris dark brown.  
~ ~~*Saxicola solaris*~~ bill & feet black. S.C.: insects.

6141. Same ♀. o.n.e. S.C.: remains of  
~~*Saxicola solaris*~~ insects, & a large white beetle larva  
(“grub”)

Not uncommon; usually seen  
near cultivated ground <sup>or grassy</sup>,  
perching on bushes or tall grass stalks.

6142. Piney grass Warbler. ♀. o.n.e. Iris  
~ <sup>circum. & cisticola wrygalls</sup> rather light brown, bill dusky brown  
above, pale flesh-color below, feet  
pinkish-buff.

6143. <sup>Myiophantornis chloris</sup> Weavers ♂. testes somewhat enlarged.

6120. <sup>peruviana</sup> Iris scarlet, bill black, feet pinkish-  
brown, S.C.: fibers from palm-nuts,  
and remains of insects.

Jan. 3.

6144. Turacous. ♂. testes somewhat enlarged.  
~ Iris rather dark brown, papillae &  
edges of eyelids scarlet, bill dull  
dark red, blackish at very tip, feet  
black. Stomach contents: 9 wild  
dates. One of a pair in trees  
along a brook. The voice is  
very similar to that of *T. emini*.  
(reiterated resonant “haw”, as well as a  
low, hoarse, protracted “haw”.)

Boma

Jan. 3. 1915.

6145 Partridge (*Pternistes*) ♂ t. n. e. *crauchii crauchii*

Iris dark brown, bill, orbits, throat & feet scarlet. Crop contained many small bulbs (abt 1/4 in. in diameter)

6146. Same ♀ [im.] o. n. e. Crop contained many small bulbs, and numbers of termitites (not winged)

These little bulbs are the same as those from the partridge at Matali. I have found that they are the bulbs of a fine grass growing commonly between the tussocks of coarse grass, and lay usually an inch or so below the surface of the ground, but are turned up in quantities in freshly cultivated land.

6147 Barbet, <sup>*Sylvius levaillantii*</sup> ♂ t. n. e. Iris light yellowish-brown (bright buff), orbits flesh-color; bill whitish; feet pink, with gray claws. S.C.: fruit.

6148. Same. <sup>*Sylvius levaillantii*</sup> ♀ o. n. e. Colors of eyes, bill & feet same as in ♂.

6149. Young of preceding. ♂ juv. S.C. fibers from palm nut, & pieces of other fruit.

not uncommon, has a prolonged



Borna

Jan 3. 1915.

"clicking" note much like that of the large red + black related species of the Nelle - which I believe I have seen here also. (OK)

6150. <sup>Darlingius major</sup> Bush Shrike. ♂. testes slightly enlarged. Iris dark brown, bill black, feet dark bluish-gray. S.C.: remains of beetles, a bug, and a millipede.

Skulls in the densest bush they can find, the male occasionally uttering short whistles (much less musical than those of the species of the Nelle), and the female answering with a harsh "kük-ik."

6151. <sup>Darlingius sp.</sup> Thrush. ♂. testes somewhat enlarged. Iris dark brown, bare skin behind eye dusky greenish, bill yellow, feet pale brownish-green. Very common
6152. Clapper Lark, <sup>mirafræ</sup> <sup>feckeri</sup> ♂. testes slightly enlarged. S.C.: small insects.

Common here; habits same as in Nelle; frequenting barren spots on hills as well as cultivated fields in hollows. Wings may be beaten 5 or 6 times.



Borna

Jan 3. 1915.

6153. Finch (*Serinus*) ♂. t. sl. e.

6154. ~~*Spinus pereini* (Vieill.)~~

Blue-gray "Warbill" with red rump. ♂ imm. Iris brown; bill light blue, dusky at tip & along cutting edges; feet rather dark blue-gray.

Crop contained small grass-seeds.

~~*Lagonosticta lalandae*~~

6155. <sup>A.</sup> Weavers. ♂. t. m. e.

6156. ~~*Lagonosticta lalandae*~~ ♀. o. m. e. (mate of no 6155)

Crop & stomach contained small seeds.

Very common here.

Jan 5. 6157. Least Bittern. ♂. testis slightly enlarged.


~~*Ixobrychus fuscus*~~; Iris light orange-red, orbit greenish-yellow; bill yellow, shading to dusky brown above, feet yellow, tinged with green.

S.C.: a minnow.

Flew up from a grassy swamp. The purple gallinule collected at Avahuli also inhabits this swamp, also the "lily-trotter", which is frequently heard uttering a reiterated "ka-ka...." in a rather weak voice.

6158. Warbler (*Calamocichla*) <sup>superficies</sup> ♂. t. e.

Iris rather light brown; bill dusky brown above, light gray below; feet greenish-gray. Numerous in



Boma

Jan 5. 1915.



the large papyrus swamp on the east side of Boma; notes precisely like those at Stanleyville & Faradje. (So far as I can judge from noises emanating from this swamp, there is no *Bradypterus* present)

6159. Clapper Lark ♀. very slightly enlarged. Stomach contained remains of numbers of termites & a small elater.  
*Sturnella erythrops pyrrhoptera*
6160. Buff-cheeked Warbler ♂. testes slightly enlarged. Iris light brown, maxilla blackish, mandible light bluish-gray; feet pinkish-buff. Rather common. Its rather shrill notes consist mainly of sounds like "chwee", "chick" etc repeated several times, then changing slightly, this to be repeated again, & so on. Two birds often call together.
6161. *Eurillas curvirostris angolensis* - Schur. Waxbill. ♂. t.e.
- s. 6157. Jan. 7. *Somateria mollissima guttulata*.
6162. Shrike ♀ imm.
- s. 6150. 6163. Reed Warbler. ♂. t. m. e. Iris <sup>nearer</sup> dark brown, maxilla dusky, mandible light gray, interior of mouth orange;  
*Acrocephalus arundinaceus*

Boma

Jan 7. 1915.

feet greenish-gray.

Hiding in some long grass on low ground, giving a prolonged series of harsh scraping sounds, repeated again and again, with short intervals. (Rather like the voice of the small European reed-warbler, but very much louder).

These notes are often heard from a papyrus swamp here, and I remember, in Dec '11, hearing them from a papyrus swamp near Paradee.

*Xanthophilus xanthops* (Hartl)

6164. <sup>A</sup> Large yellow Weaver ♂ im. Iris

light yellow, bill dusky-brown above, shading to dull yellowish beneath, feet light pinkish-brown. S.C. insects.

6165. <sup>*lybius leucocephalus*</sup> Barbet ♀. o.m.e. S.C. fruit.

J. G. 147

6166. Kingfisher. ♂. testes small. Iris

S. 6119 <sup>*Halcyon alliacea*</sup> *orientalis*.


dark brown, bill rather dark red, becoming blackish at tip, feet red on softer parts, but scales brown.

S.C.: a millipede. <sup>Shot in open grove</sup> of rubber trees (*Mauritia*)

6167. Crake. ♀. o.m.e.; an egg in oviduct

*Crecopsis egregia*

(broken) was whitish, spotted with rufous, very heavily marked on one end. Put up in long grass in



Inches

Centimeters

Boma

Jan 8. 1915.

a hollow

6168 <sup>s. 6164.</sup> *Xanthophilus xanthops* (Hartl) Large yellow Weaver. ♂ testes slightly enlarged. Iris dull yellow; bill black; feet pinkish. Stomach contained soft fruit & remains of small insects.

6169 <sup>n</sup> Same. ♂ t. s. S.C. insects.

Usually seen in pairs about but occasionally on hills  
brushes in lower-lying ground.

Has a prolong wheezy note like  
*H. cucullatus*, but is comparatively  
shy, & not apt to be seen near houses.

6170 Sunbird. ♂. <sup>chalcinotis fuliginosus</sup> testes somewhat enlarged.

6171 <sup>s. 6129</sup> Paradise Flycatcher. ♂. t. e. <sup>Empidonax resplendens</sup> Iris dark brown; bill bright blue, blackish at tip, this color running back along edges of mandibles, interior of mouth greenish-yellow; rim of eyelids, & feet, blue. Shot near same spot as no. 6159. Has notes like Paradise Flycatcher of NE Congo, but the song, instead of being whistled is a slow repetition of some notes sounding more like "gee".

6172 <sup>n</sup> *estrilda penicillata* "Waxbill". ♀. o. e. Iris dark brown bill light blue, blackish at tip. Both specimens of this bird were shot among the trees planted at

Inches

1

2

3

4

5

6

7

8

Boma  
Jan

Ka

6173

w

6174

s. 6172

6175

6176

6177

6178

6179

s. 6105

6180

s. 6105

Boma  
Jan 8. 1915.

Kalamu.

6173. Thrush (*Erythropygia*) ♂. testes slightly enlarged.
6174. Babbler, <sup>Crateropus sp.</sup> ♀. o.m.e. S.C.: palm-nut fibers & insects.  
S. 6132.
6175. Nightjar. ♂. t.m.e. Iris dark brown
6176. " ♀. o.m.e. | rim of eyelids lighter brown, bill brown with black tip; feet brown. About 6 of these nightjars were put up among bushes on a hillside near the reservoir.
6177. Skeleton of Cypohierax <sup>angolensis</sup> ♂ t.m.e [adult]  
S.C.: palm-nut pulp and piece of a small fish.
6178. Widow (*Monachus*) <sup>or Colius passer asymmetrus</sup> with bases of remiges white. ♂. testes enlarged. Iris dark brown, bill pale blue, feet blackish-brown. S.C.: small seeds. Found in a grassy valley, accompanied by 5 or 6 others of its kind, females & young males, the latter with yellow shoulder patches, white in the wing, and blue beaks.
6179. Thrush. ♂. testes slightly enlarged, skull not fully ossified. S.C. insects & their larvae. Dried brown.  
<sup>Cichladusa rufigula</sup>  
S. 6105.
6180. Crateropus <sup>sp.</sup> ♂. testes slightly enlarged.



Boma

Jan. 11. 15.

Iris scarlet outside, shading to bright yellow  
on inner edge, bill black, feet dark gray.

S.C. fibers from palm-nuts, & insect remains.

6181. Same. ♂ t. n. e. skull not fully ossified.  
<sup>Croatocephalus</sup>

Iris as in adult. S.C. palm-nut fibers & insects.

6182. Chat ♂ t. n. e. S.C. small insects

6183. " " ♂ imm. (Have noticed no whinchats  
<sup>Saxicola relax</sup>  
in the Bas-Congo, and Hirundo rustica

is either rare or absent. But the  
black (European?) swift has been  
seen frequently, Merops persicus &  
M. albicollis are common, European  
reed-warbler & Muscicapa griseola  
also noticed.

6184. Pigeon. ♂ t. s. e.

6185. Coly. ♀ o. s. n. e. Iris rather dark  
<sup>Colius n. nigricollis</sup>  
brown, naked skin before eye black,  
that behind pale blue-gray; maxilla  
pale bluish on top, elsewhere black,  
mandible buffy-whitish, a little  
black at base; feet scarlet, claw-black.  
S.C. fruit & piece of a leaf.

One of a party of 6 — in spite of  
the fact that she was laying.  
A coly's nest found at Kivamouth  
(Dec 19-'14) was built like those  
of Nelle, in a bush 9 ft from



Boma

Jan 11. 15.

ground, and contained 3 eggs, colored like those of the Nelle.

6186 Roller. ♂ t.m.e. *Eurystomus ager*. Not common.

Jan 12.

6187. Green Pigeon ♀ o.e. as they leave a tree the flight of these pigeons is always very noisy, but becomes practically silent once they are well under way. Rather common here; voice same as in Nelle & Iture.

6188. Bronze Sunbird. ♂. testes somewhat enlarged. Common

6189. Red-wattled Flycatcher. ♂. testes somewhat enlarged. Common. May give from one to five clear whistled notes.


6190. Nightjar. ♀ im. Nightjars common at dusk about Boma, probably this is the common species. They make little or no noise, neither *macrodipteryx* nor *Scotornis* noticed.

6191. <sup>xanthophilus</sup> <sup>xaanthophylax</sup> <sup>(Hartl.)</sup> Yellow Weaver. ♂ t.m.e. Iris dull yellow, bill black, feet brownish-pink. S.C.: soft fruit & insect remains.

Jan 14.

6192. Widow. ♂ t.e. *olivaceus concolor* S.G. 178. grass seeds.

6193. Same. ♀ o.m.e. Iris "dark brown, bill



Inches  
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

Borna

Jan 14. 1915

brownish white, shading to brown on culmen, feet pale buff. S.C. grass seeds.

6194. Same. ♀. o.m.e.

Common, flocks of 20 to 30 have been seen. Noiseless; rather shy.

6195. <sup>A</sup> *Uraeginthus angolensis* (C.) Small blue breasted Weaver. ♂- teste slightly enlarged. Bill light pinkish-purple shading to dusky at tip; iris gray; feet pale buff. Crop and stomach contained seeds. Seen at matadi as well as here, but usually only near towns or native villages. Thus far I have never seen one with red cheek spots; yet the present example seems perfectly adult.

6196. *T. angolensis* very common at Ntiki in June 1909. Swamp Drail. ♀. o.m.e.; laying. Iris *Excalatoria adansonii* rather light red-brown. Crop and stomach contained small seeds, a few termites, and a tiny snail. One of a pair put up on the top of a rather barren hill.

6197. <sup>s. c. 153.</sup> Pinch (Scrinus). ♂. t.n.c. <sup>mozambicus teyde</sup>

6198. " " " " ♀. o.m.e. S.C.: small seeds.

Common; has semi-musical, cheeping voice, like the species of the Vele; but does not really sing or trill.

Borna

Jan 16. 1915.

6199. Halcyonichelicuti. ♀. o. in e.

S.C. acicada. Abundant here.

Two of these king fishers will often be seen in trees some distance apart often calling unison, sitting in a stiff upright position, and raising their wings. Other members of the genus - H. senegalensis for example - will go through about the same performance, but both birds perching on the same branch.

6200. Red-headed Weaver ♂. testes <sup>Ptilaea erythrocephala</sup> somewhat enlarged. Feeding in long grass in marsh.

6201. <sup>Estilda</sup> stridula angolensis <sup>Red</sup> Waxbill. ♂. testes somewhat enlarged. Crop filled with small seeds.

6202. Reed Warbler (Calamonocilla) <sup>referred?</sup> + e. (A fan-tail warbler <sup>(Schismocerca apicalis)</sup> seen Jan 14. in low, grassy, spot.)

6203. Buphaga <sup>large</sup> ♀. o. in e. Iris yellow; base of bill bright yellow, rest scarlet; feet blackish. Stomach filled with ticks and short hairs.

6204. Same <sup>♀</sup> in e. Iris brownish gray, Buphaga laetii outer portion of bill scarlet, base of maxilla blackish above, rest of bill yellow. Stomach filled with ticks and hairs.



Boma  
Jan

Zambi  
Jan 17. 1915

Sent up to me from agricultural station at Zambi, below Boma, where they are said to be common. They are not found about the cattle near Boma, though there are herds of over 100 head, and perhaps not even on the island of Mataba, where there are many cattle.

According to Mr Drousie they were not seen at Zambi before 1908, appearing first in company with a herd installed at some distance north of the river, and then, when these were brought back to the station the birds came too.

Boma, Jan 17.

6205. Durucus. ♂ <sup>persea</sup> testes slightly enlarged

6206. " <sup>persea</sup> S.C. <sup>perst</sup> <sup>♂</sup> o.v.e.

not uncommon, though there are few woods to shelter them.

6207. Cuculus solitarius. ♂ testes slightly enlarged.

6208. Phoenicurus. Redstart. ♀ imm. Iris dark brown, bill & feet dusky-brown.

6209. Colaptes caeruleus. Widow. ♂ t.c.; skull not quite fully ossified. Crop contained small

Boma

Jan 17. 1915

grass seeds. Iris dark brown, bill light blue, feet black.

*Strilda peruviana*

6210 A Weaver. ♀. o. v. e.


6211. *Periporphon*. ♂. t. m. e. } Were found  
6212 " ♀ imm. } together.

Jan 19 Clynias <sup>resembling</sup> *macroura* ♂. testes slightly enlarged  
6213. *Urolaevicia*. ♂. testes slightly enlarged  
Very common, Kwanzaouth; Leopold-  
ville, Matadi, Boma.

6214 *Serinus*, (<sup>capistratus</sup> black face) ♂. t. e. Crp  
x stomach contained seeds.

6215. Same. ♀. ovary slightly enlarged.  
S.C.: seeds.

To capture these finches, a cage containing a captive bird is hung in trees about the village, this cage having one or two separate compartments, where small seeds such as rice are scattered, and the door of which is snapped down by a spring when a bird alights on a perch within. *Spermestes* & the red *Lagonosticta* (?) are also caught, but tho' I have found the other species of *Serinus* much more common here, I have never seen it kept in a cage here.  
(Saw one in cage however at Sierra Leone)



Boma

Jan 20. 1915.

*Sitagra monacha*, Sharpe.

6216 A Weavers. ♂. testes somewhat enlarged.

Iris dull light yellowish, bill black, feet rather light bluish-gray. Stomach contained a small caterpillar & insects. Shot at edge of papyrus swamp.

6217. Widow. ♂. skull not quite completely s. 6209. *Culicivora caerulea* <sup>rim of eyelids blue gray</sup> asified. t. m. e. Iris dark brown, <sup>bill</sup> light blue-gray, feet dusky. S. C.: grass-seeds.

6218. Weaver. ♂ t. m. e. S. C. - small seeds. s. 5985? *Ploceus capensis*

Since about the first of the month, a male in full plumage has been seen occasionally in the high grass in a <sup>large</sup> swamp. This one was shot in an opening in a papyrus swamp. The species has not been noticed elsewhere about Boma.

Jan 21. 6219. *macronyx* <sup>croceus</sup> ♂. testes slightly enlarged.

Common (going in pairs) about a pasture bordering on a large plain (swampy at certain seasons). 2 1/2 hrs. walk NW of Boma. Not noticed anywhere else.

6220. Pipit. ♀. o. o. e. Iris dark brown, <sup>Anthus pallidiventris</sup> bill dusky brown, becoming pale yellowish brown-color on base of mandible feet buff. 2 pairs seen today.

Borna

Jan 21. 1915.

in pasture mentioned under no 6219.

They were silent, and in habits like the species of the Vele.

6221. Lark ♂ t.m.e. S.C.: remains of <sup>mirafra</sup> ~~leachii~~ grasshoppers and many small termites.

6222. Coly ♂ testes somewhat enlarged. S.C.: <sup>C. n. Sigricollis</sup> leaves. Iris rather dark brown.

Jan 24.

6223. Bee-Eater. ♀ t.m.e. Iris dark brown, bill black, feet blackish. <sup>Melittophagus bullockoides</sup> 3 sitting on dead branches of a baobab fruit to be seen.


6224. Barbet. ♂ t.e. Crop stomach contained fruit. Not uncommon, call as usual, 4 or 5 notes, then a slight pause.

6225. Pyronella <sup>dominica</sup> ♂. testes slightly enlarged. In prenuptial molt. Very common, but very few have begun to molt yet. (A male in full plumage was seen at Yaronge nr Stanleyville on Dec 10; but none further down near Bumba.)

6226. <sup>Hyphantornis collaris</sup> Weaver. ♀. ovary slightly enlarged. S. G. 6143. Iris scarlet. S.C.: insects.

6227. Red start ♂ t.m.e. S.C.: small fruits & insects. Shot at same spot

" 601.



Borna

Jan 24. 1915.

as no 6208, near a wooded ravine on  
a hillside *androphilus* sp.

6228. Bullbul. ♀. t. n. e. S.C.: insects.

6229. Sunbird. ♂. t. n. e. *Cinnyris moquensis* Gmelin's  
Jan 25 *Trochiliphaga rufoviridis* Common.

6230. Paradise Flycatcher. ♂. testes slightly  
enlarged.

6231. Coly. ♀. t. n. e. S.C.: fruit (Picus)  
C. w. nigricollis  
Common, often in small flocks.  
Utters short hoarse notes, but is  
generally silent.

6232. Merops albicollis. ♂. t. n. e.

Common here at present. Many  
of the migrants from Europe  
are very little in evidence here.

*Hirundo rustica*, *Riparia*, *Saxicola*  
*Pratincola*, *Anthus trivialis*; these  
are some of the common ones that  
appear to be lacking. *Cypselus*  
*& Acrocephalus*, *Sylvia*, *Muscicapa*  
*griseola* have been noticed.

6233. <sup>Dendropicos</sup> *hartlaubii*  
Woodpecker. ♂. t. n. e. Iris dark  
red, bill blackish with base of  
mandible bluish gray, feet dull green  
S.C.: many larvae & a few pupae of insects  
This seems to be the only woodpecker  
at all common about Borna.

Borna

Jan 25, 1915.

*Antichromis minutus*

6235. Shrike ♂ t.m.e. Iris rose,  
bill black; feet lead-gray. Rather  
common in lowlands.


6236. Pratincola salax. ♀ imm.

6237. Pytelia <sup>melba</sup> ♂. testes slightly enlarged  
iris brownish-orange, bill rose  
(but without gloss), feet grayish-  
brown. Crop & stomach contained  
termites (workers) & small grass-seeds.  
not a common bird at Borna,  
probably fond of hills.

Jan 27

6238. Porphyrio <sup>alleni</sup> ♂. t.m.e. Stomach  
contained pieces of a large spider,  
small seeds, insect-remains, &  
bits of stone. Iris scarlet, bill  
scarlet but darker, frontal plate  
dusky greenish; feet dull red,  
but shading to bright scarlet  
on tibiae.

Shot in a papyrus swamp  
where both adult & immature  
specimens had been observed  
previously. Limnocorax also  
inhabits this swamp, and  
the "lily-trotter" (Jacana) may  
also be seen in open places in



Boma  
Jan 27.

it,  
abor  
Cong  
not  
but  
var  
low  
abu  
ma

*Pycnonotus* variorum TC.  
Poponochyarchus (large  
size & tail) 3.  
Bullbul like no 6073. only 3.  
Bias ~~in~~ <sup>minimus</sup> (very mitratus)  
Hirundo smithii  
Passalodrepanis (tail tail  
comes apertive under wing-coverts  
white)

Hirundo smithii  
Passalodrepanis TC.  
Large Long-tail Weaver  
T. C.  
Small-tail Shrike 1.  
♂: A medium sized. Black  
face & lower part with  
breast probably tinged with  
brown, and black of head  
not reaching nape, in  
parties.

Long-tailed Yellow Weaver.  
*Hypergerus senegalensis*  
Passer gossypiinus (not common)

6239. *Xanthophorus xanthophrys* (Hartlaub) ♀. ovary somewhat enlarged. Iris rather dull yellow, bill black, feet light brownish pink. S.C.: green seeds.
6240. Widow T. t. e. *Colius caeruleoconcolor* S.C.: small grass seeds.

Libreville, Gaboon. Feb. 12.

6241. *Pelecanus rufescens*. ♀. o. e. Iris gray, tinged with purplish, lores mottled blackish, eyelids & naked orbits orange-yellow, bordered with blackish posteriorly; bill pale lemon-yellow, becoming pale grey on outer part of mandible, tips (or "nails") of both maxilla & mandible light cadmium-yellow; pouch light rufous and bright yellow, the two colors alternating in

Boma


Jan 27. 1915.

it, as also on the Kalamu reservoir.  
About water-birds in the Bas-Congo I can say almost nothing,  
not having been out on the river,  
but there is said to be a fair  
variety of aquatic species, at  
low water at least, especially  
about the islands below Boma,  
Mataba I., etc.

6239. *Xanthophorus xanthops* (Hartl)  
Large yellow Weaver ♀. ovary somewhat enlarged. Iris rather dull yellow,  
bill black, feet light brownish pink.  
S.C.: green seeds.
6240. Widow ♂. t.e. <sup>*Colius spissus concolor*</sup> S.C.: small grass seeds.

Libreville, Gaboon. Feb. 12.

6241. *Pelecanus rufescens*. ♀. o.e. Iris gray, tinged with purplish, lores mottled blackish, eyelids & naked orbits orange-yellow, bordered with blackish posteriorly; bill pale lemon-yellow, becoming pale gray on outer part of mandible, tips (or "nails") of both maxilla & mandible light cadmium-yellow; pouch light rufous and bright yellow, the two colors alternating in



ish  
ed.  
d  
sh.  
L.  
-

Birds seen at Boma  
not secured.

Commoner [ islands in  
Smaller Birds ] on Rivers  
below Boma 4 times.  
Belcher  
Macrorhamphous (over)  
(Gularis) (few)  
Pipileashin (few)  
Pipileashin (not uncommon)  
Hypothymis 1.  
Numida (5 or 6) not common  
Actitis common.  
Jacana Rallinaeum  
Conical (white throat & breast)  
Ceryleacea chloris - C.  
c. amayensis (also at Matadi)  
Agapornis pullaria. 3.  
Lophoceros TC. nigris in  
infuscatus) 4. seen to mate.  
Rufophaea 4.

*Buteo maculius* (also at Matadi)  
*Hapaloderma marginata* 1 sp.  
famed with forman i firm  
Kakamega species are by no means

of this species, was seen on an island below Boma, Jan 30 th. none noticed on the Congo in coming down from Stanleyville to Kinshasa.

The present example was bought in the market at Libreville and was said to have been captured with the fish-net known as "l'épervier", which is thrown out over the water.

The scarcity of water-birds (as seen from the steamer at least) in the different ports at which we have called (Sandana, Loango, Cape Lopez, Gaboon R, Fernando Po (St Isabel)

not reaching stripe in  
uppereyes.


Long-tailed yellow weaver.  
Passer luteus (Vigors)  
Passerina (not common)

rather narrow cross-lines, of which  
the yellow ones are slightly raised,  
feet (including webs) rather dull  
red, paler behind metatarsi and  
beneath toes, claws light grayish-  
green. Stomach contained 3 fish,  
swallowed head first, largest  
one 10 inches long.

A flock of pelicans, apparently  
of this species, was seen on an  
island below Boma, Jan 30th.  
None noticed on the Congo in  
coming down from Stanleyville  
to Kinshasa.

The present example was  
bought in the market at Libreville  
and was said to have been  
captured with the fish-net  
known as "l'épervier", which  
is thrown out over the water.

The scarcity of water-birds  
(as seen from the steamer at  
least) in the different ports  
at which we have called  
(Sandana, Loango, Cape Lopez,  
Gaboon R., Fernando Po (St. Isabel))



is striking. There are usually a few large Terns, with light reddish beaks and black nape, which often alight on floating logs or buoys, but as a rule this is about all, tho an <sup>(Pandion haliaetus)</sup> osprey was seen at Landana, and a couple of *Gypohierax* at Fernando Po.

(St. Isidore)

At this latter town, under the roof of a balcony, there were a number of nests of *Cypselus affinis*, white rumped swifts, like that collected at Matadi, and also seen at Boma & Libreville. They were built of feathers and dry grass (or other similar material), shaped like nests of certain swallows (*H. puella*), with a rather small lateral entrance. The birds were numerous, probably 50 or 60 at least, and not shy. One dead and dried-up example was secured that hung by its neck from a nest, as tho it had been



Inches  
1  
2  
3  
4  
5  
6  
7  
8

caught by a hair or fiber  
and strangled.

In Sierra Leone harbor for the first time we saw numbers of gulls, mostly a middle sized species (*L. cirrocephalus*?), the adults of which had gray heads, under sides of wings blackish, mantle blue gray, but wing coverts darker. One adult black-backed gull (*L. fuscus*?) also noticed. There were likewise many terns, partly *Hydrochelidon* in winter dress, the others resembling Common Tern, and only black on the back of the head. A few kites (*Milvus*) also hovered over the water, and numbers of brown vultures (*Neophron*) sailed about over the town and sat on trees and house tops.

The day after leaving Sierra Leone (no land in sight) there were shearwaters, following the ship, sometimes 2, sometimes as many as 6 or 8. As we were making only 9 knots or so they would often sail down,





Mar 7  
Kottiwakes 3.



On Feb 28, Mar 2 + 3, these snakes

alight on the water like gulls, and drop far behind; easily catching up to us again.

They varied considerably as to color. A few were sooty brown all over; others were sooty brown, but whitish about the vent, and broadly margined with whitish on abdomen, rump, and under tail-coverts. Still others were brown save for the <sup>white</sup> lower breast and abdomen, the undertail-coverts being dark brown, not margined with whitish; and in one the white of the breast went a little further forward and around the back of the neck. But the commonest pattern was this: brown on crown and sides of head, back, wings, tail & undertail-coverts, whitish throat, hind neck, breast & abdomen, with a dark, rather narrow, breast band. Some of the last mentioned, and one of the pure-brown ones shows the prolonged & twisted rectrices, which were not pointed, so I believe they were all *S. pomatorhinus*.

On Feb 28, Mar 2 + 3, these shrews

(*Stercorarius pomatorhinus*)

S. Lays Feb 27-28 <sup>more</sup> 12-3.

1. All sooty-brown
2. All sooty brown, but feathers of abdomen, rump & under t. coverts broadly margined with white.

3.



4.



Nos 1 + 3 with  
elongated rectrices  
possibly *S. parasiticus*


5.



Mar 4. Kittiwakes 6 or 8 following ship,  
Mar. 6. Kittiwakes 2.

S shearwaters

l. 5



~~base of bill yellow  
underparts, incl.  
throat & upper d  
wing-coverts white  
tail (above) whitish  
on light gray with  
black tip, square &  
slightly rounded  
rather short~~

Oceanodroma 1. gray above, white below  
forked tail

Megalestris (?) 2. brown, edges with  
lighter color, bases of primaries white  
at least had middle tail feathers  
slightly elongates.  
waters were noticed, flying low




still followed us, tho' of course none were seen in the harbor at Dakar. There there were many gulls, of two species, the lesser Black-backed (*L. fuscus*) and a smaller, black-headed gull — tho' only a few were black on the head at this season — with dark red bill, which I took for *L. melanoleuca*, tho' the tip of the wing was edged with black.

But the day before we reached Teneriffe Kittiwakes replaced the Skuas, mostly young birds with black-banded tail & blackish on nape & wing coverts, even the few adults having dusky check spots.

Numbers of lesser Black-backed Gulls were again seen at Santa Cruz (Teneriffe), and other gulls of the same size, but the mantle lighter and grayer, tho' much darker than Herring Gulls, especially on the wing coverts.

About 90 miles northward, the following morning, 5 or 6 shearwaters were noticed, flying low



over the water as to be hidden frequently behind a wave. They were gray-brown above, white below (including throat & under wing-coverts) the bill mostly yellow, but dark at tip, and the rather short square tail seemed (above) whitish or light grey with black band at or near tip.

This same morning we also saw a fork-tailed petrel, gray above & white below, skimming along just above the water, which it would strike at intervals with its foot. (*Oceanodroma*?)

Towards noon (Mar 6) two Shearwaters flew about the ship, larger than those previously seen, brown with buff-edged feathers and white bases to the primaries. One at least had the middle tail-feathers slightly prolonged (*megalastis*?)

In many of the shops at Dakar where post-cards, periodicals & souvenirs are on sale one may buy bird-skins, roughly prepared & usually minus the feet. The species

I noticed thus are as follows:

"Foliotocole" (*Chrysococcyx m. aragdineus*)

Parakeet (*Palaeornis*)

Rollers (*Coracias*)

Rosy Bee-Eater (*M. nubicus*)

Oriole (a yellow-crowned sp. like that of Ule)

Glossy Starlings:

A small species of *Lamprocolius* like those of

A similar bird but with long graduated tail.

A small purple species with white belly

Sunbirds


A dark brown species with metallic  
finishes & red breast patch.

A nectarinia, metallic green, even  
on belly, with 2 elongated rectrices &  
a red breast patch.

In one of these shops there were hanging  
2 trogons (*Pharomacrus*) with long tail.  
coverts, evidently from S. America.

Every day, from Mar 6 to 9, there  
were kittiwakes sailing at our stern  
and hovering in our wake, perhaps even  
8 or 10 at a time, and on Mar 9 a large  
shua (*Megalestris*). On Mar 10 we had  
passed C. Finisterre; and were followed  
all day by a half-dozen lesser black-  
backed gulls, but no kittiwakes.





- [cont'd.]
- 157511 — *Rhynchosia fluminensis* St. Hil. 1915 Pauli, Rio Longo  
 157512 — " " " ♀ " 22, " " "  
 157513 — " " " ♂ " 23, " " "  
 157514 — " " " ♂ " 23, " " "  
 157515 — " " " ♀ " 23, " " "  
 157516 — " " " ♀ " 23, " " "  
 157517 — " " " ♂ " 23, " " "  
 157518 — *Galactochyrea amara* ♀ " 23, " " "  
 157519 — " " " Young ♂ 23, " " "  
 157520 — *Odeoniasse nemoralis* ♀ " 23, " " " Pauli "  
 157419 — " " " ♀ " 23, " " "  
 157420 — " " " Capensis ♂ " 23, " " "  
 157421 — *Burrota melanogaster* ♂ " 23, " " "  
 157572 — *Cyclista gazella* ♂ Sept. 4, " Banana "  
 157620 — *Sila capensis* ♂ " 23, " " "  
 157621 — " " " ♂ " 23, " " "  
 157622 — *Pelcones ruficervus* ♂ Jan? " Zambia (Van Seep)  
 157763 — *Polykorda typica* Young Jan 10, " Stanleyville  
 158404 — *Acis grandiflora* Dec. 1914 Bonn Dr. Ags.  
 159465 — *Comptonia* ♂ Jan 16 1915 " "  
 160625 — *Mammillaria* Binn. June 25, " Zambia  
 161187 — *Tacca callae-aurea*? June 24, " "


*0291*  
*BB*  
*BB*


163005 — *Colius fasciatus macrourus* ♀ June 24, 1915 *Zambézi* <sup>Ba</sup> *Cong*  
163002 — *Buphaga longi* ? July 4 " "  
163003 — " " ? " 4, " " "  
163004 — " " ♂ " 25 " " "



Birds collected by Lang after my departure.

6242. Gymnomyza typica, ♀ imm.  
Stanleyville, Mar. 10, 1915, Stomach  
filled with pulp of palm nuts, also  
two small kernels.  
Zambi, Lower Congo
6243. Rynchops flavirostris ♂ ad June 28, 1915-
- 6244 " " ♀ ad June 22, 1915-
- 6245-6 " " ♂ juv. June 25, 1915-
- 6247-49 " " 3 ♀ juv. " " "
6250. Galachrysa cinerea ♀ ad. June 22, "
- 6251 " " ♀ juv. " " "
- 6252 Oedicnemus vermiculatus ♀. June 28, 1915-
- 6253 " " ♀. June 30 "
- 6254 Oedicnemus capensis ♂ June 30, 1915-
- 6255 Lissotis melanogaster ♂ June 22, 1915-
- 6256 Pelecanus rufescens [♂] January?  
Skinned by Dr. Van Soegehem.
6257. Ficulus collaris smithii June 24, 1915-
- 6258 Myrmecocichla nigra ♂ June 25, 1915-
- 6259 Buphagus africanus longi [?] July 4, "
- 6260 " " " [?] July 4, "
- 6261 " " " ♂ July 25, "
- 6262 Colius passer macroura ♂ June 20, "
- Banana, Congo mouth
- 6263 Egretta garzetta ♂ Sept. 6, 1915. "ov." not  
enlarged. Lores grayish green, bill black, lower  
part of mandible near base pinkish; feet





Birds collected by H. Long (continued).


Banana, Congo mouth.

black, toes and a little of tarsus above  
them light green.

6264 Morus capensis ♂ Sept. 8. 1915-

6265 " " ♂ Sept 8, 1915.


— o o —



10

Blank

pages



Egyptian Goose. ♂ Banane Sept 6.  
1915. (A juv n. no. 157572) " or " not  
enlarged. ~~green~~ <sup>greenish</sup> green; bill black;  
lower part of mandible near base pinkish;  
feet black, toes & a little of tarsus  
above them light green.



Fish.

nouvelle Anvers, Dec 14. 14 — Purple — Dusky green above and on sides, shading to grayish-white on belly (in front of anus); iris red-brown.

Coquihattaville, Dec. 16. (Fish shaped like *Polypterus*) Dark brownish green above, sides much the same color, but with a line of large blackish blotches, and mottled somewhat between these blotches with dull orange. Throat & rest of under side whitish mottled with dusky. Dorsal, caudal, & anal fins dusky-greenish, spotted basally with blackish & dull orange, pectoral & ventral fins yellowish, spotted with blackish. Iris greenish above and below, dark brown before & behind, & with a thin inner rim of reddish-brown.

Near Coquihattaville, Dec 16. *Polypterus* — General color above brownish-green, with large dusky patches. Below, yellowish-white. Dorsal fin dusky, greenish on ray base, grayish, ventral & anal fins yellowish, caudal yellowish, spotted with black, dorsal translucent gray, marbled with blackish. Iris brown.

Coquihattaville Dec 16. Small compressed fish — white on belly, shading gradually to dusky above, but with silver fall over, and also bluish reflections above. Many blackish horizontal stripes. Fins gray, a little yellow on pectorals & ventrals. Iris silver.

Irebu, Dec 17 — Perch — General color above changing from light to dark green or brownish green, usually lighter below, whitish on belly. Spots of reddish on base of each scale on sides of body, & dorsal, caudal, & anal fins dull greenish, thickly spotted with reddish, these spots varying from dusky purplish to fairly bright red together with color of rest of body. Pectoral and ventral fins dull greenish, also tinged at times with red, a large black

spot on edge of operculum, cheeks  
with coppery reflections. Iris green above,  
silver below; upper rim red. Lower lip &  
throat whitish, often tinged with blue.  
Fins. Bright red markings often appear on  
throat and chest, as in a related species  
painted at Awakeni.

Boma, Jan 1. '15. "White-Bait" — Translucent  
whitish all over; silver reflections on  
sides of head and gill covers, also on belly;  
a longitudinal band on side of body. Fins  
whitish save caudal, which is pale yellow  
with black tip. Top of snout & head bordered  
with blackish. Iris blackish above, silver  
below, and with a narrow silver inner  
rim.

Kinshassa Dec 22. Large ground-lizard.  
Iris buffy-yellow. Crown greenish-brown  
shading to orange-red on cheeks and sides  
of lower jaw. Back and upper surface of  
tail greenish brown, with a black  
spot on the middle of each scale. Along  
each side of the back, from the eye  
nearly to the tip of the tail runs a greenish  
yellow stripe, bordered on both sides  
with black. Sides of body blackish above  
shading to greenish-gray lower down,  
and flecked with yellow. A yellow  
line runs from the ear nearly to the  
base of the fore-leg. Sides of tail  
greenish-gray, faintly flecked with  
yellow. Upper surface of front & hind  
limbs rather dark greenish-gray.  
Throat blotched with reddish & yellow,  
whole rest of under surface dull  
greenish-white.



#7000-

7001 -

7002 -

7003 -

7004 -

7005 -




## Large Mammals.

- #7000- Red colobus ♀ ad. Lubolela. Dec 17. 1914.  
TL. 127 LT. 69 LH. 17. LE 3.5
- 7001- Red colobus. ♀ imm. Lubolela Dec. 17. 1914.  
TL. 104; LT. 58.5 LH. 15. LE 3.4  
Iris in both specimens light brown, ischial callosities dusky-greenish. Stomach contained light green vegetable matter, probably chewed-up leaves.
- 7002- Hyrax ♀ ad. Matadi. Dec 27. 1914.  
TL. 522 LT. abt 12. LH. 6.8 LE. 3.5  
HS. 16 HP. 16 LB. 33. 2 embryos in uterus. Stomach contained leaves, well chewed up, and with a slightly aromatic smell. Iris blackish, skin of nose, eyelids, ears & soles of feet black. The tail does not protrude, and cannot be bent upward, hence Hyrax ♀ imm. Matadi Dec 27. 14
- 7003- TL. 43, LT. abt 1. LH. 6.0 LE. 3.0  
Stomach contained leaves, like preceding (neither had eaten any grass as far as I could see)
- 7004- Bushbuck. ♀ ad. Boma, Jan 16. (1 foetus)  
TL. 130.5; LT. 24; LH. 33; LE, 13; H.S. 65  
HP. 70; LB. 65.  
Rather common, hiding in long grass and bushes, especially in hollows.
- 7005- Reedbuck. ♂ ad. Boma. Jan 17.  
TL. 155; LT. 29; LH. 48.5; LE. 16.5  
Iris brown, rather light posteriorly, darker and with a slight purplish tinge near pupil. with is elongated horizontally. Not uncommon, especially in broad valleys with long grass.

6429

Inches





Continuation

