

AMERICAN MUSEUM *Novitates*

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, N.Y. 10024
Number 3174, 59 pp., 16 figures June 26, 1996

Catalog of Recent Type Specimens in the Department of Invertebrates, American Museum of Natural History. III. "Parasitica" (Phyla Platyhelminthes, Rhombozoa, and Pentastomida) and Gastrotricha (Supplement)

CHRISTOPHER B. BOYKO¹

ABSTRACT

A complete, annotated listing of all type specimens of "parasitica" (including Turbellaria) in the collection of the Department of Invertebrates at the American Museum of Natural History (as of December 1995) is given, as well as a supplement to the Gastrotricha. A total of 608 type lots from the four phyla is listed. Seventeen lectotypes are des-

ignated herein and six changes in generic placement are made along with notes on unjustified emendations. One new replacement name is given for a subgenus of Turbellaria. Illustrations are provided for 13 lectotypes and three neotypes. An index of genera and species is included for each phylum.

INTRODUCTION

The Department of Invertebrates at the American Museum of Natural History currently houses a collection of over 350,000 cataloged lots divided among three major categories: fossil (45,000 lots), Recent mol-

lusks (275,000 lots), and other Recent phyla (30,000 lots). All of the types are housed separately from the main body of the collection. This allows for a greater level of security than can be maintained in a large, general collec-

¹ Department of Zoology, University of Rhode Island, Kingston, Rhode Island 02881-0816.

tion area as per ICZN Recommendation 72 (ICZN, 1985: 147).

This catalog is the third in a series of comprehensive listings of all type material in the department's collection. It is also an updating and expansion of the work of Feinberg (1970, in part) which contains many inaccuracies and omissions. Included here are all AMNH holotype, allotype, lectotype, neotype, paratype, paralectotype, and syntype specimens from the three "parasitica" phyla (including the primarily free-living Turbellaria): Platyhelminthes (242 species), Rhombozoa (7 species), and Pentastomida (21 species). In addition, supplementary Gastrotricha type material (5 species) has come to light since this group was treated in Boyko (1994) and is included as well. A total of 155 holotype, 1 allotype, 23 lectotype, 9 neotype, 329 paratype, 43 paralectotype, and 49 syntype lots is listed in this publication.

This catalog includes an accounting of the type specimens that have been transferred to this institution from other sources. Very often no published record of these transfers is ever made, especially for specimens coming from private collections, such as those listed below. In several cases, types were published as having been given to the American Museum, but subsequent investigation determined that the specimens were never deposited. Other members of type series with representatives in the AMNH have also been subject to transfer between institutions over time and the most recent known repositories have been listed, as well as the most recent catalog numbers used to identify type lots in other institutions. A specific example of this kind of transfer involves the types of the turbellarian species described by Hyman (1953a). Many of these specimens were originally deposited in the Allan Hancock Foundation, but have been tracked by this author to their (previously unpublished) current repository at the Santa Barbara Museum of Natural History.

The majority of the specimens included in this catalog come from three sources: the Horace W. Stunkard Collection of platyhelminths, the Libbie H. Hyman Collection of turbellarians, and the John Teague Self Collection of pentastomids. Of these, the Hyman

and Self Collections are perhaps the best curated, the former by H. S. Feinberg immediately following Hyman's death (see Feinberg, 1970, for details) and the latter by virtue of its donation and subsequent use by J. T. Self and his associates, which continues to the present. The Stunkard Collection was, in large part, donated in 1983–84 by Stunkard. Many labeling errors have been corrected by the present author working with information from other institutions, primarily the U.S. National Parasite Collection, in order to identify, as accurately as possible, the majority of Stunkard's type material. Stunkard's occasional a posteriori labeling of specimens and designation of types and his confusing (and inconsistent) use of the terms "type" and "co-type" on his slides have undoubtedly caused some of his material to be unrecognizable as types.

The AMNH collection has been organized by phylum and "superphylum" (Protozoa). Each division is given a separate series of catalog numbers. As a result of this system of numbering, a pentastomid may have the same catalog number as a trematode, but the specimens cannot be confused because of the taxonomic distinction between phyla.

Each type lot was compared to the original description and figures in order to verify type status, confirm locality and collection data, and remove any specimens from the type collection that are not valid types. The publication of this catalog fulfills ICZN Recommendation 72G(4), which states that all museums holding type specimens have the responsibility of making a published catalog of these types available to the scientific community (ICZN, 1985: 147).

This catalog is arranged by phylum and class, based primarily on Brusca and Brusca, 1990. Within each class, the types are listed in alphabetical order by genus and species. An index is provided, which gives a genus- and species-level list within each phylum. This catalog follows Boyko and Sage (1996) by including detailed information on current synonymy and generic placement of all species listed.

In the case of 70 species treated herein, no holotype was designated in the original publication. For 23 of these species, one speci-

men in the AMNH collection is clearly either the figured specimen in the original publication or was labeled by the author of the species as being the holotype. Such specimens not subsequently published as holotype or lectotype are herein designated lectotype (any postpublication reference to a holotype where none was originally designated being the equivalent of a lectotype designation). Thirteen of these specimens are figured, along with three previously designated neotypes.

In the section on Turbellaria, several species of polyclads are transferred to new genera in order to conform to both prior published inferred generic placement and ICZN rules. This action was taken because of the many inconsistencies and errors in the papers by Faubel (1983, 1984) which inspired the critical comments of Prudhoe (1989). Prudhoe refuted portions of Faubel's monograph, especially the validity of classifying polyclads solely on the basis of the male copulatory organ. Where discrepancies occur in current generic placement or synonymy, both authors' views are presented since deciding such matters is beyond the scope of the present paper. At the very least, some adjustment in Faubel's taxonomy is required (see remarks under *Notoplana segnis* and *Planocera burchami*, for examples). In addition, one subgenus (*Kytorhynchoides*) of Turbellaria is deemed invalid for lack of designation of a type species and a replacement name is introduced, with designation of the type species. Notes on unjustified emendations and spelling errors are included for several species.

Each type listing contains four to six categories of data. Within each genus, specific names are followed by authorship data (author, year description was published, and page and figure numbers). The next category indicates the specific kind of type(s) in the collection, as well as AMNH catalog number(s), and the number of specimens in each lot. The locality category contains all pertinent collection data, including county names for U.S. locations where that can be determined. The collector category indicates the collector(s) of the specimen(s) and the date(s) they were collected, if those data are known. The remarks category contains information as to prove-

nance of specimens, location of other members of the type series, current synonymy and/or generic placement, etc. The host category is included for species that are parasites or commensals of other species.

Institutional Abbreviations

AHF	Allan Hancock Foundation, University of Southern California, Los Angeles
AM	Australian Museum, Sydney
AMNH	American Museum of Natural History
ANSP	Academy of Natural Sciences of Philadelphia, Pennsylvania
BPBM	Bernice P. Bishop Museum, Honolulu, Hawaii
BM(NH)	British Museum (Natural History) (now The Natural History Museum), London
BOC	Bingham Oceanographic Collection, now part of YPM
CASIZ	California Academy of Sciences, Invertebrate Zoology, Berkeley
FST	Faculté des Sciences de Toulouse, France
HCUI	Helminthological Collection, University of Illinois, Urbana
MACN	Museo Argentino de Ciencias Naturales "Bernardino Rivadavia," Buenos Aires, Argentina
MBL	Marine Biological Laboratory, Woods Hole, Massachusetts
MIRP	Molteno Institute for Research in Parasitology, Cambridge, England
NHMSU	Natural History Museum, Stanford University, California
NRS	Naturhistoriska Riksmuseet (Stockholm), Sektionen för Evertebratzoologi, Sweden
SBMNH	Santa Barbara Museum of Natural History, California
TIU	Tokyo Imperial University, Zoological Institute, Faculty of Science, Japan
UNS	Universidad Nacional del Sur, Laboratorio de Ecología Acuática, Bahía Blanca, Argentina
URI	University of Rhode Island, Department of Zoology, Kingston
USNM	United States National Museum (now National Museum of Natural History-NMNH), Washington, D.C.
USNPC	United States National Parasite Collection, Beltsville, Maryland
USP	University of São Paulo, Department of Zoology, Brazil
UW	University of Windsor, Department of Biology, Ontario, Canada
YPM	Yale Peabody Museum, New Haven, Connecticut

ACKNOWLEDGMENTS

I would like to acknowledge the contributions of the late Walter E. Sage III, Senior Scientific Assistant in the Department of Invertebrates at the American Museum, for his critical comments and suggestions, for help in locating pertinent references, and especially for his friendship over the past several years. He will be missed. Additional thanks to the following for data related to specific sections. Turbellaria: Liz Kools (CASIZ), Eric Lazo-Wasem (YPM), and Barbara Littman (USNM). Special Thanks to Paul Scott (SBMNH), who kindly spent time unearthing and cataloging the Hyman types there. Monogenea, Trematoda, and Cestoda: Patricia Pilitt (USNPC). Mesozoa: Linda Cole (USNM) and F. G. Hochberg (SBMNH), who supplied data from McConnaughey's original notebooks. Thanks also to Alan Harvey (AMNH), Robert Bullock (URI), and Robin Overstreet (Gulf Coast Research Laboratory, Mississippi) for reviewing the manuscript. Photographic assistance was provided by Robert Bullock (equipment) and Robert Moyle (URI). Alan Harvey and Deborah Levine assisted with the preparation of illustrations.

PHYLUM PLATYHELMINTHES

CLASS TURBELLARIA

Acanthiella chaetonotoides Rieger and Sterrer, 1975: 234–246, 248, figs. 17–24.

Holotype: AMNH 814.

Locality: Station 14460, 20 m, deep sand, 34°25.5'N, 76°20'W, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: R/V "Eastward" cruise E-10-70, Oct. 1970.

Remarks: ex Rieger and Sterrer. The repository of the paratypes was not cited. This species is the type of the genus *Acanthiella* Rieger and Sterrer, 1975, by monotypy (Rieger and Sterrer, 1975: 248).

Acanthozoon albopapillosum Hyman, 1959a: 583–584, fig. 14.

Holotype: AMNH 479.

Locality: Palau Islands, Caroline Islands, Pacific Ocean.

Collector: Unknown, July 22, 1954.

Remarks: ex Hyman. AMNH 479 is the holotype by monotypy. This species is placed in *Thysanozoon* Grube, 1840 (Prudhoe, 1985: 132).

Acerotisa californica Hyman, 1953a: 379–381, figs. 150, 152–154.

Paratypes: AMNH 657.1–657.3 (3 specimens).

Locality: On wooden blocks suspended in water at various levels, Los Angeles Harbor, Los Angeles Co., California, United States, Pacific Ocean.

Collector: D. Reish.

Remarks: ex Hyman. The holotype was in AHF (Hyman, 1953a: 381), now SBMNH 142925. The repository of additional paratypes was not cited and they could not be located in SBMNH. This species is placed in *Stylostomum* Lang, 1884, by Prudhoe (1985: 140), but is retained in *Acerotisa* Strand, 1928, by Faubel (1984: 221).

Acerotisa multiselis Hyman, 1955c: 139–140, figs. 87b, c, 88a.

Paratypes: AMNH 658.1–658.5 (2 specimens), 659 (1 specimen).

Locality: Biscayne Bay, Dade Co., Florida, United States (AMNH 658); San Juan Harbor, Puerto Rico (AMNH 659), Atlantic Ocean.

Collector: F. M. Bayer, Aug. 1, 1951 (AMNH 658), Smithsonian-Hartford Expedition, March 29, 1937 (AMNH 659).

Remarks: ex Hyman. The holotype is USNM 24627 and additional paratypes are USNM 24628, 24629 (Hyman, 1955c: 140). This species is placed in *Eurylepta* Ehrenberg, 1831 (Faubel, 1984: 220).

Afronta aurantiaca Hyman, 1944: 1–2, figs. 1–3.

Holotype: AMNH 339.

Locality: Dredged in mud, 30 ft, near biological laboratory, Mount Desert Island, Hancock Co., Maine, United States, Atlantic Ocean.

Collector: Unknown, Aug. 16, 1937.

Remarks: ex Hyman. AMNH 339 is the holotype by monotypy. This species is the type of the genus *Afronta* Hyman, 1944, by monotypy (Hyman, 1944: 1).

Alleena mexicana Hyman, 1953a: 275–276, figs. 1–3.

Holotype: AMNH 355.

Locality: Gulf of California, Mexico, Pacific Ocean.

Collector: Unknown.

Remarks: ex Hyman. Many paratypes are SBMNH 142927. This species is placed in *Latocestus* Plehn, 1896 (Prudhoe, 1985: 47).

Amphiscolops bermudensis Hyman, 1939g: 3–5, figs. 1–6.

Paratype: AMNH 660 (1 specimen).

Locality: On rooted seaweeds, the Reach, Bermuda, Atlantic Ocean.

Collector: L. H. Hyman, Aug. 1935.

Remarks: ex Hyman. The holotype was BOC 1044 (Hyman, 1939g: 5), now YPM 20136.

Anoploidium longiductum Hyman, 1960: 6–8, figs. 4–7.

Holotype: AMNH 496.

Paratypes: AMNH 497 (3 specimens), 570, 571 (1 specimen each).

Host: *Actinopyga* sp. (Holothuroidea).

Locality: Nossi-Bé, Madagascar, Indian Ocean.

Collector: A. G. Humes, June 1955.

Remarks: ex Hyman.

Anoploidium ramosum Hyman, 1960: 3–4, figs. 1–3.

Holotype: AMNH 494.

Paratypes: AMNH 495, 547 (1 specimen each), 548 (2 specimens).

Host: *Stichopus variegatus* (Holothuroidea).

Locality: Nossi-Bé, Madagascar, Indian Ocean.

Collector: A. G. Humes, June 1955.

Remarks: ex Hyman.

Bertiliella bathycola Rieger and Sterrer, 1975: 225–232, figs. 12–15.

Holotype: AMNH 812.

Paratypes: AMNH 813.1–813.4 (3 specimens).

Locality: 260 m, mud, Fanafjord, northwest of Korsnes, Espergrend, Norway, Atlantic Ocean.

Collector: Unknown, Oct. 17, 1964.

Remarks: ex Rieger and Sterrer. The repository of additional paratypes was not cited. This species is the type of the genus *Bertiliella* Rieger and Sterrer, 1975, by monotypy (Rieger and Sterrer, 1975: 232).

Bipalium adventitium Hyman, 1943: 18, figs. 23–25.

Holotype: AMNH 302.

Paratype: AMNH 661.1–661.6 (1 specimen).

Locality: Garden, Pasadena, Los Angeles Co. (AMNH 302); campus of the University of California, Berkeley, Contra Costa Co. (AMNH 661), California, United States.

Collector: M. W. de Laubenfels (AMNH 302); J. L. Mohr (AMNH 661).

Remarks: ex Hyman. AMNH 302 was fig. 23 in Hyman (1943). The repository of additional paratypes was not cited.

Bipalium costaricensis Hyman, 1939a: 430–431, figs. 49e, 50a.

Paratype: AMNH 662 (1 specimen [part]).

Locality: Gardens, 1160 m, San José, Costa Rica.

Collector: M. Valerio, June 15, 1929.

Remarks: ex Hyman. The holotype, fig. 49e in Hyman (1939a), is USNM 20409 and the remainder of the single paratype is USNM 20410 (Hyman, 1939a: 431). The specific name was emended to *costaricense* by Froelich, 1967 (see Ogren and Kawakatsu, 1987: 83–84), but neither spelling has been used consistently. This species may be a synonym of *B. kewense* Moseley, 1878 (Ogren and Kawakatsu, 1987: 84).

Bipalium pennsylvanicum Ogren, 1987: 23–27, figs. 1–8.

Holotype: AMNH 1164.

Paratype: AMNH 1165 (1 specimen).

Locality: Eastern College, St. Davids, Delaware Co., Pennsylvania, United States.

Collector: R. E. Ogren and J. K. Sheldon, June 23, 1984.

Remarks: ex Ogren. Additional paratypes are ANSP WO4521 and in the author's collection (Ogren, 1987: 26). This specific name is a nomen nudum from Kawakatsu (1986: 6).

Bradynectes sterreri Rieger, 1971a: 206–224, figs. 1–14.

Holotype: AMNH 742.1, 742.2.

Paratypes: AMNH 743.1–743.3 (2 specimens), 744.1, 744.2, 745.1, 745.2 (1 specimen each).

Locality: Kristineberg, near Klubban Station, Sweden (AMNH 742, 743); Beaufort, Carteret Co., North Carolina, United States

(AMNH 744); Robin Hood's Bay, Yorkshire, England (AMNH 745), Atlantic Ocean.

Collector: W. Sterrer, Sept. 24, 1964 (AMNH 742, 743); R. M. Rieger, April 1970 (AMNH 744), Aug. 1969 (AMNH 745).

Remarks: ex Rieger. The repository of additional paratypes was not cited. This species is the type of the genus *Bradynectes* Rieger, 1971, by monotypy (Rieger, 1971a: 232).

Cirrifera xanthoderma Riser, 1981: 141, 143, fig. 2i.

Holotype: AMNH 936.

Locality: Manomet Beach, Manomet, Plymouth Co., Massachusetts, United States, Atlantic Ocean.

Collector: N. W. Riser, May 19, 1980.

Remarks: ex Riser. The repository of the paratypes was not cited.

Coelogynopora erotica Riser, 1981: 141, figs. 1h, k.

Holotype: AMNH 935.

Locality: Liberty Point, Robbinston, Maine, United States, Atlantic Ocean.

Collector: N. W. Riser, Nov. 1974.

Remarks: ex Riser. The repository of the paratypes was not cited.

Coelogynopora hymanae Riser, 1981: 140, fig. 1i.

Holotype: AMNH 930.

Paratypes: AMNH 931.1, 931.2 (2 specimens).

Locality: 8 m from cliff face, Brandy Cove, St. Andrews, New Brunswick, Canada, Atlantic Ocean.

Collector: N. W. Riser, Jan. 6, 1981 (AMNH 930), March 1980 (AMNH 931).

Remarks: ex Riser. The repository of additional paratypes was not cited.

Coelogynopora juxtaforcipis Riser, 1981: 140–141, figs. 1g, j.

Holotype: AMNH 932.

Locality: Dionis Beach, Nantucket Co., Massachusetts, United States, Atlantic Ocean.

Collector: N. W. Riser.

Remarks: ex Riser. The repository of the paratypes was not cited.

Coelogynopora poaceaglandis Riser, 1981: 141, figs. 1c, f, 2a, e.

Holotype: AMNH 933.

Paratype: AMNH 934 (1 specimen).

Locality: Plum Island, Essex Co., Massachusetts, United States, Atlantic Ocean.

Collector: N. W. Riser, Sept. 1980 (AMNH 933), April 22, 1978 (AMNH 934).

Remarks: ex Riser.

Comprostatum insularis Hyman, 1944: 7, figs. 14–16.

Holotype: AMNH 333–337.

Paratypes: AMNH 559.1–559.3 (1 specimen), 560 (2 specimens).

Locality: Under rocks in shallow water, Key Largo, Monroe Co., Florida, United States, Atlantic Ocean.

Collector: H. Humm, 1940.

Remarks: ex Hyman. The specific name was erroneously cited by Hyman (1944) as “*venus*” in fig. 16. This species is the type of the genus *Comprostatum* Hyman, 1944, by monotypy (Hyman, 1944: 7). This species is a synonym of *Phaenocelis purpurea* (Schmarda, 1859) (Prudhoe, 1985: 155).

Crassicollum musculare Dean, 1977: 269–273, figs. 5–12.

Holotype: AMNH 870.

Paratype: AMNH 871 (1 specimen).

Locality: Intertidal coarse sand, Crowe Neck, North Trescott (AMNH 870); intertidal coarse sand, Reid State Park, Georgetown (AMNH 871), Maine, United States, Atlantic Ocean.

Collector: H. K. Dean, Dec. 1976 (AMNH 870), Sept. 10, 1976 (AMNH 871).

Remarks: ex Dean. The holotype slide data given above do not match the published type locality or “type date.” Additional paratypes are NRS 2954, 2955 (Dean, 1977: 273). This species is the type of the genus *Crassicollum* Dean, 1977, by monotypy (Dean, 1977: 273).

Cryptocelis occidentalis Hyman, 1953a: 294–296, figs. 28–31.

Paratype: AMNH 663.1–663.5 (1 specimen).

Locality: 39 fathoms, on bottom of gray sand, near Santa Cruz Island, Santa Cruz Co., California, United States, Pacific Ocean.

Collector: Allan Hancock Foundation, Oct. 31, 1940.

Remarks: ex Hyman. The holotype was in AHF (Hyman, 1953a: 296), now SBMNH 142922. The additional paratype is SBMNH 142928.

Cystiplana rubra Dean, 1977: 265–269, figs. 1–3.

Holotype: AMNH 867.

Paratypes: AMNH 868, 869 (1 specimen each).

Locality: Intertidal coarse sand, Griffith's Head Beach, Reid State Park, Georgetown (AMNH 867); intertidal coarse sand, Crowe Neck, North Trescott (AMNH 868), Maine; subtidal, 20 m, off East Point, Nahant, Essex Co., Massachusetts (AMNH 869), United States, Atlantic Ocean.

Collector: H. K. Dean, March 9, 1976 (AMNH 867), Sept. 27, 1976 (AMNH 868, 869).

Remarks: ex Dean. The holotype slide data given above do not match the published type locality. Additional paratypes are NRS 2952, 2953 (Dean, 1977: 269).

Dalyella viridis carolinensis Hyman, 1955b: 2–10, figs. 1–9.

Holotype: AMNH 445.

Paratypes: AMNH 446.1–446.5 (3 specimens), 664.1–664.18 (41 specimens).

Locality: Vernal pool in meadow of flood plain, Rocky River, Cabarrus Co., North Carolina, United States.

Collector: T. Daggy, Feb. 1952.

Remarks: ex Hyman. This subspecies was placed in a list of *D. viridis* synonyms by Smith (1991: 257–259).

Dendrocoelopsis vaginatus Hyman, 1935: 339–344, figs. 1–5.

Holotype: AMNH 665.1, 665.2.

Paratypes: AMNH 666.1–666.14 (3 specimens).

Locality: Along shore, Flathead Lake, Montana, United States.

Collector: R. T. Young, March and July, 1933.

Remarks: ex Hyman. AMNH 665 was figs. 1–4 in Hyman (1935).

Desmorhynchus angustus Hyman, 1941: 10, figs. 15–17.

Holotype: AMNH 288.

Paratypes: AMNH 667.1, 667.2 (2 specimens).

Locality: Barro Colorado Island, Panama.

Collector: E. C. Williams Jr., July 1938.

Remarks: ex Hyman. The repository of additional paratypes was not cited. This spe-

cies is placed in *Rhynchodemus* Leidy, 1851 (Ogren and Kawakatsu, 1988: 68).

Diplandros singularis Hyman, 1953a: 341–342, figs. 101–105.

Paratype: AMNH 668.1–668.3 (1 specimen).

Locality: Ensenada, Baja California, Mexico, Pacific Ocean.

Collector: E. F. Ricketts, May 31, 1939.

Remarks: ex Hyman. AMNH 668 was fig. 105 in Hyman (1953a). The holotype was NHMSU 5639 (Hyman, 1953a: 342), now CASIZ 021626. An additional paratype is SBMNH 142929. This species is the type of the genus *Diplandros* Hyman, 1953, by monotypy (Hyman, 1953a: 341).

Diporodemus indigenus Hyman, 1943: 15–17, figs. 15–18, 20.

Paratypes: AMNH 300, 669.1–669.3, 670.1–670.12 (1 specimen each).

Locality: 1830 ft, deep leaf mold, Sky Line Drive, Blue Ridge Mountains, near Fort Royal (AMNH 300); under a board in a grassy pasture, Bolding's Spring, Staunton, Augusta Co. (AMNH 669), Virginia; near the Potomac River, Georgetown, Washington, D.C. (AMNH 670), United States.

Collector: J. P. E. Morrison, Aug. 25, 1940 (AMNH 300) and Oct. 9, 1938 (AMNH 669); L. Hubricht, March 8, 1938 (AMNH 670).

Remarks: ex USNM 157515 (AMNH 300), 150650 (AMNH 669) and Hyman Collection via Hyman. The holotype, fig. 15 in Hyman (1943), is USNM 20616 and an additional paratype is USNM 20617 (Hyman, 1943: 17). The repository of additional paratypes was not cited.

Diporodemus plenus Hyman, 1941: 9–10, figs. 11–14.

Holotype: AMNH 287.1–287.8.

Locality: Barro Colorado Island, Panama.

Collector: E. C. Williams Jr., July 1938.

Remarks: ex Hyman. AMNH 287 is the holotype by monotypy.

Diporodemus yucatani Hyman, 1938a: 26–31, figs. 3–9.

Syntypes: AMNH 640.1–640.11 (2 specimens).

Locality: Under stones near mouth of Sabacha Cave, Tekax, Yucatan, Mexico.

Collector: A. S. Pearse, July 30, 1936.

Remarks: ex Hyman. This species is the type of the genus *Diporodemus* Hyman, 1938, by monotypy (Hyman, 1938a: 32).

Discocelis insularis Hyman, 1955d: 66–70, figs. 1a, b, 2a, c.

Paratype: AMNH 671.1–671.4 (1 specimen).

Locality: Raroia Atoll, Tuamotu Islands, French Polynesia, Pacific Ocean.

Collector: J. P. E. Morrison, Sept. 2, 1952.

Remarks: ex Hyman. The holotype is USNM 25939 (Hyman, 1955d: 70). This species is placed in *Thalamoplanea* Laidlaw, 1903 (Faubel, 1983: 46).

Dugesia arimana Hyman, 1957b: 2–4, figs. 1, 2.

Holotype: AMNH 465.1, 465.2.

Paratypes: AMNH 673 (5 specimens).

Locality: Under stones, Arima River, between Simla and Arima, Trinidad, Trinidad and Tobago.

Collector: L. H. Hyman, Feb. 14, 1956.

Remarks: ex Hyman. This species is placed in *Girardia* Ball, 1974 (Sluys, 1992: 13).

Dugesia diabolis Hyman, 1956: 5–8, figs. 1, 5–7.

Holotype: AMNH 468.1, 468.2.

Paratypes: AMNH 469, 674.1–674.3 (1 specimen each).

Locality: Devil's Hole, 70 mi west of Las Vegas, Clark Co., Nevada, United States.

Collector: E. E. Simmons, about Dec. 2, 1954.

Remarks: ex Hyman. This species is a synonym of *Girardia dorotocephala* (Woodworth, 1897) (Kenk, 1989: 2).

Dugesia sanchezi Hyman, 1959c: 2–7, figs. 1–3, 5–11.

Holotype: AMNH 488.1, 488.2.

Paratypes: AMNH 489.1–489.4 (3 specimens), 675.1–675.7 (4 specimens).

Locality: Rio Mapocho, near Talagante, Santiago Province, Chile.

Collector: P. Sanchez.

Remarks: ex Hyman. This species is a synonym of *Girardia festai* (Borelli, 1898) (Sluys, 1992: 5).

Dugesia titicacana Hyman, 1939a: 419–422, figs. 47a–c.

Paratype: AMNH 676.1–676.4 (1 specimen).

Locality: Isla de la Sol, Lake Titicaca, Bolivia.

Collector: M. C. James, Feb. 18, 1936.

Remarks: ex Hyman. The holotype is USNM 20402 and was the figured specimen (Hyman, 1939a: 422). AMNH 676 is the sexual specimen without copulatory apparatus cited by Hyman (1939a: 419). The repository of the additional paratype was not cited. This species is a synonym of *Girardia festai* (Borelli, 1898) (Sluys, 1992: 5).

Ectocotyla multitesticulata Fleming and Burt, 1978: 1227–1230, figs. 4, 5.

Paratypes: AMNH 918.1, 918.2, 919 (1 specimen each).

Host: *Chionoecetes opilio* (Malacostraca).

Locality: 40 fathoms, off the coast of Newfoundland (AMNH 918); off the coast of Caraquet, Bay of Chaleur, New Brunswick (AMNH 919), Canada, Atlantic Ocean.

Collector: L. C. Fleming and M. D. B. Burt, Jan. 1974 (AMNH 918), May 26, 1974 (AMNH 919).

Remarks: ex Fleming and Burt. The holotype is BM(NH) 1978.4.7.1 and an additional paratype is BM(NH) 1978.4.7.2 (Fleming and Burt, 1978: 1230).

Ectocotyla paguri Hyman, 1944: 2–3, figs. 4–8.

Syntypes: AMNH 340 (1 specimen), 341 (11 specimens), 549, 550 (1 specimen each).

Host: *Pagurus pubescens* (Malacostraca).

Locality: Biological station of University of Maine, Ellsworth, Hancock Co., Maine, United States, Atlantic Ocean.

Collector: E. G. Reinhard, Aug. 1940.

Remarks: ex Hyman. This species is the type of the genus *Ectocotyla* Hyman, 1944, by monotypy (Hyman, 1944: 2), but is a synonym of *E. hirudo* (Levinsen, 1879) (Fleming and Burt, 1978: 1224).

Enchiridium punctatum Hyman, 1953a: 386–387, figs. 160, 161.

Holotype: AMNH 390.

Locality: 8–15 m, Newport Bay, California, United States, Pacific Ocean.

Collector: G. E. and N. MacGinitie, June 12, 1938.

Remarks: ex Hyman. A paratype is SBMNH 142930.

Endocincta punctata Crezée, 1975: 810–814, figs. 16, 17, 21a, b, d, e.

Holotype: AMNH 843.

Paratypes: AMNH 844–849 (1 specimen each).

Locality: Sand flats, Wrightsville Beach, New Hanover Co. (AMNH 843–845); end of 28th Street, Bogue Sound, Morehead City, Carteret Co. (AMNH 846, 847); near Swansboro, Onslow Co. (AMNH 848); Cape Lookout, Carteret Co. (AMNH 849), North Carolina, United States, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Crezée. The repository of additional paratypes was not cited. This species is the type of the genus *Endocincta* Crezée, 1975, by monotypy (Crezée, 1975: 815).

Euplanaria microbursalis Hyman, 1931: 339–340, figs. 6–8.

Syntypes: AMNH 638.1–638.10 (3 specimens).

Locality: Running stream under rocks, Mystic, New London Co., Connecticut, United States.

Collector: L. H. Hyman, Sept. 10, 1930.

Remarks: ex Hyman. This species is placed in *Girardia* Ball, 1974 (Curino and Cazaniga, 1993: 642).

Euplanaria philadelphica Hyman, 1931: 336–337, figs. 1–5.

Syntypes: AMNH 639.1–639.24 (4 specimens).

Locality: Under stones in moderate current, Wissahicken Creek, Philadelphia, Pennsylvania, United States.

Collector: L. H. Hyman, Sept. 17, 1930.

Remarks: ex Hyman. This species is a synonym of *Girardia dorocephala* (Woodworth, 1897) (Kenk, 1989: 2).

Eurylepta californica Hyman, 1959d: 11–12, figs. 15, 16.

Holotype: AMNH 491.

Locality: Rocky shore in coralline algae, near Cambria, San Luis Obispo Co., California, United States, Pacific Ocean.

Collector: D. Montgomery, Feb. 28, 1958.

Remarks: ex Hyman. AMNH 491 is the holotype by monotypy.

Eurylepta leoparda Freeman, 1933: 138–140, figs. 30–35.

Holotype: AMNH 389.

Locality: 135 fathoms, off Speiden Island, Puget Sound, Washington, United States, Pacific Ocean.

Collector: Unknown.

Remarks: ex Freeman Collection via Hyman. AMNH 389 is the holotype by monotypy.

Euryleptodes insularis Hyman, 1953a: 375–376, figs. 146, 147.

Paratype: AMNH 677.1–677.4 (1 specimen).

Locality: 20 m, off Santa Rosa Island, California, United States, Pacific Ocean.

Collector: Allan Hancock Foundation, Aug. 10, 1939.

Remarks: ex Hyman. The holotype was in AHF (Hyman, 1953a: 376), now SBMNH 142921. The repository of the additional paratype was not cited and it could not be located in SBMNH.

Florianella bipolaris Rieger and Sterrer, 1975: 208–222, 225, figs. 1–11.

Holotype: AMNH 810.1–810.3.

Paratypes: AMNH 811.1–811.5 (2 specimens).

Locality: 32 m, 34°16.7'N, 76°24.7'W, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: B. Coull, April 1970.

Remarks: ex Rieger and Sterrer. The repository of additional paratypes was not cited. This species is the type of the genus *Florianella* Rieger and Sterrer, 1975, by monotypy (Rieger and Sterrer, 1975: 225).

Fusantrum rhamniphorum Crezée, 1975: 804–809, figs. 11c, d, 14, 15.

Holotype: AMNH 838.

Paratypes: AMNH 839–842 (1 specimen each).

Locality: Sand flats, Wrightsville Beach, New Hanover Co. (AMNH 838–840); estuary, White Oak River, opposite Swansboro, Onslow Co. (AMNH 841, 842), North Carolina, United States, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Crezée. The repository of additional paratypes was not cited. This species is the type of the genus *Fusantrum* Crezée, 1975, by monotypy (Crezée, 1975: 809).

Geocentrophora tropica Hyman, 1941: 1–5, figs. 1–4.

Syntypes: AMNH 283 (3 specimens), 678.1–678.3 (1 specimen).

Locality: Forest floor, Barro Colorado Island, Panama.

Collector: E. C. Williams Jr., July 1938.

Remarks: ex Hyman. The repository of additional syntypes was not cited.

Geoplana andina Hyman, 1962: 7–10, figs. 6–10.

Holotype: AMNH 505.

Paratypes: AMNH 506.1–506.24 (2 specimens).

Locality: Under logs in forest, 1500 m, Bitate, Departamento del Valle, Colombia.

Collector: E. Kozloff, June 20, 1954.

Remarks: ex Hyman. This species is placed in *Notogynaphallia* Ogren and Kawakatsu, 1990 (Ogren and Kawakatsu, 1990: 140).

Geoplana aphalla Hyman, 1941: 6–9, figs. 6, 10.

Holotype: AMNH 286.1, 286.2.

Locality: Forest floor, Barro Colorado Island, Panama.

Collector: E. C. Williams Jr., July 1938.

Remarks: ex Hyman. AMNH 286 is the holotype by monotypy. This species is placed in *Pasipha* Ogren and Kawakatsu, 1990 (Ogren and Kawakatsu, 1990: 145).

Geoplana arkalabamensis Ogren and Darlington, 1991: 228–232, figs. 1–14.

Holotype: AMNH 1593.

Locality: Garden, 513 Johnson Avenue, Monroeville, Monroe Co., Alabama, United States.

Collector: L. Hubricht, July 29, 1981.

Remarks: ex Ogren. Paratypes are in the senior author's collection and "will be deposited" at AMNH (Ogren and Darlington, 1991: 232); they had not been at the time of this writing.

Geoplana contaminensis Hyman, 1955e: 22–23, figs. 29–31.

Holotype: AMNH 449.1–449.9.

Locality: 300 m, Contamana, Ucayali River, Peru.

Collector: American Museum Expedition, March 1927.

Remarks: ex Hyman. AMNH 449 is the holotype by monotypy. This species is placed in *Amaga* Ogren and Kawakatsu, 1990 (Ogren and Kawakatsu, 1990: 93).

Geoplana panamensis Hyman, 1941: 6, fig. 5.

Holotype: AMNH 285.

Locality: Forest floor, Barro Colorado Island, Panama.

Collector: E. C. Williams Jr., 1938.

Remarks: ex Hyman. The repository of the paratypes was not cited. This species is placed in *Pseudogeoplana* Ogren and Kawakatsu, 1990 (Ogren and Kawakatsu, 1990: 158).

Geoplana septemlineata Hyman, 1939b: 116–119, figs. 1–3.

Holotype: AMNH 631.

Paratypes: AMNH 275 (4 specimens), 632.1–632.10 (2 specimens).

Locality: Wet forests under stones and fallen leaves, 2400 ft, Kipapa Ridge; Lulumahu Mountain, 1800 ft; Koolan Mountains, 1800 ft, Oahu, Hawaii, United States.

Collector: F. X. Williams, Sept. 5, 1938.

Remarks: ex Hyman. This species is the type of the genus *Endeavouria* Ogren and Kawakatsu, 1991, by monotypy (Ogren and Kawakatsu, 1991: 69).

Geoplana subpallida Hyman, 1939b: 119–120, fig. 4.

Holotype: AMNH 633.

Locality: Wet forests, 1800 ft, Koolan Mountains, Oahu, Hawaii, United States.

Collector: F. X. Williams, 1938.

Remarks: ex Hyman. AMNH 633 is the holotype by monotypy. This species is placed in *Australopacifica* Ogren and Kawakatsu, 1991 (Ogren and Kawakatsu, 1991: 56).

Geoplana ucayalensis Hyman, 1955e: 24, figs. 32–35.

Holotype: AMNH 451.

Locality: 300 m, Contamana, Ucayali River, Peru.

Collector: American Museum Expedition, March 1927.

Remarks: ex Hyman. AMNH 451 is the holotype by monotypy. This species is placed in *Pseudogeoplana* Ogren and Kawakatsu, 1990 (Ogren and Kawakatsu, 1990: 161).

Geoplana vaga Hyman, 1943: 3, figs. 3–5.

Holotype: AMNH 294.1–294.7.

Locality: Garden, Pasadena, Los Angeles Co., California, United States.

Collector: W. M. de Laubenfels.

Remarks: ex Hyman. AMNH 294 is the holotype by monotypy. This taxon is a subspecies of *Caenoplana coerulea* Moseley, 1877 (Ogren and Kawakatsu, 1991: 63).

Girardia canai Curino and Cazzaniga, 1993: 633–639, figs. 1–9.

Paratypes: AMNH 1595.1–1595.9, 1596.1–1596.12 (1 specimen each).

Locality: Mapocho River, near Talagante, Region Metropolitana (AMNH 1595) and Estero Renaca, south of Conçon, V Region (AMNH 1596), Chile.

Collector: A. C. Curino, N. J. Cazzaniga, and P. Sanchez, Oct. 26, 1986.

Remarks: ex Curino and Cazzaniga. The holotype is MACN 33438 and additional paratypes are MACN 33439 and in UNS (Curino and Cazzaniga, 1993: 634).

Gnesioceros sargassicola var. *lata* Hyman, 1939g: 13, fig. 18.

Paratypes: AMNH 679.1–679.10 (3 specimens).

Locality: Among rooted seaweeds, the Reach, Bermuda, Atlantic Ocean.

Collector: L. H. Hyman, July–Aug. 1935.

Remarks: ex Hyman. The holotype was BOC 1051 (Hyman, 1939g: 13), now YPM 20147. As ICZN (1985) does not allow for varieties, this taxon should be considered a subspecies as *G. sargassicola lata*.

Gnesioceros verrilli Hyman, 1939c: 146–148, figs. 14–16.

Paratypes: AMNH 680.1–680.8 (4 specimens).

Locality: Eel grass, Devil's Foot Island, Woods Hole, Barnstable Co., Massachusetts, United States, Atlantic Ocean.

Collector: Unknown, Oct. 28, 1931.

Remarks: ex Hyman. This species was described as a "new name" for *Imogine oculifera* (Girard, 1853) of Verrill, 1892 (Hyman, 1939c: 146). This was actually a description of a new species and the "neotype" selected by Hyman and deposited at MBL must be considered the holotype. This species is considered a synonym of *G. sargassicola* (Mertens, 1832) by Prudhoe (1985: 166), but of *G. floridana* (Pearse, 1938) by Faubel (1983: 118).

Hoploplana californica Hyman, 1953a: 344, figs. 106, 107.

Holotype: AMNH 379.

Paratype: AMNH 575.1–575.4 (1 specimen).

Locality: 40–50 ft, Corona Del Mar, Orange Co., California, United States, Pacific Ocean.

Collector: G. E. and N. MacGinitie, July 9, 1947.

Remarks: ex Hyman. The repository of additional paratypes was not cited and they could not be located in SBMNH.

Hoploplana luracola Smith, 1961: 69–70, figs. 1–4.

Holotype: AMNH 501.

Host: *Nerita* (*Ritena*) *scabricosta ornata* (Gastropoda).

Locality: 09°00'N, 79°30'W, Balboa Park, Panama City, Panama, Pacific Ocean.

Collector: E. H. Smith, Dec. 1960.

Remarks: ex Smith. Paratypes were cited as being in USNM and the author's collection (Smith, 1961: 70), but none can be located in USNM (Littman, personal commun.).

Hoploplana usaguia Smith, 1960: 385–387, figs. 1–4.

Paratype: AMNH 499 (1 specimen).

Locality: 23°27'S, 14 km west of Ubatuba, São Paulo, Brazil, Atlantic Ocean.

Collector: E. Smith, July 1960.

Remarks: ex Smith. The holotype is in Smith's personal collection and additional paratypes are USNM 30249 and in USP (Smith, 1960: 387).

Indistylachus hewatti Hyman, 1955c: 123–124, figs. 81d, 83a.

Paratype: AMNH 681 (1 specimen [part]).

Locality: Under beds of *Mytilus*, between tides, East Point Beach, Boqueron, Puerto Rico, Atlantic Ocean.

Collector: W. G. Hewatt, March 13, 1946.

Remarks: ex Hyman. The holotype is USNM 24620 and the paratype is USNM 24621 (Hyman, 1955c: 124). AMNH 681 is the copulatory region of USNM 24621. This species is the type of the genus *Indistylachus* Hyman, 1955, by monotypy (Hyman, 1955c: 123). This species is placed in *Styl-ochocestus* Laidlaw, 1903 (Faubel, 1983: 52).

Kenkia rhynchida Hyman, 1937b: 457–462, figs. 1–6.

Syntypes: AMNH 641.1–641.10 (3 specimens).

Locality: Malheur Cave, Oregon, United States.

Collector: C. L. Hubbs, July 27–28, 1934.

Remarks: ex Hyman. The additional syntype is USNM 20229. This species is the type

of the genus *Kenkia* Hyman, 1937, by monotypy (Hyman, 1937b: 473).

Kytorhynchella karlingi Rieger, 1974: 55, figs. 1i, 4a, 6a-c.

Holotype: AMNH 748.

Locality: In fine sand, 11 m, in front of Shackleford Banks, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: R. M. Rieger.

Remarks: ex Rieger. The repository of the paratypes was not cited.

Kytorhynchella meixneri Rieger, 1974: 54-55, figs. 1a-g, 2a-c, 6d, 8a, 9d, 12a, 13a-c, 14b, 15e, 16a, b.

Holotype: AMNH 746.1, 746.2.

Paratypes: AMNH 747.1-747.12 (6 specimens).

Locality: Sand flat, 300 ft north of the drawbridge on Highway 76, Wrightsville Beach, New Hanover Co., North Carolina, United States, Atlantic Ocean.

Collector: R. M. Rieger.

Remarks: ex Rieger. The repository of additional paratypes was not cited. This species is the type of the genus *Kytorhynchella* Rieger, 1974, by original designation (Rieger, 1974: 54).

Kytorhynchella riedli Rieger, 1974: 55, figs. 1h, 8b, 14a, 15f.

Holotype: AMNH 749.1, 749.2.

Locality: Station 14207, 32 m, 34°16'N, 76°24'W, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: R/V "Eastward" cruise E-4-70, April 1970.

Remarks: ex Rieger. The repository of the paratype was not cited.

Kytorhynchus (Kytorhynchoides) macrostylus Rieger, 1974: 57-58, figs. 5b, 6f, 9e, 11b, 12e, 15a, b, 16c.

Holotype: AMNH 755.1, 755.2.

Locality: Station 14460, 20 m, 34°25.5'N, 76°20'W, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: R/V "Eastward" cruise E-10-70, May 1970.

Remarks: ex Rieger. The repository of the paratype was not cited. The subgenus *Kytorhynchoides* Rieger, 1974, is not valid and is an unavailable name because no type

species was designated in the original publication (Rieger, 1974: 57), an action made mandatory by ICZN article 13b (ICZN, 1985: 37). This species is placed in the subgenus *Eukytorhynchus* new name for *Kytorhynchoides* Rieger, 1974 (see *Kytorhynchus microstylus* for type species designation).

Kytorhynchus (Kytorhynchoides) microstylus Rieger, 1974: 57, figs. 4c, 5a, c-e, 7a-g, 9c, f, 11a, 12d, 14d, f, 15d.

Holotype: AMNH 753.

Paratypes: AMNH 754A, 754B (1 specimen each).

Locality: Station 14457, 30 m, 34°18'N, 73°16'W, off Beaufort, Carteret Co., North Carolina, United States (AMNH 753); 15 m, Tobacco Bay, Bermuda (AMNH 754), Atlantic Ocean.

Collector: R/V "Eastward" cruise E-10-70, May 1970 (AMNH 753); R. M. Rieger (AMNH 754).

Remarks: ex Rieger. The repository of additional paratypes was not cited. The subgenus *Kytorhynchoides* Rieger, 1974, is not valid and is an unavailable name because no type species was designated in the original publication (Rieger, 1974: 57), an action made mandatory by ICZN Article 13b (ICZN, 1985: 37). This species is hereby designated the type of the subgenus *Eukytorhynchus* new name for *Kytorhynchoides* Rieger, 1974 (see Rieger, 1974: 57 for diagnosis of both the species and subgenus).

Kytorhynchus (Kytorhynchus) oculatus Rieger, 1974: 56, figs. 3a, 4b, 9a, b, 10, 12b, 14c, 15c.

Holotype: AMNH 750.

Paratype: AMNH 751 (1 specimen).

Locality: Station 14207, 28 m, 34°12'N, 76°14'W, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: R/V "Eastward" cruise E-4-70, April 1970.

Remarks: ex Rieger. The repository of additional paratypes was not cited. This species is the only named one in the subgenus *Kytorhynchus* Rieger, 1974, and is therefore the type of the subgenus by monotypy. Since the nominotypical subgenus contains only a single species, this species is also the type of the genus *Kytorhynchus* Rieger,

- 1974, by monotypy as per ICZN Article 68d(i) (ICZN, 1985: 129–130).
- Latocestus brasiliensis* Hyman, 1955e: 12, figs. 19, 20.
Paratype: AMNH 682 (1 specimen).
Locality: Off São Francisco, Brazil, Atlantic Ocean.
Collector: W. Schmitt, Oct. 30, 1925.
Remarks: ex Hyman. The holotype is USNM 26929 (whole mount) and the additional paratype is USNM 26929a (sections) (Hyman, 1955e: 14).
- Latocestus galapagensis* Hyman, 1953b: 183–185, figs. 1, 2.
Paratype: AMNH 683 (1 specimen).
Locality: Galápagos Islands, Ecuador, Pacific Ocean.
Collector: R/V "Velero III," 1933–1934.
Remarks: ex Hyman. The holotype was in AHF (Hyman, 1953b: 183), now SBMNH 142932. The repository of additional paratypes was not cited and they could not be located in SBMNH. This species is placed in *Eulatocestus* Faubel, 1983 (Faubel, 1983: 64).
- Leptoplana inquieta* Heath and McGregor, 1912: 470–472, figs. 8, 14, 43.
Neotype: AMNH 369.
Locality: Monterey Bay, California, United States, Pacific Ocean.
Collector: Unknown.
Remarks: ex Hyman. The neotype (fig. 1, herein) was designated by Hyman (1953a: 321) because none of the four syntypes could be located. This species is placed in *Pleioiplana* Faubel, 1983 (Faubel, 1983: 117).
- Leptoplana limnoriae* Hyman, 1953a: 313–314, figs. 57–59.
Paratype: AMNH 545.1–545.4 (1 specimen).
Locality: In burrows of *Limnoria* [Isopoda], California Yacht Club, Los Angeles Harbor, Los Angeles Co., California, United States, Pacific Ocean.
Collector: D. Reish, 1950.
Remarks: ex Hyman. The holotype was in AHF (Hyman, 1953a: 314), now SBMNH 142923. The repository of additional paratypes was not cited and they could not be located in SBMNH. This species is placed in *Stylochoplana* Stimpson, 1857 (Prudhoe, 1985: 171).
- Leptostylochus novacambrensis* Hyman, 1959b: 3–5, figs. 5, 6.
Paratypes: AMNH 684.1–684.4 (2 specimens).
Locality: Intertidal zone, Long Reef, near Collaroy, New South Wales, Australia, Pacific Ocean.
Collector: E. Pope, 1955–1956.
Remarks: ex Hyman. The holotype is AM W.3686 (Hyman, 1959b: 5). The repository of additional paratypes was not cited.
- Longiprostatum rickettsi* Hyman, 1953a: 300–301, figs. 37, 38.
Holotype: AMNH 363.1–363.4.
Locality: Angeles Bay, Gulf of California, Mexico, Pacific Ocean.
Collector: E. F. Ricketts, April 1, 1940.
Remarks: ex Hyman. AMNH 363 is the holotype by monotypy. This species is the type of the genus *Longiprostatum* Hyman, 1953, by monotypy (Hyman, 1953a: 300).
- Macginitella delmaris* Hyman, 1953a: 337–339, figs. 95, 96.
Holotype: AMNH 377.1–377.5.
Locality: Corona del Mar, Orange Co., California, United States, Pacific Ocean.
Collector: B. Ewing, Feb. 1948.
Remarks: ex Hyman. AMNH 377 is the holotype by monotypy. This species is the type of the genus *Macginitella* Hyman, 1953, by monotypy (Hyman, 1953a: 337).
- Macroatrium setosum* Riser, 1981: 143, 145, figs. 2b–d, f, g.
Holotype: AMNH 937.
Paratype: AMNH 938 (1 specimen).
Locality: Dionis Beach, Nantucket Co. (AMNH 937); Folly Cove, Anisquan (AMNH 938), Massachusetts, United States, Atlantic Ocean.
Collector: N. W. Riser, Dec. 1976 (AMNH 937), April 6, 1977 (AMNH 938).
Remarks: ex Riser. The repository of additional paratypes was not cited. This species is the type of the genus *Macroatrium* Riser, 1981, by monotypy (Riser, 1981: 143).
- Macrocotyla glandulosa* Hyman, 1956: 9–14, figs. 8–17.
Holotype: AMNH 470.

Paratypes: AMNH 471 (7 specimens), 685.1–685.10, 686.1, 686.2 (2 specimens each), 687.1–687.16 (3 specimens).

Locality: Stream, below outlet of a cave, Rock Bridge, 5 mi south of Columbia, Boone Co., Missouri, United States.

Collector: P. W. Frank, April 30, 1955.

Remarks: ex Hyman. The species name was misspelled “*glandularis*” in the figure captions. This species is the type of the genus *Macrocotyla* Hyman, 1956, by monotypy (Hyman, 1956: 8). This species is placed in *Kenkia* Hyman, 1937 (Kenk, 1989: 5).

Marcusia ernesti Hyman, 1953a: 296–297, figs. 32, 33.

Holotype: AMNH 361.1–361.4.

Paratypes: AMNH 542, 543.1–543.5 (1 specimen each).

Locality: Point Marcial Reef (AMNH 543); Gulf of California (AMNH 361, 542), Mexico, Pacific Ocean.

Collector: G. E. MacGinitie, Dec. 1947–Feb. 1948 (AMNH 361); Unknown (AMNH 542); E. F. Ricketts, March 1940 (AMNH 543).

Remarks: ex Hyman. Additional paratypes are SBMNH 142926. This species is the type of the genus *Marcusia* Hyman, 1953, by monotypy (Hyman, 1953a: 296). This species is placed in *Pericelis* Laidlaw, 1902, by Prudhoe (1985: 175) but is retained in *Marcusia* by Faubel (1984: 199).

Mesostoma andrewsi Hyman, 1957a: 2–6, figs. 1–7.

Holotype: AMNH 472.

Paratype: AMNH 688 (1 specimen).

Locality: Freshwater pools on wet tundra and in vegetation along streams, St. Matthew’s Island, United States.

Collector: R. Rausch, Aug.–Sept. 1954.

Remarks: ex Hyman. The repository of additional paratypes was not cited. The type locality was incorrectly given by Hyman (1957a) as “St. Andrew’s Island” (see Sluys, 1989: 186). This species is a synonym of *M. platygastricum* Hofsten, 1924 (Janssen and Faubel, 1992: 35).

Mesostoma appinum Kolasa and Schwartz, 1988: 329–330, figs. 1, 4–6.

Holotype: AMNH 1448.

Locality: Appin Pond, New South Wales, Australia.

Collector: Unknown, Nov. 1983.

Remarks: ex Kolasa and Schwartz. Paratypes are in UW (Kolasa and Schwartz, 1988: 329). This specific name was a nomen nudum from both Kolasa (1984, as *M. appia*) and Kolasa (1987).

Mesostoma arctica Hyman, 1938c: 1–8, figs. 1–6.

Holotype: AMNH 263.1, 263.2.

Paratypes: AMNH 264 (2 specimens), 689.1–689.5 (4 specimens).

Locality: Fresh water pools on the tundra, near Churchill, Manitoba, Canada.

Collector: H. E. McClure, July 1936.

Remarks: ex Hyman. The repository of additional paratypes was not cited. This specific name was emended to *arcticum* to agree with the neuter genus (Hyman, 1955b: 25).

Mesostoma californicum Hyman, 1957a: 6–10, figs. 8–15.

Holotype: AMNH 473.

Paratypes: AMNH 474.1, 474.2, 576.1–576.9 (5 specimens each), 690.1, 690.2 (4 specimens).

Locality: Temporary rain pools, near Carmel, Monterey Co., California, United States.

Collector: E. Larsen and R. Tetley, Feb. 1956.

Remarks: ex Hyman.

Mesostoma curvipenis Hyman, 1955b: 20–25, figs. 15–19.

Holotype: AMNH 448.

Paratypes: AMNH 577.1, 577.2 (2 specimens).

Locality: Vernal pool in swampy place, Brandywine Creek, New Castle Co., Delaware, United States.

Collector: R. G. Schmieder, April 15, 1945.

Remarks: ex Hyman.

Mesostoma macropenis Hyman, 1939d: 646–650, figs. 1–4.

Holotype: AMNH 578.

Paratypes: AMNH 579 (1 specimen), 580.1–580.5 (6 specimens).

Locality: Sandy bottom, 4–10 m, South Fish-tail Bay, Douglas Lake, Michigan, United States.

Collector: G. M. Moore, July–Aug. 1937.

Remarks: ex Hyman. This taxon is a subspecies of *M. ehrenbergi* (Focke, 1836) (Hyman, 1957a: 14).

Mesostoma macroprostatum Hyman, 1939f: 629–630, figs. 1–4.

Paratypes: AMNH 691.1–691.5 (3 specimens).

Locality: 9700 ft, temporary pond, Medicine Bow Mountains, Wyoming, United States.

Collector: I. H. Blake, July 2–4, 1939.

Remarks: ex Hyman. The holotype, fig. 1 in Hyman (1939f), is USNM 20527 and additional paratypes are USNM 20793, 20794 (Hyman, 1939f: 630). This species is placed in *Bothromesostoma* Braun, 1885 (Smith, 1991: 260).

Mesostoma magnum Kolasa and Schwartz, 1988: 333–334, figs. 2, 3, 7–11.

Holotype: AMNH 1449.1, 1449.2.

Locality: Pond along Bett's Creek, Kosciuszko National Park, New South Wales, Australia.

Collector: P. D. N. Hebert, Nov. 1982.

Remarks: ex Kolasa and Schwartz. Paratypes are in UW (Kolasa and Schwartz, 1988: 333).

Mesostoma vernale Hyman, 1955b: 16–20, figs. 10–14.

Holotype: AMNH 447.1, 447.2.

Paratypes: AMNH 692.1–692.5 (3 specimens).

Locality: Two adjacent pools in a meadow on the flood plain, Rocky River, Cabarrus Co., North Carolina, United States.

Collector: T. Daggy, Feb. 25, 1952.

Remarks: ex Hyman. The repository of additional paratypes was not cited.

Mexistylodus laevis Hyman, 1953a: 293–294, fig. 27.

Holotype: AMNH 360.1–360.6.

Locality: Puerto Penasco, Sonora, Mexico, Pacific Ocean.

Collector: G. E. and N. MacGinitie, Dec. 1947.

Remarks: ex Hyman. AMNH 360 is the holotype by monotypy. This species is placed in *Ommatoplana* Laidlaw, 1903, by Prudhoe (1985: 175), but is made the type of the genus *Latoplana* Faubel, 1983, by monotypy as *L. levis* (sic) (Faubel, 1983: 64).

Mexistylodus tuberculatus Hyman, 1953a: 291–293, figs. 25, 26.

Holotype: AMNH 359.

Paratype: AMNH 541.1–541.3 (1 specimen).

Locality: Miramar Beach, Guaymas, Sonora, Mexico, Pacific Ocean.

Collector: G. E. and N. MacGinitie, Feb. 9, 1948.

Remarks: ex Hyman. The additional paratype is SBMNH 142917. This species is the type of the genus *Mexistylodus* Hyman, 1953, by original designation (Hyman, 1953a: 291), but this genus is considered synonymous with *Ommatoplana* Laidlaw, 1903 (Hyman, 1955a: 9), although there is disagreement about this (see Prudhoe, 1985: 66). The combination *O. tuberculata* is preoccupied by *O. tuberculata* Laidlaw, 1903, so the new name *O. mexicana* Hyman, 1955, was proposed for this species (Hyman, 1955a: 9).

Microplana rufocephalata Hyman, 1954b: 8–11, figs. 4, 5.

Holotype: AMNH 422.1–422.11.

Paratype: AMNH 423 (1 specimen).

Locality: 2600 ft, Pine Mountain, near Harlan, Harlan Co., Kentucky, United States.

Collector: L. Hubricht, May 25, 1952.

Remarks: ex Hyman. AMNH 422 was fig. 5 and AMNH 423 was fig. 4 in Hyman (1954b).

Monocelis durhami Hyman, 1964: 248–250, figs. 1–5.

Holotype: AMNH 539.

Paratypes: AMNH 540.1–540.5 (3 specimens).

Locality: Brackish water, Durham, Strafford Co., New Hampshire, United States.

Collector: W. Bullock.

Remarks: ex Hyman. The repository of additional paratypes was not cited.

Monosolenia asymmetrica Hyman, 1953a: 359–361, figs. 127–129.

Holotype: AMNH 381.

Paratypes: AMNH 693.1–693.3 (2 specimens).

Locality: Cape San Lucas, Gulf of California, Baja California, Mexico, Pacific Ocean.

Collector: E. F. Ricketts, March 18, 1940.

Remarks: ex Hyman. This species is the type of the genus *Monosolenia* Hyman, 1953, by monotypy (Hyman, 1953a: 359).

Myomacrostromum bichaeta Rieger, 1986: 42–43, figs. 3a, c, 4b, 5a.

Holotype: AMNH 1235.

Paratype: AMNH 1236 (1 specimen).

Locality: Sandbank, 1 mi from mouth of New River Inlet, New River estuary, Onslow Co., North Carolina, United States, Atlantic Ocean.

Collector: R. M. Rieger, Summer 1972.

Remarks: ex Rieger. The repository of additional paratypes was not cited.

Myomacrostromum unichaeta Rieger, 1986: 40–42, figs. 3b, d, 4a, c, 5b, 6.

Holotype: AMNH 1233.

Paratype: AMNH 1234 (1 specimen).

Locality: Sandbank, 1 mi from mouth of New River Inlet, New River estuary, Onslow Co., North Carolina, United States, Atlantic Ocean.

Collector: R. M. Rieger, summer 1972.

Remarks: ex Rieger. The repository of additional paratypes was not cited. This species is the type of the genus *Myomacrostromum* Rieger, 1986, by original designation (Rieger, 1986: 44).

Myopea crassula Crezée, 1975: 792–799, figs. 7–10, 11a, b.

Holotype: AMNH 825.

Paratypes: AMNH 826–834 (1 specimen each).

Locality: Intertidal sand and muddy sand flats, Wrightsville Beach, New Hanover Co. (AMNH 825–830); opposite Swansboro, Onslow Co. (AMNH 831–833); Cape Lookout, Carteret Co. (AMNH 834), North Carolina, United States, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Crezée. The repository of additional paratypes was not cited. This species is the type of the genus *Myopea* Crezée, 1975, by original designation (Crezée, 1975: 804).

Myopea latafaucium Crezée, 1975: 799–803, figs. 12, 13.

Holotype: AMNH 835.

Paratypes: AMNH 836, 837 (1 specimen each).

Locality: Radio Island, opposite Morehead City, Carteret Co. (AMNH 835, 836); sand flat, Swansboro, Onslow Co. (AMNH 837), North Carolina, United States, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Crezée. The repository of additional paratypes was not cited.

Myozonaria jenneri Rieger and Tyler, 1974: 171, figs. 1b, 2b, 3a, 4b, c, 5, 6a–c, 7a, 8a, c, 9c, 10, 13, 14c, d, 15c, d.

Lectotype: AMNH 789.

Paralectotype: AMNH 790 (1 specimen).

Locality: Bahia Honda Key, Monroe Co., Florida, United States, Atlantic Ocean.

Collector: R. M. Rieger, May 1971.

Remarks: ex Rieger and Tyler. Rieger marked slide AMNH 789 “holotype.” Since no holotype was designated by Rieger and Tyler (1974), AMNH 789 is here designated lectotype. This specific name is a nomen nudum from Rieger (1971b: 249).

Nemertinoides elongatus Riser, 1987: 60–64, figs. 1–10.

Holotype: AMNH 1238.

Locality: Upper tidal zone, coarse sand, 41°49'N, 70°33'W, Ellisville Beach, Barnstable Co., Massachusetts, United States, Atlantic Ocean.

Collector: N. W. Riser, Sept. 10, 1984.

Remarks: ex Riser. A paratype is NRS 3591. The repository of additional paratypes was not cited. This species is the type of the genus *Nemertinoides* Riser, 1987, by monotypy (Riser, 1987: 60).

Nesion arcticum Hyman, 1956: 2–4, figs. 2–4.

Holotype: AMNH 466.1, 466.2.

Paratypes: AMNH 467.1, 467.2 (5 specimens), 581.1–581.6 (3 specimens).

Locality: Intertidal zone, St. Matthew's Island, United States.

Collector: R. Rausch, Sept. 6, 1954.

Remarks: ex Hyman. The type locality was incorrectly given by Hyman as “St. Andrew's Island” (Sluys, 1989: 186). This species is the type of the genus *Nesion* Hyman, 1956, by monotypy (Hyman, 1956: 2).

Nexilis epichitonius Holleman and Hand, 1962: 20–22, figs. 1–3.

Holotype: AMNH 503.1–503.5.

Paratypes: AMNH 504.1–504.9 (12 specimens).

Host: *Mopalia hindsi* (Polyplacophora).

Locality: 1 mi south Stinson Beach, Marin Co., California, Pacific Ocean.

Collector: J. Holleman, July 2, 1960.

Remarks: ex Holleman and Hand. Additional paratypes are USNM 46054, 46055. This species is the type of the genus *Nexilis* Holleman and Hand, 1962, by monotypy (Holleman and Hand, 1962: 20).

Notoplana celeris Freeman, 1933: 125–128, figs. 16–18.

Lectotype: AMNH 372.1–372.3.

Locality: Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, 1926–1929.

Remarks: ex Freeman Collection via Hyman. The lectotype was designated by Hyman (1953a: 330). This species was placed in *Notocomplana* Faubel, 1983 (Faubel, 1983: 113–114). See remarks under *Notoplana segnis* for justification in making this species *Pucelis celeris* (Freeman, 1933) **n. comb.**

Notoplana inquieta Freeman, 1933: 120–122, figs. 9, 10, 38.

Syntypes: AMNH 367.1–367.5 (2 specimens).

Locality: Parker's Reef, Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, 1926–1929.

Remarks: ex Freeman Collection via Hyman. This species is a synonym of *Stylochoplana chloranota* (Boone, 1929) (Faubel, 1983: 101).

Notoplana inquilina Hyman, 1955a: 1–4, figs. 1–3.

Holotype: AMNH 582.1, 455.1, 455.2.

Paratypes: AMNH 582.2 (2 specimens).

Locality: In association with hermit crabs, 8–16 m, Cowlitz Bay, Waldron Island, Puget Sound, Washington, United States, Pacific Ocean.

Collector: P. Illg, Aug. 12, 1952.

Remarks: ex Hyman. AMNH 455 is the copulatory apparatus of the partial holotype specimen AMNH 582.1, which is on the same slide as AMNH 582.2.

Notoplana natans Freeman, 1933: 123–125, figs. 14, 15.

Lectotype: AMNH 373.

Paralectotypes: AMNH 694.1–694.3 (5 specimens).

Locality: False Bay, Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, July 11, 1929.

Remarks: ex Freeman Collection via Hyman.

The lectotype was designated by Hyman (1953a: 329). The repository of additional paralectotypes was not cited. This species was placed in *Notocomplana* Faubel, 1983 (Faubel, 1983: 113–114). See remarks under *Notoplana segnis* for justification in making this species *Pucelis natans* (Freeman, 1933) **n. comb.**

Notoplana sanguinea Freeman, 1933: 122–123, figs. 11–13.

Neotype: AMNH 371.1–371.4.

Locality: Turn Rock, Puget Sound, Washington, United States, Pacific Ocean.

Collector: F. A. Pitelka, July 4, 1940.

Remarks: ex Hyman. The neotype was designated by Hyman (1953a: 328). This species was placed in *Notocomplana* Faubel, 1983 (Faubel, 1983: 113, 115). See remarks under *Notoplana segnis* for justification in making this species *Pucelis sanguinea* (Freeman, 1933) **n. comb.**

Notoplana sanjuaniana Freeman, 1933: 128–130, figs. 19, 20, 22.

Lectotype: AMNH 374.

Locality: Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, 1926–1929.

Remarks: ex Freeman Collection via Hyman. Hyman (1953a: 332) cited AMNH 374 as the "holotype." Since no holotype was designated by Freeman (1933), Hyman's action constituted a lectotype designation. The repository of the paralectotypes was not cited. This species was placed in *Notocomplana* Faubel, 1983 (Faubel, 1983: 113, 115). See remarks under *Notoplana segnis* for justification in making this species *Pucelis sanjuaniana* (Freeman, 1933) **n. comb.**

Notoplana segnis Freeman, 1933: 130–133, figs. 23–25, 37.

Lectotype: AMNH 378.

Locality: Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, 1926–1929.

Remarks: ex Freeman Collection via Hyman. Hyman (1953a: 337) cited AMNH 378 as the "holotype." Since no holotype was designated by Freeman (1933), Hyman's action constituted a lectotype designation. The repository of paralectotypes was not

cited. This species is a synonym of *Freemaniana litoricola* (Heath and McGregor, 1912) (Prudhoe, 1985: 178). The monotypic genera *Freemaniana* Hyman, 1953, and *Pucelis* Marcus, 1947, were both considered junior synonyms of *Notocomplana* Faubel, 1983, by Faubel (1983: 113–114). This is in conflict with ICZN regulations and, by the rule of priority, both *Notocomplana* and *Freemaniana* must be junior synonyms of *Pucelis* (Prudhoe, 1989: 68). If Faubel (1983: 113–115) was correct in his congeneric placing of species under *Notocomplana*, then this species becomes *Pucelis litoricola* (Heath and McGregor, 1912) **n. comb.**

Oligocladius albus Freeman, 1933: 140–143, fig. 21.

Holotype: AMNH 391.

Locality: On hydroids at Old Station, Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, 1926–1929.

Remarks: ex Freeman Collection via Hyman. AMNH 391 is the holotype by monotypy. This species was placed in *Acerotisa* Strand, 1926, as *A. alba* by Prudhoe (1985: 179), but in *Eurylepta* Ehrenberg, 1831, by Faubel (1984: 220).

Paracatenula erato Sterrer and Rieger, 1974: 86, figs. 2s–u, 9, 10a–d, j, 14b, c.

Holotype: AMNH 775.1, 775.2.

Locality: “Ann McCrary’s mud flat,” Wilmington, New Hanover Co., North Carolina, United States, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Sterrer and Rieger. The repository of the paratype was not cited.

Paracatenula urania Sterrer and Rieger, 1974: 85–86, figs. 2v, w, 11e.

Holotype: AMNH 772.

Paratypes: AMNH 773.1–773.3, 774.1, 774.2 (1 specimen each).

Locality: On tubes of an onuphid polychaete, fine sand flat, Harbor Island Banks, channel side near bridge highway 76, Wrightsville Beach, north of Wilmington, New Hanover Co., North Carolina, United States, Atlantic Ocean.

Collector: C. Jenner, June 1970.

Remarks: ex Sterrer and Rieger. The repository of additional paratypes was not cited.

This species is the type of the genus *Paracatenula* Sterrer and Rieger, 1974, by original designation (Sterrer and Rieger, 1974: 78).

Paramyozonaria bermudensis Rieger, 1977: 215, figs. 4b, 5, 6f–k.

Lectotype: AMNH 905.

Paralectotypes: AMNH 906 (7 specimens), 907 (1 specimen), 908 (8 specimens), 909 (2 specimens).

Locality: In sand free of sea grass, 1–2 m, shallow subtidal, Tobacco Bay, Bermuda, Atlantic Ocean.

Collector: R. M. Rieger, July 1973.

Remarks: ex Rieger. Rieger marked slide AMNH 905 “holotype.” Since no holotype was designated by Rieger (1977), AMNH 905 is here designated lectotype. This specific name is a nomen nudum from Rieger (1971b: 249).

Paramyozonaria simplex Rieger and Tyler, 1974: 169–170, figs. 1a, 2a, 4a, 6d, 7b, c, 8b, 9a, b, d, 11, 12, 14a, b, 15a, b.

Lectotype: AMNH 782.

Paralectotypes: AMNH 783–786 (1 specimen each), 787.1, 787.2 (2 specimens), 788 (1 specimen).

Locality: New River Inlet, North Carolina, United States, Atlantic Ocean.

Collector: R. M. Rieger, July 1971.

Remarks: ex Rieger and Tyler. Rieger marked slide AMNH 782 “holotype.” Since no holotype was designated by Rieger and Tyler (1974), AMNH 782 is here designated lectotype. This specific name is a nomen nudum from Rieger (1971b: 249).

Parathylacorhynchus reidi Dean, 1980: 5–8, figs. 1–5.

Holotype: AMNH 925.

Paratype: AMNH 926 (1 specimen).

Locality: Griffith’s Head Beach, Reid State Park, Georgetown, Maine, United States, Atlantic Ocean.

Collector: H. K. Dean, Jan. 7, 1976 (AMNH 925), Jan. 10, 1976 (AMNH 926).

Remarks: ex Dean. Additional paratypes are NRS 3200 (Dean, 1980: 5). This species is the type of the genus *Parathylacorhynchus* Dean, 1980, by monotypy (Dean, 1980: 5).

Paromalostomum coronum Spurr, 1983: 334–346, figs. 1–11.

Holotype: AMNH 922.

Paratypes: AMNH 923, 924 (1 specimen each).

Locality: Intertidal zone, Griffith's Head Beach, Reid State Park, Georgetown, Maine, United States, Atlantic Ocean.

Collector: S. J. Spurr, April 1978 (AMNH 922); Unknown, Sept. 19, 1976 (AMNH 923); Unknown, Jan. 5, 1977 (AMNH 924).

Remarks: ex Spurr. Additional paratypes are NRS 3083–3085 (Spurr, 1983: 338).

Pericelis hymanae Poulter, 1974: 95–105, figs. 1, 4, 6–10.

Paratypes: AMNH 756 (2 specimens).

Locality: Oahu, Hawaii, United States, Pacific Ocean.

Collector: J. L. Poulter, July 25, 1961.

Remarks: ex Poulter. The holotype, figs. 1, 4, 6, 7 in Poulter (1974), is USNM 45278 and additional paratypes are USNM 45279–45281, CASIZ 024013–024017, and in BPBM and BM(NH) (Poulter, 1974: 94–95).

Phaenocelis mexicana Hyman, 1953a: 299, figs. 34–36.

Holotype: AMNH 362.

Paratypes: AMNH 392.1–392.3, 544.1–544.3 (1 specimen each).

Locality: San Carlos Bay (AMNH 362, 392); Miramar Beach, Guaymas (AMNH 544), Sonora, Mexico, Pacific Ocean.

Collector: E. F. Ricketts, April 4, 1940 (AMNH 362, 392), G. E. and N. MacGinitie, Feb. 9, 1948 (AMNH 544).

Remarks: ex Hyman. An additional paratype is SBMNH 142931. This species was placed in *Notocomplana* Faubel, 1983 (Faubel, 1983: 113–114). See remarks under *Notoplana segnis* for justification in making this species *Pucelis mexicana* (Hyman, 1953) *n. comb.*

Phagocata cavernicola Hyman, 1954c: 563–566, figs. 65a–c.

Paratypes: AMNH 695.1–695.3 (2 specimens).

Locality: Evac Cave, near Hillside, Pennsylvania, United States.

Collector: R. C. Hoffmaster, Jan. 19, 1951.

Remarks: ex Hyman. The holotype is USNM 24610 and the additional paratypes are USNM 24611, 24612 (Hyman, 1954c: 566). This species may be a synonym of *P.*

m. morgani (Stevens and Boring, 1906) (Kenk, 1989: 3).

Phagocata gracilis monopharyngea Hyman, 1945: 475–476, fig. 1.

Holotype: AMNH 325.

Paratypes: AMNH 326, 327 (1 specimen), 696.1, 696.2 (2 specimens).

Locality: Outlet of tile drain, Haskins, Iowa, United States.

Collector: L. Hubricht, April 24, 1942.

Remarks: ex Hyman. The repository of additional paratypes was not cited. This subspecies is a synonym of *P. velata* (Stringer, 1909) (Kenk, 1989: 2).

Phagocata oregonensis Hyman, 1963a: 2, figs. 1, 2.

Holotype: AMNH 514.1–514.4.

Paratypes: AMNH 515 (3 specimens).

Locality: Vicinity of Crystal Springs, Eastmoreland District, Portland, Multnomah Co., Oregon, United States.

Collector: E. Kozloff.

Remarks: ex Hyman.

Phagocata subterranea Hyman, 1937b: 474–475, figs. 14, 15.

Syntypes: AMNH 644 (2 specimens).

Locality: Donaldson's Cave, near Mitchell, Lawrence Co., Indiana, United States.

Collector: A. E. Emerson, Nov. 12, 1934.

Remarks: ex Hyman. The sectioned syntype is USNM 20230. This species is a synonym of *P. gracilis* (Haldeman, 1840) (Kenk, 1989: 3).

Phagocata woodworthi Hyman, 1937a: 305–306, figs. 3, 4.

Syntypes: AMNH 645.1–645.17 (7 specimens).

Locality: Swan's Island, Maine (AMNH 645.1–645.9); New Haven, New Haven Co., Connecticut (AMNH 645.10); near Yonkers, Westchester Co., New York (AMNH 645.11–645.13); Princeton, Mercer Co., New Jersey (AMNH 645.14–645.17), United States.

Collector: Unknown, May 1936 (AMNH 645.11–645.17).

Remarks: ex Hyman. This species was described as a "nov. nom." for *P. gracilis* (Haldeman, 1840) of Girard, 1851, Woodworth, 1891, and Hyman, 1931 (Hyman,

1937a: 305). This was actually a description of a new species and type material is accordingly recognized here.

Plagiostomum album Hyman, 1938b: 24.

Lectotype: AMNH 338.

Locality: Under stones in tidal zone, Mount Desert Island, Hancock Co., Maine, United States, Atlantic Ocean.

Collector: Unknown, Aug. 1937.

Remarks: ex Hyman. Hyman (1944: 4–5) cited AMNH 338 as the “type.” Since no holotype was designated by Hyman (1938b), Hyman’s action (of 1944) constituted a lectotype designation. The repository of the paralectotypes was not cited. The species was figured by Hyman (1944: figs. 9–11).

Planocera burchami Heath and McGregor, 1912: 461–462, figs. 9, 27, 44.

Neotype: AMNH 382.

Locality: 20 m, Monterey Bay, California, United States, Pacific Ocean.

Collector: G. E. and N. MacGinitie, April 12, 1931.

Remarks: ex Hyman. The neotype (fig. 2, herein) was designated by Hyman (1953a: 359), because none of the five syntypes could be located. This species is the type of the genus *Koinostylochus* Faubel, 1983, by original designation (Faubel, 1983: 60). Faubel (1983: 60) stated that this is “the type of the genus and the only species”; he then proceeded to list nine more species in this “monotypic” genus.

Polycelis tibetica Hyman, 1934: 7–10, pl. 1; pl. 2, figs. 1–4.

Syntypes: AMNH 647.1–647.22 (10 specimens).

Locality: Kashmir (AMNH 647.1); L25: stream (7.8–8.6°C), 3570 m, Leh, Kashmir (AMNH 647.2–647.6); K71: stream (7.0°C), 2590 m, west of Sonamarg, Kashmir (AMNH 647.7–647.14); K76: stream (19.0°C), 3080 m, 1 mi west of Dras, Kashmir (AMNH 647.15–647.18); K78: stream (8.0°C), 2819 m, Karbu, between Dras and Kargil, Kashmir (AMNH 647.19–647.22).

Collector: Yale North India Expedition, no date (AMNH 647.1), June 21 (AMNH 647.2–647.6), May 19 (AMNH 647.7–647.14), May 21 (AMNH 647.15–647.18), May 22 (AMNH 647.19–647.22), 1932.

Remarks: ex Hyman. The repository of additional syntypes was not cited. This specific name is preoccupied by *P. tibetica* (Zabusov, 1911), with which this species may be identical (Kenk, 1974: 57).

Probursa veneris Hyman, 1944: 6, figs. 12, 13.

Holotype: AMNH 328.

Paratypes: AMNH 329–332 (1 specimen), 551.1–551.3 (3 specimens), 552.1, 552.2, 553.1, 553.2, 554, 555.1, 555.2, 556–558 (1 specimen each).

Locality: In empty *Venus* [= *Mercenaria*] shells [Gastropoda], Fire Island, Suffolk Co., New York, United States, Atlantic Ocean.

Collector: M. C. Meyer, Sept. 4, 1940.

Remarks: ex Hyman. This species is the type of the genus *Probursa* Hyman, 1944, by monotypy (Hyman, 1944: 6).

Procerodes pacifica Hyman, 1954d: 1–4, figs. 1–3.

Holotype: AMNH 443.1–443.3.

Paratypes: AMNH 444 (40 specimens), 591 (1 specimen), 592 (38 specimens).

Locality: on kelp (*Macrocystis*), base of Sunset Cliffs, near San Diego, San Diego Co., California, United States, Pacific Ocean.

Collector: T. E. Bowman, June 21, 1953.

Remarks: ex Hyman.

Proschizorhynchus nahantensis Doe, 1974: 101–105, figs. 1–8, 13, 15, 17.

Holotype: AMNH 780.

Paratypes: AMNH 781.1, 781.2 (2 specimens).

Locality: Canoe Beach, Marine Science Institute, Nahant, Essex Co., Massachusetts, United States, Atlantic Ocean.

Collector: D. A. Doe, Aug. 12, 1972.

Remarks: ex Doe. Additional paratypes are NRS 2659, 2660 (Doe, 1974: 105).

Proschizorhynchus papillatus Doe, 1974: 105–108, figs. 9–12, 14–16.

Holotype: AMNH 778.

Paratypes: AMNH 779.1, 779.2 (2 specimens).

Locality: Plum Cove Beach, Lanesville, Massachusetts, United States, Atlantic Ocean.

Collector: D. A. Doe, Jan. 28, 1973.

Remarks: ex Doe. Additional paratypes are NRS 2661, 2662 (Doe, 1974: 108).

Prostheceraeus(?) *bellostriatus* Hyman, 1953a: 373–375, figs. 144, 145.

Syntype: AMNH 368.1–368.9 (1 specimen).

Locality: In tank, Kerckhoff Marine Laboratory, Corona del Mar, Orange Co., California, United States.

Collector: G. E. and N. MacGinitie, May 1938.

Remarks: ex Hyman. The repository of the additional syntype was not cited and it could not be located at SBMNH. This species is the type of the genus *Praestheceraeus* Faubel, 1984, by monotypy (Faubel, 1984: 226).

Prostheceraeus zebra Hyman, 1955f: 266–267, figs. 7, 8.

Holotype: AMNH 463.

Paratypes: AMNH 697, 698.1–698.11 (1 specimen each).

Locality: First culvert, Palisadoes Road (AMNH 463), Palisadoes Road or Ocho Rios (AMNH 698), Jamaica; on leaf of green algae, 3 m, St. Marks Light, Wakulla Co., Florida, United States (AMNH 697), Atlantic Ocean.

Collector: L. H. Hyman, Feb. 1955 (AMNH 463); H. Humm, June 23, 1953 (AMNH 697), 1955 (AMNH 698).

Remarks: ex Hyman. AMNH 463 was fig. 7 in Hyman (1955f). The repository of additional paratypes was not cited and they could not be located in SBMNH.

Prosthiostomum (*Lurymare*) *katoi* Poulter, 1976: 323–327, figs. 7–11, 19.

Paratype: AMNH 851 (1 specimen).

Locality: Reef, 157°51'W, 21°17'20"N, Ala Moana Park, Oahu, Hawaii, United States, Pacific Ocean.

Collector: Unknown, June 4, 1962.

Remarks: ex Poulter. The holotype, figs. 7b, 9a, 10, 11 in Poulter (1976), is USNM 52698 and additional paratypes are CASIZ 002350, USNM 52697, 52699, 52700, and in BM(NH) (Poulter, 1976: 323). This species is placed in the genus *Lurymare* Marcus and Marcus, 1968 (Faubel, 1984: 235).

Prosthiostomum latocelis Hyman, 1953a: 382–384, figs. 155–157.

Paratypes: AMNH 699, 700 (1 specimen each).

Locality: Laguna Beach, Orange Co. (AMNH 699); 30–90 m, California Islands (AMNH

700), California, United States, Pacific Ocean.

Collector: M. W. de Laubenfels, Nov. 27, 1939 (AMNH 699); Allan Hancock Foundation, 1940–1941 (AMNH 700).

Remarks: ex Hyman. AMNH 700 contains one whole mount plus a partial specimen. The holotype was in AHF (Hyman, 1953a: 384), now SBMNH 142918. Additional paratypes are SBMNH 142920.

Prosthiostomum (*Prosthiostomum*) *montiporae* Poulter, 1976: 317–322, figs. 1–6, 18, 22.

Paratypes: AMNH 850 (2 specimens).

Host: *Montipora verrucosa* (Anthozoa).

Locality: 157°47'21"W, 21°26'10"N, Kaneohe Bay, Oahu, Hawaii, United States, Pacific Ocean.

Collector: P. L. Jokiel, Oct. 14, 1971.

Remarks: ex Poulter. The holotype, figs. 1b, 2b, c in Poulter (1976), is USNM 52695 and additional paratypes are USNM 52696 and in BM(NH) (Poulter, 1976: 318). This specific name is a nomen nudum from Jokiel and Townsley (1975: 362).

Prosthiostomum multiceles Hyman, 1953a: 384–386, figs. 158, 159.

Paratype: AMNH 701.1–701.3 (1 specimen).

Locality: Santa Catalina Island, Orange Co., California, United States, Pacific Ocean.

Collector: Allan Hancock Foundation, Aug. 4, 1941.

Remarks: ex Hyman. The holotype was in AHF (Hyman, 1953a: 386), now SBMNH 142919. The repository of the additional paratype was not cited and it could not be located in SBMNH.

Pseudoceros bajae Hyman, 1953a: 365, figs. 135, 136.

Holotype: AMNH 385.

Paratype: AMNH 407 (1 specimen).

Locality: Puerto Peñasco, Gulf of California, Sonora, Mexico, Pacific Ocean.

Collector: G. E. and N. MacGinitie, Dec. 1947.

Remarks: ex Hyman. The repository of additional paratypes was not cited and they could not be located in SBMNH. This species is the type of the genus *Cryptobiceros* Faubel, 1984, by monotypy (Faubel, 1984: 215). This species is placed in *Pseudobi-*

ceros Faubel, 1984 (Newman and Cannon, 1994: 239–240), making *Cryptobiceros* a synonym of *Pseudobiceros*.

Pseudoceros canadensis Hyman, 1953a: 368, figs. 138, 139.

Holotype: AMNH 388.

Locality: Clayoquot, British Columbia, Canada, Pacific Ocean.

Collector: E. F. Ricketts, June 7, 1945.

Remarks: ex Hyman. The repository of the paratype was not cited and it could not be located in SBMNH.

Pseudoceros corallophilus Hyman, 1954a: 223–224, fig. 2.

Holotype: AMNH 416.

Locality: Heron Island, Great Barrier Reef, Queensland, Australia, Pacific Ocean.

Collector: J. M. Schweitzer, summer 1952.

Remarks: ex Hyman. AMNH 416 is the holotype by monotypy. This species is a synonym of *P. bimarginatus* Meixner, 1907 (Newman and Cannon, 1994: 217–218).

Pseudoceros griseus Hyman, 1959d: 10–11, figs. 13, 14.

Holotype: AMNH 490.

Locality: 1 ft, sandy bottom, Newport Harbor, California, United States, Pacific Ocean.

Collector: W. E. Pequegnat, Sept. 1, 1958.

Remarks: ex Hyman. AMNH 490 is the holotype by monotypy.

Pseudoceros mexicanus Hyman, 1953a: 363–365, figs. 132–134.

Holotype: AMNH 384.

Paratypes: AMNH 406, 599.1–599.10 (1 specimen each).

Locality: Puerto Peñasco, Gulf of California, Sonora, Mexico, Pacific Ocean.

Collector: G. E. and N. MacGinitie, Dec. 1947.

Remarks: ex Hyman. The repository of additional paratypes was not cited and they could not be located in SBMNH.

Pseudoceros texanus Hyman, 1955f: 264–266, figs. 5, 6.

Holotype: AMNH 464.

Paratypes: AMNH 600.1–600.6, 601, 602.1–602.14 (1 specimen each).

Locality: Port Aransas, Aransas Co., Texas, United States, Atlantic Ocean.

Collector: W. C. Guest, July–Aug. 1954.

Remarks: ex Hyman. AMNH 601 was the figured specimen in Hyman (1955f).

Pseudostylochus ostreophagus Hyman, 1955a: 4–7, figs. 4–7.

Holotype: AMNH 452.

Paratypes: AMNH 453.1–453.8 (7 specimens), 454.1–454.7 (4 specimens).

Locality: Miyagi Prefecture, Japan (AMNH 452, 453); Puget Sound, Washington, United States (AMNH 454), Pacific Ocean.

Collector: C. E. Woelke, April 1, 1954 (AMNH 452, 453) and Nov. 1953 (AMNH 454).

Remarks: ex Hyman. This species is placed in *Koinostylochus* Faubel, 1983 (Faubel, 1983: 60).

Retronectes atypica Doe and Rieger, 1977: 1–7, figs. 1, 2.

Holotype: AMNH 915.

Paratypes: AMNH 916, 917 (1 specimen each).

Locality: Exposed beach, 500 m from inlet mouth, 20–50 cm sediment depth, clean medium sand, Bogue Bank, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: D. A. Doe, April 1977.

Remarks: ex Doe and Rieger. The repository of additional paratypes was not cited.

Retronectes clio Sterrer and Rieger, 1974: 80–81, figs. 2a–c, 10i, 11c, 13f.

Holotype: AMNH 771.

Locality: 1.5 m, fine sand, Klubban Beach, Kristineberg, Sweden, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Sterrer and Rieger. The repository of the paratypes was not cited.

Retronectes thalia Sterrer and Rieger, 1974: 78–79, figs. 2h–j, 3a–c, 4a, b, 5–8, 10e–h, 11a, b, d.

Holotype: AMNH 769.

Paratype: AMNH 770.1, 770.2 (1 specimen).

Locality: Station 14208, 28 m, 34°12.5'N, 76°14'W, off Beaufort, Carteret Co., North Carolina, United States, Atlantic Ocean.

Collector: R/V "Eastward" cruise E-4-70, May 1970.

Remarks: ex Sterrer and Rieger. The repository of additional paratypes was not cited. This species is the type of the genus *Retro-*

- nectes* Sterrer and Rieger, 1974, by original designation (Sterrer and Rieger, 1974: 78).
- Rhynchodemus americanus* Hyman, 1943: 7–9, figs. 9–12.
 Syntypes: AMNH 296.1–296.8 (5 specimens).
 Locality: Greenhouse, Forest Park, St. Louis, St. Louis Co., Missouri, United States.
 Collector: L. Hubricht, Jan. 18, 1936.
 Remarks: ex Hyman. The repository of additional syntypes was not cited. This species is a synonym of *R. sylvaticus* (Leidy, 1851) (Ogren, 1989: 44).
- Rhynchodemus oahuensis* Hyman, 1939b: 120–122, figs. 6–8.
 Holotype: AMNH 634.
 Paratypes: AMNH 635, 636.1, 636.2 (1 specimen each), 637.1–637.5 (3 specimens).
 Locality: Wet forests under stones and fallen leaves, 2400 ft, Kipapa Ridge; 1800 ft, Koolan Mountains, Oahu, Hawaii, United States.
 Collector: F. X. Williams, 1938.
 Remarks: ex Hyman.
- Solenofilomorpha funilis* Crezée, 1975: 775–783, figs. 1–4, 21c.
 Holotype: AMNH 815.
 Paratypes: AMNH 816–820 (1 specimen each).
 Locality: Estuary, White Oak River, opposite Swansboro, Onslow Co. (AMNH 815–818); by bridge, highway 76, Wrightsville Beach, New Hanover Co. (AMNH 819, 820), North Carolina, United States, Atlantic Ocean.
 Collector: Unknown.
 Remarks: ex Crezée. The species name was misspelled by Crezée (1975) as "*funilia*" in fig. 21c. The repository of additional paratypes was not cited.
- Solenofilomorpha guaymensis* Crezée, 1975: 784–787, figs. 5, 6.
 Holotype: AMNH 821.
 Paratypes: AMNH 822–824 (1 specimen each).
 Locality: 1 m, clean, medium-fine quartz sand, southwestern shore, Bahia de Baco-chibampo, near Guaymas, Sonora, Mexico, Pacific Ocean.
 Collector: Unknown, Aug. 1971.
- Remarks: ex Crezée. The repository of additional paratypes was not cited.
- Speophila buchanani* Hyman, 1937b: 468–469, figs. 16–19.
 Syntypes: AMNH 648.1–648.12 (6 specimens).
 Locality: Hovey's Cathedral Domes, Fox's Alley and Becky's Alley, underside of stones, stream (55°F, pH 7.6), 300 ft down, 4th level, Mammoth Cave, Kentucky, United States.
 Collector: J. W. Buchanan, Oct.–Nov. 1935.
 Remarks: ex Hyman. This species is placed in *Sphalloplana* Beauchamp, 1931, and the subgenus *Speophila* Hyman, 1937 (Kenk, 1989: 5).
- Speophila hoffmasteri* Hyman, 1954c: 570–573, figs. 66a, 67b, 68.
 Paratype: AMNH 702.1–702.6 (1 specimen).
 Locality: Blowing Cave, Pendleton Co., West Virginia, United States.
 Collector: R. E. Hoffmaster.
 Remarks: ex Hyman. The holotype is USNM 24616 and the additional paratype is USNM 24617 (Hyman, 1954c: 573). This species is placed in *Sphalloplana* Beauchamp, 1931, and the subgenus *Speophila* Hyman, 1937 (Kenk, 1989: 5).
- Speophila hubrichti* Hyman, 1945: 479–480, figs. 7, 8.
 Holotype: AMNH 320.
 Paratypes: AMNH 321–324 (1 specimen), 703.1, 703.2 (2 specimens), 704.1–704.16 (3 specimens).
 Locality: Kohn's Cave, Ste. Genevieve Co., Missouri (AMNH 320–324, 703); Morrison's Cave, near Burksville, Monroe Co., Illinois (AMNH 704), United States.
 Collector: L. Hubricht, Sept. 13, 1941 (AMNH 320–324, 703), June 1, 1936 (AMNH 704).
 Remarks: ex Hyman. Additional paratypes are USNM 20640. This species is placed in *Sphalloplana* Beauchamp, 1931, and the subgenus *Speophila* Hyman, 1937 (Kenk, 1989: 5).
- Speophila pricei* Hyman, 1937b: 462–466, figs. 7–11, 20, 21.
 Paratypes: AMNH 649.1–649.24 (4 specimens).
 Locality: Refton Cave, John Hess Farm, 1 mi

- northwest Refton, along south bank of Pequea Creek, Lancaster, Lancaster Co. (AMNH 649.1–649.20); Brownstone Cave, 1 mi east Hummelstown, Dauphin Co. (AMNH 649.21–649.24), Pennsylvania, United States.
Collector: J. W. Price, Jan. 20 (AMNH 649.1), April 17 (AMNH 649.2–649.20), May 23 (AMNH 649.21–649.24), 1936.
Remarks: ex Hyman. The holotype, fig. 8 in Hyman (1937b), is USNM 20228 and an additional paratype is USNM 20236. AMNH 649.25–649.31, previously considered types (see Feinberg, 1970: 49–50), are here excluded from the type series because they were not cited by Hyman (1937b) and there is no indication that they were collected prior to the species description. This species is the type of the genus *Speophila* Hyman, 1937, by original designation (Hyman, 1937b: 473). This species is placed in *Sphalloplana* Beauchamp, 1931, and the subgenus *Speophila* Hyman, 1937 (Kenk, 1989: 5).
- Sphalloplana georgiana* Hyman, 1954c: 566–570, figs. 65d, 66b, 67a.
Paratypes: AMNH 705.1–705.16 (4 specimens).
Locality: Waterfall Cave, Trenton, Walker Co., Georgia, United States.
Collector: C. E. Mohr, Dec. 8, 1950.
Remarks: ex Hyman. The holotype is USNM 24614 and the anterior part of a paratype is USNM 24615 (Hyman, 1954c: 570). One specimen from AMNH 705 is the remainder of the partial paratype in USNM. This species is placed in the subgenus *Sphalloplana* Beauchamp, 1931 (Kenk, 1989: 4).
- Sphalloplana kansensis* Hyman, 1945: 478–479, fig. 6.
Holotype: AMNH 311–313.
Locality: Purity Springs, near Augusta, Butler Co., Kansas, United States.
Collector: L. Hubricht, May 18, 1942.
Remarks: ex Hyman. AMNH 311–313 is the holotype by monotypy. This species is placed in the subgenus *Sphalloplana* Beauchamp, 1931 (Kenk, 1989: 4).
- Sphalloplana mohri* Hyman, 1939e: 276–278, figs. 2–5.
Holotype: AMNH 650.
Paratypes: AMNH 651.1–651.24 (2 specimens).
Locality: Ezell's Cave, San Marcos, Hays Co., Texas, United States.
Collector: C. E. Mohr and K. Dearolf, June 19 (AMNH 651), 21 (AMNH 650), 1938.
Remarks: ex Hyman. This species is the type of the subgenus *Polypharyngea* Carpenter, 1971, by monotypy (Carpenter, 1971: 1284; Kenk, 1989: 5).
- Sphalloplana virginiana* Hyman, 1945: 477–478, figs. 4, 5.
Holotype: AMNH 314.
Paratypes: AMNH 315–319 (1 specimen), 706 (3 specimens).
Locality: Showalter's Cave, near Lexington, Rockbridge Co., Virginia, United States.
Collector: J. A. Fowler, Oct. 30, 1943.
Remarks: ex Hyman. Additional paratypes are USNM 53407–53415. This species is placed in the subgenus *Speophila* Hyman, 1937 (Kenk, 1989: 4).
- Stylochoplana inquilina* Hyman, 1950: 55–56, fig. 1.
Paratypes: AMNH 707.1–707.4 (2 specimens).
Locality: In association with hermit crab/anemone complex, 16 fathoms, Oahu, Hawaii, United States, Pacific Ocean.
Collector: Unknown.
Remarks: ex Hyman. The holotype is BPBM 268 (Hyman, 1950: 58).
- Stylochoplana longastyletta* Freeman, 1933: 119, figs. 6–8.
Neotype: AMNH 370.
Locality: Turn Rock, Puget Sound, Washington, United States, Pacific Ocean.
Collector: F. A. Pitelka.
Remarks: ex Hyman. The neotype (fig. 3, herein) was designated by Hyman (1953a: 327). This species is placed in *Notoplana* Laidlaw, 1903 (Hyman, 1953a: 325).
- Stylochoplana longipenis* Hyman, 1953a: 305–307, figs. 45–48.
Holotype: AMNH 365.1–365.3.
Paratype: AMNH 708 (1 specimen).
Locality: On rocky shore, Miramar Beach, Guaymas, Sonora, Mexico (AMNH 365); Point Fermin, San Pedro, Los Angeles Co., California, United States (AMNH 708), Pacific Ocean.

Collector: G. E. and N. MacGinitie, Feb. 9, 1948 (AMNH 365); D. Reish, Dec. 1950–Feb. 1951 (AMNH 708).

Remarks: ex Hyman. An additional paratype is SBMNH 142924. This species is placed in *Phaenoplana* Faubel, 1983 (Faubel, 1983: 104).

Stylochoplana reishi Hyman, 1959d: 5–7, figs. 8–10.

Holotype: AMNH 492.

Paratype: AMNH 607.1–607.3 (1 specimen).

Locality: Los Angeles Harbor, Newport Station, Los Angeles Co., California, United States, Pacific Ocean.

Collector: D. Reish, Jan. 1954.

Remarks: ex Hyman. AMNH 492 is fig. 8 in Hyman (1959d). The repository of additional paratypes was not cited and they could not be located in SBMNH. This species is placed in *Armatoplana* Faubel, 1983 (Faubel, 1983: 102–103).

Stylochoplana viridis Freeman, 1933: 118, figs. 3–5.

Lectotype: AMNH 376.1–376.4.

Locality: False Bay or near Brown's Island, Puget Sound, Washington, United States, Pacific Ocean.

Collector: D. Freeman, 1926–1929.

Remarks: ex Freeman Collection via Hyman. Hyman (1953a: 334) cited AMNH 376 as the "holotype." Since no holotype was designated by Freeman (1933), Hyman's action constituted a lectotype designation. The repository of the paralectotype was not cited. This species is the type of the genus *Triplana* Faubel, 1983, by monotypy (Faubel, 1983: 104).

Stylochus californicus Hyman, 1953a: 285, figs. 18, 19.

Holotype: AMNH 358.

Paratypes: AMNH 408.1–408.3 (3 specimens), 533 (1 specimen).

Locality: Laguna Beach or San Diego (AMNH 358); burrows of *Barnea pacifica* [Bivalvia], Anaheim Slough, 20 mi north Corona del Mar, Orange Co. (AMNH 408, 533), California, United States, Pacific Ocean.

Collector: G. E. and N. MacGinitie, July 8, 1934 (AMNH 408, 533).

Remarks: ex Hyman. The repository of additional paratypes was not cited and they

could not be located in SBMNH. This species is placed in the subgenus *Stylochus* Ehrenberg, 1831 (Prudhoe, 1985: 202).

Stylochus franciscanus Hyman, 1953a: 280–282, figs. 8, 9.

Holotype: AMNH 356.

Paratypes: AMNH 534, 535.1–535.3 (1 specimen each), 536.1–536.4 (4 specimens).

Locality: Corona del Mar, Orange Co. (AMNH 356); San Pedro, Los Angeles Co. (AMNH 534); San Francisco Bay (AMNH 535, 536), California, United States, Pacific Ocean.

Collector: G. E. and N. MacGinitie, Nov. 27, 1937 (AMNH 356), D. Freeman (AMNH 534), "Albatross" (AMNH 535, 536).

Remarks: ex Hyman. The repository of additional paratypes was not cited and they could not be located in SBMNH. This species is placed in the subgenus *Stylochus* Ehrenberg, 1831 (Prudhoe, 1985: 203).

Stylochus tripartitus Hyman, 1953a: 282–283, figs. 10–13.

Holotype: AMNH 357.1–357.3.

Paratypes: AMNH 409.1–409.6 (1 specimen), 538 (1 specimen [part]).

Locality: El Mogote, Baja California, Mexico (AMNH 409); Newport Bay (AMNH 538); Pacific Grove, Monterey Co. (AMNH 357), California, United States, Pacific Ocean.

Collector: E. F. Ricketts, March 22, 1940 (AMNH 409); L. H. Hyman, Aug. 1936 (AMNH 357); G. E. and N. MacGinitie, Sept. 1932 (AMNH 538).

Remarks: ex Hyman. The repository of the additional paratype was not cited and it could not be located in SBMNH. This species is placed in the subgenus *Imogene* Girard, 1853 (Prudhoe, 1985: 206).

Syndesmis glandulosa Hyman, 1960: 9–12, figs. 8–12.

Syntypes: AMNH 498.1, 498.2 (11 specimens), 608.1–608.4 (15 specimens).

Host: *Diadema setosum* (Echinoidea).

Locality: Nossi-Bé, Madagascar, Indian Ocean.

Collector: A. G. Humes, June 1955.

Remarks: ex Hyman. This species is placed in *Syndisyrinx* Lehman, 1946 (Cannon, 1987: 302).

Thysanozoon californicum Hyman, 1953a: 362–363, figs. 130, 131.

Holotype: AMNH 383.

Locality: Corona del Mar, Orange Co., California, United States, Pacific Ocean.

Collector: G. E. and N. MacGinitie, May 12, 1933.

Remarks: ex Hyman. AMNH 383 is the holotype by monotypy.

Thysanozoon flavotuberculatum Hyman, 1939g: 16–17, fig. 20.

Paratype: AMNH 709 (1 specimen).

Locality: On rooted seaweeds, the Reach, Bermuda, Atlantic Ocean.

Collector: L. H. Hyman, Aug. 1935.

Remarks: ex Hyman. The holotype was BOC 1059 (Hyman, 1939g: 17), now YPM 20152.

Zygantropiana styliifera Hyman, 1953a: 308–310, figs. 52–54.

Holotype: AMNH 366.

Locality: Rocky littoral, Puerto Refugio, Angel de la Guardia Island or Tiberon Island, Gulf of California, Mexico, Pacific Ocean.

Collector: E. F. Ricketts, April 2–3, 1940.

Remarks: ex Hyman. The repository of the paratype was not cited and it could not be located in SBMNH. This species is placed in *Alloioiplana* Plehn, 1906 (Faubel, 1983: 101).

CLASS MONOGENEA

Caballerocotyla klawei Stunkard, 1962: 883–884, figs. 1–7.

Paratypes: AMNH 860 (1 specimen), 939.1–939.30 (3 specimens).

Host: *Neothunnus macropterus* (Osteichthyes).

Locality: 18°24'N, 104°38'W, Mexico, Pacific Ocean.

Collector: W. L. Klawe, March 29, 1957.

Remarks: ex Stunkard. The holotype is USNPC 59865 (Stunkard, 1962: 889). This species is the type of the genus *Nasicola* Yamaguti, 1968, by monotypy (Yamaguti, 1968: 53–55).

Gyrodactylus fairporti Van Cleave, 1921: 36–37, figs. 1, 2a, 2c.

Syntype: AMNH 1433 (1 specimen).

Host: *Ameiurus melas* or *Cyprinus carpio* (Osteichthyes).

Locality: Fairport, Iowa, United States.

Collector: H. J. Van Cleave, Aug. 20, 1920.

Remarks: ex Van Cleave via Stunkard. Additional syntypes are USNPC 37522 (Salley et al., 1978: 48) and were in the collection of the author (Van Cleave, 1921: 37).

Octobothrium minus Olsson, 1876: 10.

Syntype: AMNH 13 (1 specimen).

Host: *Gadus melanostomus* (Osteichthyes).

Locality: Bergen, Norway.

Collector: P. Olsson, Aug.–Sept. 18??.

Remarks: ex Olsson Collection. This species is a synonym of *Declidophora minor* (Olsson, 1868) (Euzet and Trilles, 1961: 293, fig. 1).

Oculotrema hippopotami Stunkard, 1924b: 436–440, pl. 16.

Paratypes: AMNH 142, 1470.1–1470.4 (1 specimen each).

Host: *Hippopotamus* sp. (Mammalia).

Locality: Giza Zoological Gardens, Cairo, Egypt.

Collector: A. Looss.

Remarks: ex Stunkard. The holotype is in MIRP (Stunkard, 1924b: 440). This species is the type of the genus *Oculotrema* Stunkard, 1924, by monotypy (Stunkard, 1924b: 440).

Polystoma floridanum Stunkard, 1924a: 100–103, figs. 7, 8.

Lectotype: AMNH 1464.

Paralectotype: AMNH 1465 (1 specimen).

Host: *Pseudemys floridana* (Testudinata).

Locality: Lakeland, Polk Co., Florida, United States.

Collector: Unknown.

Remarks: ex Stunkard. Stunkard marked slide AMNH 1464 “type” and AMNH 1465 “co-type.” Since no types were designated by Stunkard (1924a), AMNH 1464 is here designated lectotype. AMNH 1464 was figured by Stunkard (1924a: fig. 7) and is fig. 4, herein. An additional paralectotype is USNPC 37519 (Salley et al., 1978: 50). This species is a synonym of *Neopolystoma orbiculare* (Stunkard, 1916) (Price, 1939: 87).

Polystoma integerrimum nearcticum Paul, 1938: 495–496, figs. 1, 2.

Holotype: AMNH 281.

Paratype: AMNH 282 (1 specimen).

Host: *Hyla versicolor* and/or *H. cinerea* (Amphibia).

Locality: Weston, Fairfield Co., Connecticut, United States.

Collector: A. A. Paul.

Remarks: ex Paul. AMNH 281 was fig. 1 and AMNH 282 was fig. 2 in Paul (1938). An additional paratype is USNPC 41164 (Salley et al., 1978: 94). This subspecific name is a nomen nudum from Paul (1935) (Price, 1939: 82). This subspecies has been elevated to full species rank (Price, 1939: 82).

Polystoma multifalx Stunkard, 1923c: 357.

Lectotype: AMNH 883.1.

Paralectotypes: AMNH 882 (1 specimen), 883.2–883.5 (3 specimens), 1466.1–1466.10 (13 specimens).

Host: *Pseudemys floridana* (Testudinata).

Locality: Lakeland, Polk Co., Florida, United States.

Collector: C. R. Halter, 1923.

Remarks: ex Stunkard. Stunkard marked slide AMNH 882 "type" and AMNH 883.1 "co-type." Since no types were designated by Stunkard (1923c), a lectotype designation is desirable. Unfortunately, the "type" specimen is badly broken so the "co-type," AMNH 883.1 (fig. 5, herein), is here designated lectotype. The species was figured by Stunkard (1924a: figs. 1–6). The repository of additional paralectotypes was not cited. This species is placed in *Polystomoides* Ward, 1917 (Price, 1939: 85–86).

Polystoma opacum Stunkard, 1916: 23.

Paratype: AMNH 1469.1–1469.4 (1 specimen).

Host: *Malacoclemmys lesueurii* (Testudinata).

Locality: Newton, Newton Co., Texas, United States.

Collector: Unknown, Jan. 25, 1914.

Remarks: ex Stunkard. The holotype was in HCUI (Stunkard, 1916: 26), now USNPC 51801 (not 37520 as per Salley et al., 1978: 98). The repository of the additional paratype was not cited. The species was figured by Stunkard (1917: figs. 15–21, 37, 38, 40). This species is a synonym of *Polystomoides coronatum* (Leidy, 1888) (Price, 1939: 84–85).

Polystoma orbiculare Stunkard, 1916: 23.

Paratypes: AMNH 1434.1–1434.3 (3 speci-

mens), 1467, 1468.1, 1468.2 (1 specimen each).

Host: *Pseudemys scripta* (AMNH 1434.1, 1434.3, 1468) and *Chrysemys marginata* (AMNH 1434.2, 1467) (Testudinata).

Locality: Raleigh, Wake Co., North Carolina (*P. scripta*) and Creston, Union Co., Iowa (*C. marginata*), United States.

Collector: Unknown, 1895 (*C. marginata*) and May 20, 1915 (*P. scripta*).

Remarks: ex Stunkard. The holotype was in HCUI (Stunkard, 1916: 26), now USNPC 52173 (Salley et al., 1978: 100). The repository of additional paratypes was not cited. The species was figured by Stunkard (1917: figs. 1–14, 34–36, 43, 44). This species is the type of the genus *Neopolystoma* Price, 1939, by original designation (Price, 1939: 87–88).

Polystomoides oris Paul, 1938: 501, fig. 3.

Holotype: AMNH 280.

Host: *Chrysemys picta* (Testudinata).

Locality: Cold Spring, Suffolk Co., New York, United States.

Collector: A. A. Paul, Jan. 11, 1936.

Remarks: ex Paul. A paratype is USNPC 41165 (Salley et al., 1978: 100).

Sphyranura oligorchis Alvey, 1933: 140.

Syntypes: AMNH 1432.1–1432.5 (5 specimens).

Host: *Necturus maculosus* (Amphibia).

Locality: Pennsylvania, United States.

Collector: Unknown.

Remarks: ex Alvey via Stunkard. The species was figured by Alvey (1936: pl. 10, fig. 1).

Sphyranura polyorchis Alvey, 1936: 246–247, fig. 2.

Holotype: AMNH 1431.

Host: *Necturus maculosus* (Amphibia).

Locality: Meadville, Crawford Co., Pennsylvania, United States.

Collector: Unknown.

Remarks: ex Alvey via Stunkard. AMNH 1431 was the figured specimen in Alvey (1936). This species may be a synonym of *S. osleri* Wright, 1879 (McAllister et al., 1991: 139), but *S. polyorchis* type material was not studied by McAllister et al., as its repository was not previously known to the scientific community.

CLASS TREMATODA

Acanthochasmus diploporus Stunkard, 1931: 722–723, figs. 1, 2.

Lectotype: AMNH 875.

Paralectotypes: AMNH 875A, 876 (1 specimen each), 877 (2 specimens), 878–880 (1 specimen each), 881.1–881.4 (4 specimens).

Host: *Alligator mississippiensis* (Crocodylia).

Locality: United States.

Collector: Unknown, 1925.

Remarks: ex Stunkard. Stunkard marked slide AMNH 875 “holotype.” Since no holotype was designated by Stunkard (1931), AMNH 875 is here designated lectotype (fig. 6, herein). AMNH 875 and 875A are on the same slide, but are clearly distinguished. The repository of additional paralectotypes was not cited. This species is the type of the genus *Proctoecaecum* Baugh, 1956, by monotypy (Baugh, 1956: 305), but is a synonym of *P. coronarium* (Cobbold, 1861) (Brooks, 1980: 351).

Acanthostomum minimum Stunkard, 1938a: 39–40, fig. 2.

Holotype: AMNH 268.

Paratype: AMNH 852 (1 specimen).

Host: *Rhamdia guatemalensis* (Osteichthyes).

Locality: Kaua Cave (AMNH 268); San Isidro Cave (AMNH 852), Yucatan, Mexico.

Collector: A. S. Pearse, June 16 (AMNH 268) and July 3 (AMNH 852), 1936.

Remarks: ex Stunkard. AMNH 268 was the figured specimen in Stunkard (1938a). This species is the type of the genus *Stunkardiella* Lamothe and Ponciano, 1986 (Lamothe and Ponciano, 1986: 309) but was placed in the genus *Acanthostomum* Looss, 1899, and the subgenus *Blairium* Brooks and Holcman, 1993, by Brooks and Holcman (1993: 214), who did not cite Lamothe and Ponciano (1986).

Allassostoma magnum Stunkard, 1916: 24–25.

Paratype: AMNH 983 (1 specimen).

Host: *Pseudemys troostii* (Testudinata).

Locality: Havana, Mason Co., Illinois, United States.

Collector: H. W. Stunkard.

Remarks: ex Stunkard. The holotype was in HCUI (Stunkard, 1916: 26), now USNPC

49978 (vial specimen). See remarks under *A. parvum* for a discussion of alcohol-preserved holotypes designated by Stunkard. An additional probable paratype is USNPC 51358 (Pilitt, personal commun.). The species was figured by Stunkard (1917: figs. 59–65). This species is the type of the genus *Allassostoma* Stunkard, 1916, by original designation (Stunkard, 1916: 24). *Allassostoma* (Stunkard, 1924a: 103) is an unjustified emendation; the original spelling *Allassostoma* should be retained (e.g., Sey, 1991: 203).

Allassostoma parvum Stunkard, 1916: 24–25.

Paratype: AMNH 792 (1 specimen).

Host: *Chelydra serpentina* (Testudinata).

Locality: Urbana, Champaign Co., Illinois, United States.

Collector: T. B. Magath.

Remarks: ex Stunkard. The holotype was in HCUI (Stunkard, 1916: 26), now USNPC 49979 (vial specimen). The additional paratype is USNPC 51357. USNPC 49980 (Salley et al., 1978: 106) is not a type specimen as it was collected after description. Although Stunkard rarely designated alcohol specimens as holotypes (but see *Stephanopharynx secundus*), the evidence indicates that he did so on this occasion as USNPC 51357 is sectioned and he did not designate a sectioned specimen as holotype unless he had only a single specimen. The species was figured by Stunkard (1917: figs. 66–71). This species is the type of the subgenus *Allassostomoides* Stunkard, 1924, by monotypy (Stunkard, 1924a: 103), now elevated to generic rank as *Allassostomoides parvus* (Brooks and Mayes, 1975: 404). Sey (1991: 204) was incorrect in using the spelling *Allassostomoides*; the original spelling should be retained.

Amphistomum bicaudatum Poirier, 1909: 581–582, figs. 4, 5.

Lectotype: AMNH 526 (1 specimen).

Paralectotype: AMNH 527.1–527.5 (1 specimen).

Host: *Loxodon africanus* (Mammalia).

Locality: Fort Archimbault, Chad.

Collector: M. Kerandel.

Remarks: ex Roubaud via Stunkard (see Stunkard, 1929a: 260). AMNH 526, fig. 27 in Stunkard (1929a), is here designated lec-

totype (fig. 7, herein). This species is placed in *Brumptia* Travassos, 1921, as *B. bicaudata* (Stunkard, 1929a: 258).

Anenterotrema auritum Stunkard, 1938a: 42–45, fig. 4.

Holotype: AMNH 267.

Paratypes: AMNH 862–866 (1 specimen each), 1023.1–1023.10 (10 specimens), 1025.1–1025.6 (6 specimens).

Host: *Micronycteris megalotis mexicana* (Mammalia).

Locality: Xmahit Cave, Tekax (AMNH 267, 862–866, 1023); Xconsacab, Tizamin (AMNH 1025), Yucatan, Mexico.

Collector: A. S. Pearse, July 31 (AMNH 267, 862–866, 1023) and Aug. 11 (AMNH 1025), 1936.

Remarks: ex Stunkard. AMNH 267 was the figured specimen in Stunkard (1938a). An additional paratype is USNPC 71223, erroneously cited as "holotype" by Salley et al. (1978: 17). This species is the type of the genus *Anenterotrema* Stunkard, 1938, by original designation (Stunkard, 1938a: 42).

Anenterotrema singulare Stunkard, 1938a: 45–46, fig. 5.

Holotype: AMNH 271.

Host: *Natalus mexicanus* (Mammalia).

Locality: Balaam Canche Cave, Chichen Itza, Yucatan, Mexico.

Collector: A. S. Pearse, June 22, 1936.

Remarks: ex Stunkard. AMNH 271 is the holotype by monotypy.

Ascocotyle (Parascocotyle) diminuta Stunkard and Haviland, 1924: 4–5, fig. 1.

Lectotype: AMNH 137.

Paralectotypes: AMNH 139, 140 (1 specimen each).

Host: *Rattus norvegicus* (Mammalia).

Locality: Clason Point Dump, New York City, New York, United States.

Collector: H. W. Stunkard, April 2, 1924.

Remarks: ex Stunkard. Stunkard marked slide AMNH 137 "type." Since no holotype was designated by Stunkard and Haviland (1924), AMNH 137 is here designated lectotype (fig. 8, herein). The repository of additional paralectotypes was not cited. These specimens are poor, moribund representatives of the species, but are the only extant types. It is possible that the species

discussed by Stunkard and Uzmann (1955) as *Phagicola diminuta* is in fact a different species from the *A. (P.) diminuta* of Stunkard and Haviland (1924) (Overstreet, personal commun.). If this is true, then the Stunkard and Uzmann (1955) species will need to be described, if not synonymous with any other. This species is placed in the genus *Phagicola* Faust, 1920, by Price (1932: 89), but *Phagicola* is considered a subgenus of *Ascocotyle* by de Núñez (1993: 197).

Asymphyllodora amnicolae Stunkard, 1959: 571–574, figs. 1–7.

Syntypes: AMNH 1547.1–1547.14 (20+ specimens), 1548.1–1548.15 (71+ specimens), 1549.1–1549.3 (7+ specimens), 1550.1–1550.22 (38+ specimens).

Host: *Amnicola limosa* (Gastropoda).

Locality: Woods Hole (AMNH 1547, 1549, 1550); Wing Pond, Falmouth (AMNH 1548), Barnstable Co., Massachusetts, United States, Atlantic Ocean.

Collector: Woods Hole Marine Laboratory, 1954 (AMNH 1547), Sept. 1955 (AMNH 1548), 1957 (AMNH 1549), and July 1959 (AMNH 1550).

Remarks: ex Stunkard. The repository of additional syntypes was not cited. In Stunkard (1959), one of the specimens from AMNH 1549.1 was fig. 5; one of the specimens from AMNH 1550.2 was likely fig. 3; one of the specimens from AMNH 1550.17 was likely fig. 2; AMNH 1550.18 was likely fig. 1; and the adult on AMNH 1550.19 was likely fig. 7.

Botulus cablei Stunkard, 1965a: 488–491, figs. 1–4.

Paratypes: AMNH 652.1–652.95 (5 specimens).

Host: *Alepisaurus borealis* (Osteichthyes).

Locality: 26°04.9'S, 101°47'W, Easter Island, Chile, Pacific Ocean.

Collector: W. L. Klawe, "Shoyo Maru," Jan. 23, 1963.

Remarks: ex Stunkard. The holotype whole mount specimen, fig. 1 in Stunkard (1965a), and five vial paratypes are USNPC 60934 (Stunkard, 1965a: 491). This species is a synonym of *B. microporus* (Monticelli, 1889) (Gibson and Bray, 1977: 208–209).

Cephalogonimus amphiumae Chandler, 1923: 1–3, fig. 1.

Paratype: AMNH 653 (1 specimen).

Host: *Amphiuma means* (Amphibia).

Locality: Louisiana, United States.

Collector: Unknown, May 1922.

Remarks: ex Chandler via Stunkard. The holotype is USNPC 25171 (Chandler, 1923: 2).

Cephalogonimus compactus Stunkard, 1924a: 106–109, fig. 10.

Holotype: AMNH 654.

Host: *Pseudemys floridana* (Testudinata).

Locality: Florida, United States.

Collector: C. R. Halter.

Remarks: ex Stunkard. The description of this species was based solely on the sexually mature specimen in fig. 10 (Stunkard, 1924a: 106). AMNH 654 is therefore the holotype by monotypy. AMNH 655 is a whole mount of one of the additional, non-type specimens mentioned in Stunkard (1924a: 106).

Cercaria loossi Stunkard, 1929b: 106.

Syntypes: AMNH 982 (1 specimen), 1351.1–1351.20 (15+ specimens).

Host: *Hydroides dianthus* (Polychaeta).

Locality: Woods Hole, Barnstable Co., Massachusetts, United States, Atlantic Ocean.

Collector: Woods Hole Marine Laboratory, 1929.

Remarks: ex Stunkard. Not “*C. loossi* Sinitzyn, 1905” of Doss (1968: 316), a combination not found anywhere in Sinitzyn (1906) (not 1905 as given by Doss; the spelling of the author’s name depends on the translation). Sinitzyn never referred to anything other than the cercaria of *Gorgodera loossi* Sinitzyn, 1906, which is a synonym of *G. cygnoides* (Zeder, 1800) (Yamaguti, 1971: 345). In any case, nowhere in Sinitzyn (1906) does the combination “*C. loossi*” occur and the first author to use it for a species was Stunkard (1929b). This species is probably a larval form of a blood fluke (Sanguinicolidae) which matures in marine fish (Overstreet, personal commun.).

Cercaria roscovita Stunkard, 1932: 333–336, fig. 5.

Syntypes: AMNH 1354.1–1354.12 (3 specimens).

Host: *Littorina saxatilis nigrolineata* (Gastropoda).

Locality: Roscoff, Finistère, France, Atlantic Ocean.

Collector: Unknown, Aug. 1931.

Remarks: ex Stunkard. This species is placed in *Renicola* Cohn, 1904 (Stunkard, 1971: 110).

Cladorchis gigas MacCallum, 1917: 867–870, pls. 108–110, text fig. 1.

Syntypes: AMNH 144 (5 specimens), 160 (9 specimens), 161 (45 specimens), 162 (14 specimens), 524.1–524.38, 525.1–525.56 (1 specimen each).

Host: *Loxodon africanus* (Mammalia).

Locality: Vankerckhovenille, Belgian Congo [Zaire].

Collector: H. Lang and J. Chapin, April, 1912.

Remarks: ex AMNH Belgian Congo Expedition. AMNH 524 was considered the “type” specimen, at least in the collection records (see Feinberg, 1970: 5). No holotype was designated by MacCallum (1917) and there is no evidence to support the idea that AMNH 524 was intended by the author to be the “type”; it is treated here as merely another syntype. This species is a synonym of *Brumptia bicaudata* (Poirier, 1909) (Stunkard, 1929a: 258).

Cyclocoelum phasidi Stunkard, 1929a: 269–271, figs. 36, 37.

Holotype: AMNH 166.

Paratypes: AMNH 531.1–531.8, 532 (1 specimen each).

Host: *Guttera plumifera schubotzi* (Aves).

Locality: Ngayu (AMNH 166) and Niapu (AMNH 531, 532), Belgian Congo [Zaire].

Collector: H. Lang and J. Chapin, Dec. 20, 1909 (AMNH 166), Nov. 1913 (AMNH 531, 532).

Remarks: ex AMNH Belgian Congo Expedition. AMNH 166 was fig. 36 in Stunkard (1929a). This specific name is a nomen nudum from MacCallum (1921: 168). This species is placed in the subgenus *Cyclocoelum* Brandes, 1892 (Dollfus, 1948: 134).

Dictyangium chelydrae Stunkard, 1943: 144–148, figs. 1–6.

Holotype: AMNH 1121.

Paratypes: AMNH 1120.1–1120.35 (33 specimens), 1128.1–1128.14 (22 specimens), 1129.1–1129.15 (7 specimens).

Host: *Chelydra serpentina* (Testudinata).

Locality: Louisiana, United States.

Collector: Unknown.

Remarks: ex Stunkard. AMNH 1121 was fig. 1 in Stunkard (1943). The repository of additional paratypes was not cited. This species is the type of the genus *Dictyangium* Stunkard, 1943, by monotypy (Stunkard, 1943: 144).

Distoma pseudoechinatum Olsson, 1876: 21–22, figs. 45–49.

Syntype: AMNH 22 (1 specimen).

Host: *Larus marinus* (Aves).

Locality: Lund, Sweden.

Collector: P. Olsson, Nov. 29, 1867.

Remarks: ex Olsson Collection. This species is placed in *Stephanoprora* Odhner, 1902 (Gupta, 1963: 125).

Haematotrema parvum Stunkard, 1923a: 373–374.

Lectotype: AMNH 133 (1 specimen).

Paralectotypes: AMNH 117, 121 (1 specimen each), 127 (3 specimens), 133A (1 specimen).

Host: *Chrysemys picta* (Testudinata).

Locality: Cold Spring on Hudson, Putnam Co., New York, United States.

Collector: H. W. Stunkard.

Remarks: ex Stunkard. The species was figured by Stunkard (1923b: figs. 45–49). AMNH 133, the circled specimen on the slide illustrated in fig. 45 in Stunkard (1923a), was marked "type" by Stunkard and was mounted on the same slide as AMNH 133A. Since no holotype was designated by Stunkard (1923a), AMNH 133 is here designated lectotype (fig. 9, herein). The circled specimen on AMNH 127 was fig. 49 in Stunkard (1923a). The repository of additional paralectotypes was not cited. This species is the type of the genus *Haematotrema* Stunkard, 1923, by monotypy (Stunkard, 1923a: 373). This species is placed in *Spirorchis* MacCallum, 1918, as *S. parvus* (Byrd, 1939: 125–126).

Hapalorhynchus gracilis Stunkard, 1922: 2–4, figs. 1, 2.

Holotype: AMNH 125.

Paratypes: AMNH 1269.1–1269.4 (4 specimens).

Host: *Chelydra serpentina* (Testudinata).

Locality: North Judson, Starke Co., Indiana, United States.

Collector: H. W. Stunkard.

Remarks: ex Stunkard. AMNH 125 was fig. 1 and AMNH 1269.2 was fig. 2 in Stunkard (1922). The repository of additional paratypes was not cited. This species is the type of the genus *Hapalorhynchus* Stunkard, 1922, by monotypy (Stunkard, 1922: 1).

Henotosoma haematobium Stunkard, 1922: 5–7, fig. 3.

Holotype: AMNH 126.

Paratypes: AMNH 1264.1–1264.10 (14 specimens).

Host: *Chelydra serpentina* (Testudinata).

Locality: Raleigh, Wake Co., North Carolina, United States.

Collector: Unknown.

Remarks: ex Stunkard. AMNH 126 was the figured specimen in Stunkard (1922). The repository of additional paratypes was not cited. This species is the type of the genus *Henotosoma* Stunkard, 1922, by monotypy (Stunkard, 1922: 5). This species is placed in *Spirorchis* MacCallum, 1918 (Byrd, 1939: 125).

Heterochinostomum magnovatum Stunkard and Haviland, 1924: 7–9, figs. 2–5.

Lectotype: AMNH 138.

Paralectotypes: AMNH 141.1–141.8 (10 specimens).

Host: *Rattus norvegicus* (Mammalia).

Locality: Clason Point Dump, New York City, New York, United States.

Collector: H. W. Stunkard, Nov. 1923.

Remarks: ex Stunkard. Stunkard marked slide AMNH 138, fig. 2 in Stunkard and Haviland (1924), "type" and AMNH 141.1 "co-type." Since no holotype was designated by Stunkard and Haviland (1924), AMNH 138 is here designated lectotype (fig. 10, herein). The repository of additional paralectotypes was not cited. This species is placed in *Echinochasmus* Dietz, 1909, as *E. magnovatus* (Price, 1931: 6).

Himasthla compacta Stunkard, 1960: 541–544, figs. 1, 2, 18–24.

Paratypes: AMNH 1522.1–1522.7 (15 specimens), 1526.1–1526.10 (30 specimens).

Host: *Hydrobia minuta* (Gastropoda) (AMNH 1522) and *Larus argentatus* (Aves) (AMNH 1526).

Locality: Boothbay, Sagadahoc Co., Maine, United States.

Collector: W. R. Welch, Dec. 8, 1959 (AMNH 1522) and Unknown, 1958 (AMNH 1526).
 Remarks: ex Stunkard. The holotype, figs. 1 and 18 in Stunkard (1960), is USNPC 39444 (Stunkard, 1960: 542). AMNH 1522.1 was fig. 21, AMNH 1522.6 was fig. 22, and two specimens from AMNH 1526.2 were figs. 23, 24 in Stunkard (1960). An additional paratype is USNPC 39444.

Himasthla littorinae Stunkard, 1966a: 368–370, figs. 1–4.

Paratypes: AMNH 1263.1–1263.17 (73+ specimens).

Host: *Littorina saxatilis* (Gastropoda).

Locality: Woods Hole, Barnstable Co., Massachusetts, United States, Atlantic Ocean.

Collector: Woods Hole Marine Laboratory, summer 1965.

Remarks: ex Stunkard. The holotype, fig. 1 in Stunkard (1966a), is USNPC 65610 and additional paratypes are USNPC 65611 (Stunkard, 1966a: 370).

Hirudinella beebei Chandler, 1937: 348–352, figs. 1, 2.

Holotype: AMNH 1411.1–1411.120.

Host: *Acanthocybium petus* (Osteichthyes).

Locality: Bermuda, Atlantic Ocean.

Collector: W. Beebe, June 30, 1935.

Remarks: ex Chandler. AMNH 1411 is the holotype by monotypy. This species is a synonym of *H. ventricosa* (Pallas, 1774) (Nigrelli and Stunkard, 1947: 189).

Liolope copulans Cohn, 1902: 877–880, figs. 1–3.

Syntype: AMNH 1490 (1 specimen).

Host: *Cryptobranchus japonicus* (Amphibia).

Locality: Komami, Japan.

Collector: G. W. Müller, 1897.

Remarks: ex Cohn via Pearse to Stunkard. The locality data is from the slide. The repository of additional syntypes was not cited. This species is the type of the genus *Liolope* Cohn, 1902, by monotypy (Cohn, 1902: 877).

Lobatozoum bilobatum Hyman, 1963b: 193–195, fig. 1.

Syntypes: AMNH 480 (2 host tissue samples; many specimens).

Host: *Epinephelus mystacinus* (Osteichthyes).

Locality: 287 m, North Bimini, Atlantic Ocean.

Collector: M. Gordon, Aug. 12, 1952.

Remarks: ex Hyman. The repository of additional syntypes was not cited.

Mesocoelium japonicum Goto and Ozaki, 1930: 77–78, fig. 4.

Syntype: AMNH 1493 (1 specimen).

Host: *Pseudosalmandra stejnegeri* (Amphibia).

Locality: Yawata Machi, Kyushu, Japan.

Collector: A. S. Pearse, April 10, 1929.

Remarks: ex Pearse via Stunkard. This species is a synonym of *M. breviaecum* Ochi in Goto and Ozaki, 1929 (Freitas, 1963: 267).

Mesocoelium pearsei Goto and Ozaki, 1930: 76–77, fig. 3.

Syntype: AMNH 1494 (1 specimen).

Host: *Pseudosalmandra stejnegeri* (Amphibia).

Locality: Kagoshima, Japan.

Collector: A. S. Pearse, Nov. 12, 1929.

Remarks: ex Pearse via Stunkard. This species is a synonym of *M. breviaecum* Ochi in Goto and Ozaki, 1929 (Freitas, 1963: 267).

Microphallus limuli Stunkard, 1951: 308–312, figs. 1–4.

Paratypes: AMNH 1043.1–1043.14 (14+ specimens).

Host: *Limulus polyphemus* (Merostomata).

Locality: Pleasant Bay, Massachusetts, United States, Atlantic Ocean.

Collector: T. H. Waterman, Summer 1950.

Remarks: ex Stunkard. The holotype is USNPC 47592 (Stunkard, 1951: 312). The repository of additional paratypes was cited as USNPC 47592, but only one specimen (the holotype) exists with this catalog number (Pilitt, personal commun.).

Neopechona cablei Stunkard, 1980: 637–640, figs. 1–10.

Holotype: AMNH 927.

Paratypes: AMNH 928 (3 specimens), 929.1–929.50 (167+ specimens).

Host: *Stenotomus chrysops* (Osteichthyes) (AMNH 927, 928, 929.21–929.39, 929.44, 929.46–929.49), *Mitrella lunata* (Gastropoda) (AMNH 929.1–929.20, 929.40–929.43) and *Podocoryne carnea* (Hydrozoa) (AMNH 929.45, 929.50).

Locality: Woods Hole, Barnstable Co. (AMNH 927, 928); Quinsett and Hadley

- Harbors (AMNH 929), Massachusetts, United States, Atlantic Ocean.
 Collector: J. Valois, April 20, 1979 (AMNH 927, 928) and 1977–1979 (AMNH 929).
 Remarks: ex Stunkard. The holotype is the largest specimen on the same slide as AMNH 928 (Stunkard, 1980: 636). AMNH 929.4, 929.26, 929.33, and 929.39 are marked "drawn" and may have served as the basis for several of the figures in Stunkard (1980). AMNH 930–943 are identified on the slides as specimens of other species; Stunkard (1980: 636) erred in listing these lots as being members of this type series. This species is placed in *Opechona* Looss, 1907 (Bray and Gibson, 1990: 176).
- Notocotylus atlanticus* Stunkard, 1966b: 508–513, figs. 1–7.
 Paratypes: AMNH 968.1–968.10 (158+ specimens), 1404.1–1404.4 (39+ specimens).
 Host: *Hydrobia salsa* (Gastropoda).
 Locality: Oyster Pond, Falmouth and Nobska Pond, Woods Hole, Barnstable Co., Massachusetts, United States, Atlantic Ocean.
 Collector: Woods Hole Marine Laboratory, 1963–1966.
 Remarks: ex Stunkard. The holotype, fig. 1 in Stunkard (1966b), is USNPC 61184 and additional paratypes are USNPC 61185 (Stunkard, 1966b: 506).
- Parabascoides yucatanensis* Stunkard, 1938a: 34–37, fig. 1.
 Holotype: AMNH 269.
 Paratypes: AMNH 1026.1–1026.4 (4 specimens).
 Host: *Natalus mexicanus* (Mammalia).
 Locality: Balaam Canche Cave, Chichen Itza, Yucatan, Mexico.
 Collector: A. S. Pearse, June 23, 1936.
 Remarks: ex Stunkard. AMNH 269 was the figured specimen in Stunkard (1938a). The repository of the additional paratype was not cited. This species is the type of the genus *Parabascoides* Stunkard, 1938, by monotypy (Stunkard, 1938a: 34).
- Paramonostomum parvum* Stunkard and Dunihue, 1931: 184–185, fig. 4.
 Syntypes: AMNH 1145.1–1145.8 (7 specimens).
 Host: "Long Island duck" (Aves).
 Locality: Purchased, New York market, New York, United States.
 Collector: Unknown, Dec. 1921.
 Remarks: ex Stunkard. The repository of additional syntypes was not cited.
- Paratelorchis dollfusi* Stunkard and Franz, 1977: 384–385, fig. 1.
 Holotype: AMNH 872.
 Paratypes: AMNH 873, 874 (3 specimens each).
 Host: *Regina alleni* (Squamata).
 Locality: Paynes Prairie, near Gainesville, Alachua Co., Florida, United States.
 Collector: R. Franz.
 Remarks: ex Stunkard. AMNH 872 is the circled specimen on the same slide as AMNH 873 and was the figured specimen in Stunkard and Franz (1977). The repository of additional paratypes was not cited. This species is the type of the genus *Paratelorchis* Stunkard and Franz, 1977, by original designation (Stunkard and Franz, 1977: 383–384). This generic name is preoccupied by *Paratelorchis* Mehra and Bokhari, 1932, so Stunkard (1979: 142) erected *Auritelorchis* as a replacement name. This species is placed in *Telorchis* Luehe, 1899 (MacDonald and Brooks, 1989: 2308).
- Phyllodistomum enterocolpium* Holl, 1930: 450, figs. 1–3.
 Paratype: AMNH 1489 (1 specimen).
 Host: *Diemictylus pyrrhogaster* (Amphibia).
 Locality: Kichijoji, near Tokyo, Japan.
 Collector: A. S. Pearse.
 Remarks: ex Holl via Stunkard. The holotype is USNPC 8083 and additional paratypes are USNPC 8084 (Salley et al., 1978: 46) and in TIU (Holl, 1930: 451). This species is a synonym of *P. (Catoptroides) patellare* (Sturges, 1897) (Pigulevsky in Skrjabin, 1953: 380).
- Plagitura parva* Stunkard, 1933: 134.
 Lectotype: AMNH 178.
 Paralectotypes: AMNH 1389.1–1389.68 (many specimens), 1390.1, 1390.2 (6 specimens).
 Host: *Triturus viridescens* (Amphibia).
 Locality: Cold Spring on Hudson, Putnam Co., New York, United States.
 Collector: Unknown.
 Remarks: ex Stunkard. Stunkard (1936: 356) designated the "type specimen" (fig. 2 in

Stunkard [1936]); fig. 11, herein) as being in AMNH and slide AMNH 178 was marked "type." Since no holotype was designated by Stunkard (1933), his action (1936) constituted a lectotype designation. The species was figured by Stunkard (1936: figs. 2–11), where AMNH 1389.67 was fig. 9 and AMNH 1389.68 was fig. 7. The many specimens from AMNH 1389, primarily cercariae and sporocysts, are considered likely paralectotypes, despite the lack of adequate data on many of the slide labels, because data on several slides match the data from the original species description.

Probolitrema californiense Stunkard, 1935: 362–364, fig. 1.

Holotype: AMNH 179.

Paratypes: AMNH 1487.1, 1487.2 (2 specimens).

Host: *Myliobatus californicus* (Chondrichthyes).

Locality: Elkhorn Slough, Monterey Bay, California, United States, Pacific Ocean.

Collector: G. E. MacGinitie, July 20, 1927.

Remarks: ex Stunkard. The repository of additional paratypes was not cited.

Pterygotomaschalos attenuatus Stunkard, 1923d: 358.

Lectotype: AMNH 656.

Paralectotypes: AMNH 1540.1–1540.8 (11 specimens).

Host: *Chelydra serpentina* (Testudinata).

Locality: Florida, United States.

Collector: C. R. Halter.

Remarks: ex Stunkard. Stunkard marked slide AMNH 656 "type." Since no holotype was designated by Stunkard (1923d), AMNH 656 is here designated lectotype (fig. 12, herein). The species was figured by Stunkard (1924a: fig. 9), where the lectotype was the figured specimen. The repository of additional paralectotypes was not cited. This species is the type of the genus *Pterygotomaschalos* Stunkard, 1923, by monotypy (Stunkard, 1923d: 358). This species is a synonym of *Auridistomum chelydrae* (Stafford, 1900) (Skrjabin, 1962: 31).

Spirorchis elegans Stunkard, 1923b: 192–193, figs. 2, 3.

Holotype: AMNH 134 (1 specimen).

Host: *Pseudemys elegans* (Testudinata).

Locality: Havana, Mason Co., Illinois, United States.

Collector: H. W. Stunkard, Nov. 18, 1913.

Remarks: ex Stunkard. The type series was restricted to the "longer" specimen illustrated in fig. 2, therefore AMNH 134 is the holotype by monotypy (Stunkard, 1923b: 192). The other (non-type) specimen cited by Stunkard (1923b: 192, fig. 3) is AMNH 1516.

Spirorchis picta Stunkard, 1923b: 193–194, fig. 6.

Lectotype: AMNH 131.

Paralectotype: AMNH 1515 (1 specimen).

Host: *Chrysemys picta* (Testudinata).

Locality: Vicinity of New York City, New York, United States.

Collector: H. W. Stunkard.

Remarks: ex Stunkard. Stunkard marked slide AMNH 131, fig. 6 in Stunkard (1923b), "type." Since no holotype was designated by Stunkard (1923b), AMNH 131 is here designated lectotype (fig. 13, herein). The repository of additional paralectotypes was not cited. This species is a synonym of *S. elegans* Stunkard, 1923 (Byrd, 1939: 125).

Spirorchis scripta Stunkard, 1923b: 190–191, figs. 4, 5.

Lectotype: AMNH 128.

Paralectotypes: AMNH 130 (1 specimen), 1511.1–1511.5 (4 specimens).

Host: *Pseudemys scripta* (Testudinata).

Locality: Raleigh, Wake Co., North Carolina, United States.

Collector: Unknown, Nov. 10, 1914 (AMNH 128) and Oct. 21, 1916 (AMNH 130, 1511).

Remarks: ex Stunkard. Stunkard marked slide AMNH 128, fig. 4 in Stunkard (1923b), "type." Since no holotype was designated by Stunkard (1923b), AMNH 128 is here designated lectotype (fig. 14, herein). AMNH 130 was fig. 5 in Stunkard (1923b). The repository of additional paralectotypes was not cited.

Stephanopharynx secundus Stunkard, 1929a: 252–254, figs. 18–20.

Holotype: AMNH 165.

Paratypes: AMNH 153, 528.1–528.9, 529.1–529.9, 530.1–530.5 (1 specimen each).

Host: *Redunca bohor* (Mammalia).

Locality: Faradje, Belgian Congo [Zaire].

Collector: H. Lang and J. Chapin, Feb. 10, 1912.

Remarks: ex AMNH Belgian Congo Expedition. AMNH 165 is an alcohol-preserved specimen. This species is a synonym of *S. compactus* Fiscoeder, 1901 (Eduardo, 1986: 60).

Sterrhurus branchialis Stunkard and Nigrelli, 1934: 535–540, figs. 1, 2.

Lectotype: AMNH 884 (1 specimen).

Paralectotypes: AMNH 884A (6 specimens), 885.1–885.24 (76+ specimens).

Host: *Trichiurus lepturus* (Osteichthyes).

Locality: Near Long Island, New York, United States, Atlantic Ocean.

Collector: Unknown.

Remarks: ex Stunkard. Stunkard marked slide AMNH 884, fig. 2 in Stunkard and Nigrelli (1934), "type." Since no holotype was designated by Stunkard and Nigrelli (1934), AMNH 884 is here designated lectotype (fig. 15, herein). AMNH 884 and 884A are on the same slide, but are clearly distinguished. The repository of additional paralectotypes was not cited. This species is placed in *Lecithochirium* Luehe, 1901 (Gushanskaya in Skrjabin, 1955: 371).

Unicaecum ruszkowskii Stunkard, 1925: 317.

Paralectotypes: AMNH 1272.1–1272.5 (5 specimens), 1273.1–1273.9 (4 specimens).

Host: *Pseudemys scripta* (AMNH 1272) and *P. scripta elegans* (AMNH 1273) (Testudinata).

Locality: Raleigh, Wake Co., North Carolina (AMNH 1272) and New Orleans, Orleans Parish, Louisiana (AMNH 1273), United States.

Collector: Unknown, 1916 (AMNH 1272).

Remarks: ex Stunkard. The species was figured by Stunkard (1927: figs. 1–3). Stunkard (1927) designated the specimen in fig. 1 as the "spécimen type monté en entier." Since no holotype was designated by Stunkard (1925), his action (1927) constituted a lectotype designation. This species is the type of the genus *Unicaecum* Stunkard, 1925, by monotypy (Stunkard, 1925: 317).

Uniserialis breviserialis Stunkard, 1967: 270–273, figs. 1, 3–7.

Paratypes: AMNH 969.1–969.36 (64 specimens).

Host: Domestic "Pekin" (sic) ducklings (Aves).

Locality: Woods Hole, Barnstable Co., Massachusetts, United States.

Collector: Woods Hole Marine Laboratory, Aug. 1965–Aug. 1966.

Remarks: ex Stunkard. The holotype is USNPC 61186 and additional paratypes are USNPC 61187 (Stunkard, 1967: 273). One of the specimens from AMNH 969.18 was fig. 4 in Stunkard (1967). This species is placed in *Notocotylus* Diesing, 1839 (Bisset, 1977: 370).

Vasatrema amydae Stunkard, 1926: 165.

Paralectotype: AMNH 791 (1 specimen).

Host: *Amyda ferox* or *A. spinifera* (Testudinata).

Locality: Indiana or Florida, United States.

Collector: Unknown.

Remarks: ex Stunkard. The "type" specimen was figured by Stunkard (1928: fig. 1). Since no holotype was designated in Stunkard (1926), the designation of the specimen figured in Stunkard (1928) as the type constituted a lectotype designation. The repository of the figured lectotype specimen and of additional paralectotypes was not cited. This species is the type of the genus *Vasatrema* Stunkard, 1926, by monotypy (Stunkard, 1926: 165). *Vasotrema* Stunkard, 1928 (Stunkard, 1928: 303), is an unjustified emendation; the original spelling *Vasatrema* should be retained.

Vasotrema attenuatum Stunkard, 1928: 309–312, figs. 4–6.

Paratypes: AMNH 806.1–806.17 (24 specimens), 807.1–807.5 (5 specimens).

Host: *Amyda ferox* or *A. spinifera* (Testudinata).

Locality: Indiana or Florida, United States.

Collector: H. W. Stunkard.

Remarks: ex Stunkard. The "spécimen type, entièrement monté" was fig. 4 in Stunkard (1928). The repository of the figured holotype specimen was not cited. The genus should be spelled *Vasatrema* (see remarks for the preceding species).

Vasotrema robustum Stunkard, 1928: 312–316, figs. 7–9.

Paratypes: AMNH 808 (1 specimen), AMNH 809.1–809.3 (3 specimens).

Host: *Amyda spinifera* (Testudinata).

Locality: Indiana, United States.

Collector: H. W. Stunkard.

Remarks: ex Stunkard. The "spécimen type, entièrement monté" was fig. 7 in Stunkard (1928). The repository of the figured holotype specimen was not cited. The genus should be spelled *Vasatrema* (see remarks under *V. amydae*).

Wardius zibethicus Barker and East in Barker, 1915: 192–193, pl. 2, fig. 3.

Syntypes: AMNH 1592.1–1592.76 (24 specimens).

Host: *Fiber zibethicus* (Mammalia).

Locality: Nebraska, United States.

Collector: Unknown.

Remarks: ex Dr. Barker's widow via Stunkard. Additional syntypes are USNPC 50562, 51864 (Salley et al., 1978: 146). This species is the type of the genus *Wardius* Barker and East in Barker, 1915, by monotypy (Barker, 1915: 192).

Zalophotrema hepaticum Stunkard and Alvey, 1929: 106–107.

Holotype: AMNH 170.

Paratypes: AMNH 1157.1–1157.25 (3 specimens).

Host: *Zalophus californianus* (Mammalia).

Locality: New York Aquarium, Kings Co., New York, United States.

Collector: Unknown, 1929.

Remarks: ex Stunkard. The species was figured by Stunkard and Alvey (1930: pl. 33). This species is the type of the genus *Zalophotrema* Stunkard and Alvey, 1929, by monotypy (Stunkard and Alvey, 1929: 107).

Zeugorchis longicirrus Odlaug, 1938: 173–175, fig. 1.

Lectotype: AMNH 265.

Host: *Natrix sipedon* (Squamata).

Locality: Louisiana, United States.

Collector: Unknown.

Remarks: ex Odlaug via Stunkard. AMNH 265 is marked "type." Since no holotype was designated by Odlaug (1938), AMNH 265 is here designated lectotype (fig. 16, herein). The repository of the paralectotypes was not cited. This species is placed in *Dasymetra* Nicoll, 1911 (Byrd and Denton, 1938: 388–389).

Zygocotyle ceratosa Stunkard, 1916: 26.

Paratypes: AMNH 1277.1–1277.7 (3 specimens).

Host: *Anas platyrhynchos* (Aves).

Locality: Rock Co., Nebraska, United States.

Collector: Unknown, 1908.

Remarks: ex Stunkard. The holotype was in HCUI (Stunkard, 1916: 26), now USNPC 51865 (Salley et al., 1978: 31). The repository of additional paratypes was not cited. The species was figured by Stunkard (1917: figs. 72–79). This species is the type of the genus *Zygocotyle* Stunkard, 1916, by original designation (Stunkard, 1916: 26), but is a synonym of *Z. lunata* (Diesing, 1836) (Sey, 1991: 264).

CLASS CESTODA

Amphilina bipunctata Riser, 1948: 479–485, figs. 1–8.

Paratype: AMNH 1232 (1 specimen).

Host: Sturgeon (Osteichthyes).

Locality: Dodson, Oregon, United States.

Collector: C. D. Duncan, Aug. 1923.

Remarks: ex Riser via Stunkard. The holotype is USNPC 46362 (Riser, 1948: 479). This species is a synonym of *A. japonica* Goto and Ishii, 1936 (Schmidt, 1986: 468).

Nybelinia lamontae Nigrelli, 1938: 8–10, figs. 3, 4.

Holotype: AMNH 523.

Host: *Xiphias gladius* (Osteichthyes).

Locality: Louisbourg, Cape Breton Island, Nova Scotia, Canada, Atlantic Ocean.

Collector: AMNH Lerner–Cape Breton Island Expedition, July 30, 1936.

Remarks: ex AMNH Dept. Ichthyology. The original catalog number for this specimen was AMNH (Dept. Ichthyology) 13769 (Nigrelli, 1938: 9). AMNH 523 was the figured specimen in Nigrelli (1938).

Oochoristica anniellae Stunkard and Lynch, 1944: 165–168, figs. 1–3.

Lectotype: AMNH 1065.

Paralectotype: AMNH 1061.1–1061.9 (1 specimen).

Host: *Anniella pulchra nigra* (Squamata).

Locality: Castroville, Monterey Co., California, United States.

Collector: C. M. Miller, 1940.

Remarks: ex H. Kirby via Stunkard. Stunkard marked slide AMNH 1065 "type." Since no holotype was designated by Stunkard and Lynch (1944), AMNH 1065 is here designated lectotype.

Oochoristica parva Stunkard, 1938a: 46–48, figs. 6–8.

Holotype: AMNH 270.

Paratype: AMNH 1027.1–1027.3 (1 specimen).

Host: *Coleonyx elegans* (Squamata).

Locality: Gongora, Oxkutzcab, Yucatan, Mexico.

Collector: A. S. Pearse, July 17, 1936.

Remarks: ex Stunkard. AMNH 270 was the figured specimen in Stunkard (1938a). This specific name is preoccupied by *O. parva* (Janicki, 1904) and was replaced by *O. parvula* Stunkard, 1938 (Stunkard, 1938b: 554).

Paratriotaenia oedipomidatis Stunkard, 1965b: 545–547, figs. 1–5.

Syntypes: AMNH 1348.1–1348.7 (10 specimens).

Host: *Oedipomidas oedipus* (Mammalia).

Locality: Primate colony, Tulane University, New Orleans, Louisiana, United States.

Collector: F. Sogandares-Bernal, 1963.

Remarks: ex Stunkard. Stunkard designated USNPC 60933 as "type and cotype" (Stunkard, 1965b: 550), but no specimen on this slide, nor any figure in the publication, is marked in such a way as to indicate that it is the holotype. In the absence of a clearly defined holotype, all members of the type series must be regarded as syntypes. Stunkard (1965c: 996) made this species the type of the genus *Paratriotaenia* Stunkard, 1965, by subsequent designation (Stunkard, 1965b: 550). This was unnecessary as it is the type of the genus by monotypy.

Tetrabothrium norvegicum Olsson, 1868: 3–4, figs. 55–61.

Syntypes: AMNH 11 (4 specimens).

Host: *Spinax niger* (Osteichthyes).

Locality: Bergen, Norway.

Collector: Unknown, Sept. 4, 1867.

Remarks: ex Olsson Collection. This species is the type of the genus *Aporhynchus* Nybelin, 1918, by monotypy as *A. norvegicus* (Schmidt, 1986: 60).

Trienophorus robustus Olsson, 1893: 20–21, figs. 28–32.

Syntypes: AMNH 8.1–8.3 (3 specimens).

Host: *Esox lucius* (Osteichthyes).

Locality: Lake Refsinden, Jemtland, near Hafra, Sweden.

Collector: P. Olsson, Aug. 8, 1875.

Remarks: ex Olsson Collection. This species is a synonym of *T. crassus* Forel, 1868 (Wardle and McLeod, 1952: 638).

PHYLUM RHOMBOZOA

Conocyema adminicula McConnaughey, 1949b: 12–13, pl. 1, fig. 15; pl. 3, figs. 40–45.

Syntypes: AMNH 1085.2 (several specimens).

Host: *Octopus* sp. [= *O. rubescens*, YPM 19459 (Hochberg, personal commun.)] (Cephalopoda).

Locality: 35 fathoms, off Point Loma, San Diego Co., California, United States, Pacific Ocean.

Collector: R/V "E. W. Scripps," Oct. 9, 1947[1946?].

Remarks: ex McConnaughey via Stunkard. These specimens are on the same slide as AMNH 1085.1: syntypes of *Dicyema balamuthi* McConnaughey, 1949 (McConnaughey, 1949b: 10). This is slide L-11-6, cited as being in the author's collection. Additional syntypes [6 slides] were also cited as being in the author's collection (McConnaughey, 1949b: 12), now in SBMNH.

Dicyema balamuthi McConnaughey, 1949b: 10, pl. 4, figs. 46, 47; pl. 5, figs. 56–64; pl. 6, figs. 71–79.

Syntypes: AMNH 1085.1 (several specimens).

Host: *Octopus* sp. [= *O. rubescens*, USNM 816376 (Hochberg, personal commun.)] (Cephalopoda).

Locality: 35 fathoms, off Point Loma, San Diego Co., California, United States, Pacific Ocean.

Collector: R/V "E. W. Scripps," Oct. 9, 1946[1947?].

Remarks: ex McConnaughey via Stunkard. These specimens are on the same slide as

AMNH 1085.2: syntypes of *Conocyema adminicula* McConnaughey, 1949 (McConnaughey, 1949b: 10). This is slide L-2-6 cited as being in the author's collection. Additional syntypes [6 slides] were also cited as being in the author's collection (McConnaughey, 1949b: 10), now in SBMNH. This species is a synonym of *D. apollyoni* Nouvel, 1947 (Hochberg, personal commun.).

Dicyema sullivanii McConnaughey, 1949a: 122-123, figs. 1-7.

Syntypes: AMNH 1086 (several specimens). Host: *Octopus bimaculatus* [= *O. bimaculoides* (Hochberg, personal commun.)] (Cephalopoda).

Locality: Near Rosarita Beach, Baja California Norte, Mexico, Pacific Ocean.

Collector: B. H. McConnaughey, Oct. 24, 1946.

Remarks: ex McConnaughey via Stunkard. AMNH 1086 bears the author's original collection number L-14-13 and also contains specimens of *Dicyemeneea californica* McConnaughey, 1941. Additional syntypes [13 slides] were in the author's collection, L-14 series (McConnaughey, 1949a: 122), now in SBMNH.

Dicyemeneea abelis McConnaughey, 1949b: 5, pl. 1, figs. 8-10; pl. 2, figs. 24-27.

Syntypes: AMNH 1095 (several specimens). Host: *Octopus bimaculatus* [= *O. bimaculoides* (Hochberg, personal commun.)] (Cephalopoda).

Locality: "Coast of Southern and Lower California" [= Laguna Beach], United States, Pacific Ocean.

Collector: B. H. McConnaughey, July 16, 1939.

Remarks: ex McConnaughey via Stunkard. AMNH 1095 bears the author's original collection number 106-7. Additional syntypes [6 slides] were in the author's collection, 106 series (McConnaughey, 1949b: 5), now in SBMNH [3 slides] and the collection of H. Nouvel (presumed to be at FST) [1 slide] with 1 slide unaccounted for (Hochberg, personal commun.).

Dicyemeneea brevicephala McConnaughey, 1941: 64, figs. 4, 5, 7, 9, 10.

Paratypes: AMNH 1097.1, 1097.2 (several specimens).

Host: *Octopus apollyon* [= *O. rubescens*, USNM 816350 (Hochberg, personal commun.)] (Cephalopoda).

Locality: Marine View Rocks, San Mateo Co., California, United States, Pacific Ocean.

Collector: E. Swann, June 8, 1939.

Remarks: ex McConnaughey via Stunkard. The holotype is USNM 22680 (McConnaughey, 1941: 64). Additional paratypes [47 slides] are USNM 22681 [1 slide] and were in the collections of the University of California [1 slide] and the author (McConnaughey, 1941: 64), now distributed as follows: SBMNH [27 slides], H. Nouvel Collection (presumed to be at FST) [2 slides], University of California [1 slide, presumed lost], Short Collection [1 slide], Mohr Collection [2 slides, presumed to be at University of Southern California], with 13 slides unaccounted for (Hochberg, personal commun.).

Dicyemeneea californica McConnaughey, 1941: 63-64, figs. 1-3, 6, 8.

Paratypes: AMNH 1096 (several specimens). Host: *Octopus bimaculatus* [= *O. bimaculoides* (Hochberg, personal commun.)] (Cephalopoda).

Locality: Laguna Beach, Orange Co., California, United States Pacific Ocean.

Collector: B. H. McConnaughey, July 2, 1939.

Remarks: ex McConnaughey via Stunkard. The holotype is USNM 22682 (McConnaughey, 1941: 64) and additional paratypes [1 slide] are USNM 22683 (Cole, personal commun.). Additional paratypes [29 slides] were in the collections of the University of California [1 slide, presumed lost], the author, and H. Nouvel (presumed to be at FST) [1 slide] (McConnaughey, 1949b: 4). 21 slides from the author's collection are now in SBMNH, the remainder have not been traced (Hochberg, personal commun.).

Dicyemeneea granularis McConnaughey, 1949b: 4, pl. 1, fig. 2; pl. 2, figs. 28, 29.

Syntypes: AMNH 1090 (several specimens). Host: *Octopus bimaculatus* (Cephalopoda).

Locality: "Southern California" [= Scripps Point, La Jolla], United States, Pacific Ocean.

Collector: D. Fox, July 2, 1946.

Remarks: ex McConnaughey via Stunkard. AMNH 1090 bears the author's original

collection number L-8-6. Additional syn-
types [39 slides] were in the author's col-
lection, L-8 series (McConnaughey, 1949b:
4), now in SBMNH [37 slides], YPM [1
slide], and the collection of J. Gwinn [1
slide, status unknown] (Hochberg, person-
al commun.). This species is a synonym of
D. californica McConnaughey, 1941
(Hochberg, personal commun.).

PHYLUM PENTASTOMIDA

CLASS PENTASTOMATA

Kiricephalus clelii Riley and Self, 1980: 133,
fig. 5.

Holotype: AMNH 332.

Paratypes: AMNH 329 (4 specimens), 332A
(1 specimen).

Host: *Clelia clelia* (Squamata).

Locality: Trinidad, Trinidad and Tobago.

Collector: C. O. R. Everard, date unknown
(AMNH 332, 332A), Sept. 28, 1970
(AMNH 329).

Remarks: ex Self. AMNH 332 and 332A were
the figured specimens in Riley and Self
(1980).

Kiricephalus gabonensis Riley and Self, 1980:
134, figs. 7a, b.

Holotype: AMNH 140.

Paratype: AMNH 140A (1 specimen).

Host: *Miodon gabonensis* (Squamata).

Locality: Metet, Cameroon.

Collector: H. R. Hill.

Remarks: ex Self. AMNH 140 was the figured
specimen in Riley and Self (1980).

Parasambonia bridgesi Stunkard and Gan-
dal, 1968: 50–54, figs. 1–6.

Holotype: AMNH 12924.

Paratypes: AMNH 12925.1–12925.32 (14
specimens), 12926 (11 specimens).

Host: *Pseudechis porphyriacus* (Squamata).

Locality: Bronx Zoo (host ex Australia), Bronx
Co., New York, United States.

Collector: Unknown, Jan. 14, 1966.

Remarks: ex Stunkard. AMNH 12924 was
fig. 1 in Stunkard and Gandal (1968).
AMNH 12926, cited by Stunkard and
Gandal (1968: 54), was never deposited in
AMNH; its repository is unknown. This
species is the type of the genus *Parasam-*
bonia Stunkard and Gandal, 1968, by

monotypy (Stunkard and Gandal, 1968:
50).

Parasambonia minor Riley and Self, 1982:
128–129, fig. 4.

Holotype: AMNH 192.

Paratypes: AMNH 192.1 (10 specimens),
192.2 (5 specimens), 428A (6 specimens).

Host: *Austrelaps superbus* (Squamata).

Locality: Australia.

Collector: W. Larking, May 26, 1954; Un-
known (AMNH 428A).

Remarks: ex Self. The AMNH catalog num-
bers were cited incorrectly by Riley and
Self (1982). AMNH 192 was fig. 4 in Riley
and Self (1982) and cannot be located in
the AMNH collection.

Porocephalus basiliscus Riley and Self, 1979:
30–31, figs. 2–4.

Holotype: AMNH 248.

Paratypes: AMNH 23 (approx. 75 speci-
mens), 249 (3 specimens).

Host: *Crotalus basiliscus basiliscus* (Squa-
mata).

Locality: Mexico.

Collector: L. M. Klauber, Sept. 15, 1930
(AMNH 248); L. M. Klauber and G. O.
Wiley, 1929–1934 (AMNH 23); H. R. Hill
Collection (AMNH 249).

Remarks: ex Self. The designation of AMNH
249 (3 males) as "holotype" by Riley and
Self (1979) was clearly erroneous; they are
paratypes. The total number of specimens
in AMNH 23 does not agree with the pub-
lished total of 59 specimens divided be-
tween AMNH 23 and BM(NH)
1978.2.20.29–51 (Riley and Self, 1979: 31).

Porocephalus tortugensis Riley and Self, 1979:
30.

Holotype: AMNH 322.1.

Paratypes: AMNH 322.2 (2 specimens), 323
(7 specimens), 324 (4 specimens).

Host: *Crotalus tortugensis* (Squamata).

Locality: Tortuga Island, Mexico.

Collector: L. M. Klauber, June (AMNH 323),
Dec. 21 (AMNH 322, 324), 1937.

Remarks: ex Self. The designation of AMNH
322.2 (2 males) as "holotype" by Riley and
Self (1979) was clearly erroneous; they are
paratypes. Additional paratypes are
BM(NH) 1978.2.20.6–.28 (Riley and Self,
1979: 30).

Raillietiella aegypti Ali, Riley and Self, 1982a: 171–174, figs. 1a, 2c, e, 3.

Holotype: AMNH 371.1.

Paratypes: AMNH 361.1 (2 specimens), 364.1, 370.1 (1 specimen each), 371.2, 371.3 (3 specimens), 372.1, 377.1, 455 (1 specimen each).

Host: *Agama stellio* (Squamata) (AMNH 370.1, 371.1–371.3, 372.1); *A. mutabilis* (AMNH 364.1); *Eumeces schneideri* (Squamata) (AMNH 361.1); *Scincus officinalis* (Squamata) (AMNH 377.1, 455).

Locality: El Aimiriya, Alexandria Governorate (AMNH 361.1), Egypt.

Collector: R. E. Kuntz, March 20 (AMNH 377.1), 25 (AMNH 370.1, 371.1–371.3), 1952, July 7, 1965 (AMNH 372.1); M. N. Kaiser, May 5 (AMNH 361.1), June 1 (AMNH 364.1), 1966.

Remarks: ex Self. AMNH 371.1 was fig. 3a and AMNH 372.1 was fig. 3c in Ali et al. (1982a). The type series was restricted to the specimens listed above on the basis of the selective designation of specimens as types by Ali et al. (1982a: 173; table 3).

Raillietiella bufonis Ali, Riley and Self, 1982b: 282, figs. 1b, c.

Holotype: AMNH 470.11A.

Paratypes: AMNH 470.1–470.10, 470.11B, 470.12–470.25 (approx. 52 specimens).

Host: *Bufo lemur* (Amphibia).

Locality: Puerto Rico.

Collector: Garcia-Diaz, Aug. 31, 1966.

Remarks: ex Self. AMNH 470.11A was fig. 1c in Ali et al. (1982b) and is on the same slide as AMNH 470.11B (3 specimens). AMNH 470.3 was fig. 1b in Ali et al. (1982b).

Raillietiella cartagensis Ali, Riley and Self, 1985: 113, figs. 1c, d.

Holotype: AMNH 521.1A.

Paratypes: AMNH 519.1–519.4 (4 specimens), 521.1B, 521.2, 521.3 (3 specimens).

Host: *Gonatodes* sp. (Squamata) (AMNH 521); *Hemidactylus* sp. (Squamata) (AMNH 519).

Locality: Cartagena, Colombia.

Collector: J. K. Frenkel.

Remarks: ex Self. AMNH 521.1A, fig. 1c, d in Ali et al. (1985), is the circled specimen on the same slide as AMNH 521.1B. AMNH 519.1, 519.2 cannot be located in the AMNH collection.

Raillietiella crotali Ali, Riley and Self, 1984a: 92, fig. 1c.

Holotype: AMNH 516.1.

Paratypes: AMNH 516.2–516.5 (14 specimens).

Host: *Crotalus ruber* (Squamata).

Locality: Pond Island, Gulf of California, Mexico.

Collector: Allan Hancock Expedition (H. R. Hill), Feb. 8, 1936.

Remarks: ex Self. AMNH 516.4 cannot be located in the AMNH collection.

Raillietiella monarchus Ali, Riley and Self, 1984b: 154, fig. 1b.

Paratype: AMNH 503 (1 specimen).

Host: *Gekko monarchis* (Squamata).

Locality: Lan Yu Island, Taiwan.

Collector: R. E. Kuntz, March 1959.

Remarks: ex Self. The holotype is BM(NH) 1982.9.20.2 and additional paratypes are BM(NH) 1982.9.20.1, 1982.9.20.3, 1982.9.20.4 (Ali et al., 1984: 154).

Raillietiella scincoides Ali, Riley and Self, 1984b: 155, fig. 2a.

Holotype: AMNH 471.1.

Paratypes: AMNH 471.2–471.6 (5 specimens).

Host: *Tiliqua scincoides* (Squamata).

Locality: Murray River, South Australia, Australia.

Collector: Unknown.

Remarks: ex Self.

Sebekia microhamus Self and Rego, 1985: 37, fig. 3.

Holotype: AMNH 550.

Paratype: AMNH 550A (1 specimen).

Host: *Caiman sclerops* (Crocodylia).

Locality: Brazil.

Collector: Inst. de Oswalda Cruz.

Remarks: ex Self. Figure 3 in Self and Rego (1985) was said to be that of "AMNH 514"; this was an error as the figured specimen is from the type series.

Sebekia mississippiensis Overstreet, Self and Vliet, 1985: 266–269, figs. 1–6, 9–27.

Holotype: AMNH 529A.1.

Allotype: AMNH 529A.2.

Paratypes: AMNH 194.1–194.7 (approx. 16 specimens), 475, 483 (1 specimen each), 488A.1, 488A.2 (2 specimens), 490A (1 specimen), 491 (approx. 25 specimens), 492A (6 specimens), 501 (approx. 25 spec-

imens), **529A.3**, **530** (1 specimen each), **530A.1**, **530A.2** (2 specimens), **531** (1 specimen), **532.1**, **532.2** (4 specimens), **533.1**, **533.2** (2 specimens), **534** (3 specimens), **535.1–535.3** (16 specimens), **536** (1 specimen), **537.1–537.3** (8 specimens), **538** (1 specimen), **539.1**, **539.2** (7 specimens), **541.1**, **541.2** (6 specimens), **542.1**, **542.2** (4 specimens), **543.1–543.4** (5 specimens), **544** (1 specimen), **545.1**, **545.2** (7 specimens), **564** (1 specimen), **579A** (4 specimens), **580** (2 specimens), **581** (1 specimen).

Host: *Alligator mississippiensis* (Crocodylia) (all but AMNH 579A); *Gambusia affinis* (Osteichthyes) (AMNH 579A).

Locality: St. Augustine (AMNH 475, 491, 492A); Lake Apopka, Orange Co. (AMNH 564), Florida (AMNH 529A, 530, 579A, 580, 581); Blackbeard Island National Wildlife Refuge, Georgia (AMNH 194, 490A); 4 mi east of Grand Chenier, Louisiana, Superior Canal (AMNH 483, 536, 538, 542); Jackson Co. (AMNH 501, 531), Mississippi; Rockefeller Refuge (AMNH 488A, 537, 539, 543–545); 10 mi east of Grand Lake (AMNH 532–534), Vermilion Parish (AMNH 541), Louisiana (AMNH 535), United States (AMNH 530A).

Collector: K. A. Vliet, Feb. (AMNH 492A), Aug. (AMNH 529A, 530), Sept. 10 (AMNH 475, 491), 1983; D. R. Deakins, Sept. 1 (AMNH 194), Sept. 17 (AMNH 490A), 1967 (AMNH 194); R. M. Overstreet, N. Whatley, and A. Glass, Sept. 5 (AMNH 533), Sept. 6 (AMNH 483, 536, 538, 542), 1972; R. M. Overstreet, May 20, 1970 (AMNH 488A, 544), Sept. 5, 1972 (AMNH 532), Sept. 11, 1973 (AMNH 539), 1973–1974 (AMNH 501), date unknown (AMNH 531, 534); Unknown (AMNH 530A); R. M. Overstreet, D. Brooks, and N. Whatley, Sept. 26, (AMNH 535), Sept. 27 (AMNH 541), 1975; R. M. Overstreet, and D. Brooks, date unknown (AMNH 537); R. M. Overstreet, and R. Palmer, May 20, 1970 (AMNH 543); R. M. Overstreet, and N. Whatley, Sept. 11, 1973 (AMNH 545); J. M. Kinsella, June 9, 1971 (AMNH 564); W. Boyce, Fall 1983 (AMNH 580, 581), Jan. 1984 (AMNH 579A).

Remarks: ex Self. AMNH 529A.3, 530 cannot be located in the AMNH collection. Only the holotype and allotype were cor-

rectly cited by Overstreet et al. (1985) (cited catalog numbers AMNH 575 and 576 are not paratypes of this species). Self deposited the remainder of the type series at AMNH and identified them as being paratypes. Using the external evidence provided by Self as to the number of specimens in the type series, these specimens are considered paratypes as per ICZN Recommendation 72B (ICZN, 1985: 141).

Waddycephalus calligaster Riley and Self, 1981: 250.

Holotype: AMNH 425B.

Host: *Dendrelaphis calligasto* (Squamata).

Locality: Australia.

Collector: Unknown.

Remarks: ex Self. AMNH 425B is the holotype by monotypy. This specimen was incorrectly cited as "AMNH 425" by Riley and Self (1981).

Waddycephalus komodoensis Riley and Self, 1981: 250, fig. 7.

Holotype: AMNH 272.

Paratypes: AMNH 184 (19+ specimens), 185 (1 specimen), 272A, 272B (4 specimens).

Host: *Dendrophis pictus* (Squamata).

Locality: Komodo (AMNH 272, 272A, 272B), Loho Liang (AMNH 184), Indonesia; University of Florida, Gainesville, Florida, United States (AMNH 185).

Collector: R. Franz (AMNH 185), Sept. (AMNH 184), Oct. 9 (AMNH 272, 272A, 272B), 1969.

Remarks: ex Self. Three lots were cited by Riley and Self (1981) as "AMNH 272"; they have since been segregated. AMNH 272 was the figured specimen in Riley and Self (1981).

Waddycephalus longicauda Riley and Self, 1981: 248, fig. 4a.

Holotype: AMNH 188.

Paratypes: AMNH 188A (6 specimens).

Host: *Aspedites melanocephalus* (Squamata).

Locality: Laura, North Queensland, Australia.

Collector: H. F. W. Ehrmann, March 1968.

Remarks: ex Self. AMNH 188 was the figured specimen in Riley and Self (1981). The type series comprises 11 specimens, but only 7 are in AMNH.

Waddycephalus porphyriacus Riley and Self, 1981: 248.

Holotype: AMNH 190A.1.

Paratypes: AMNH 190A.2 (7 specimens).

Host: *Pseudechis porphyriacus* (Squamata).

Locality: Unknown.

Collector: Unknown, Oct. 1917.

Remarks: ex Self. The type series was cited as "AMNH 190A" by Riley and Self (1981) but the holotype was clearly indicated and has since been segregated. Only 7 of the 8 paratypes are in AMNH.

Waddycephalus punctulatus Riley and Self, 1981: 250, fig. 9.

Holotype: AMNH 183.

Paratypes: AMNH 183A (5 specimens), 193 (7 specimens), 426 (3 specimens).

Host: *Dendrophis punctulatus* (Squamata).

Locality: Wallaville (AMNH 183, 183A, 183B), Australia (AMNH 193, 426).

Collector: T. L. Bancroft, June 1933 (AMNH 183, 183A, 183B); W. M. Vestjens, Nov. 1, 1971 (AMNH 193); Unknown (AMNH 426).

Remarks: ex Self. AMNH 183 was fig. 9a and a specimen from AMNH 183A was fig. 9b in Riley and Self (1981). Several type lots were cited as "AMNH 183" by Riley and Self (1981), but they have since been segregated. The type series comprises 19 specimens, but only 16 are in AMNH.

Waddycephalus scutata Riley and Self, 1981: 248.

Holotype: AMNH 410A.

Paratypes: AMNH 410B (9 specimens).

Host: *Notechis scutata* (Squamata).

Locality: St. Francis Island, South Australia, Australia.

Collector: Unknown.

Remarks: ex Self. The type series was cited as "AMNH 410" by Riley and Self (1981) but the holotype was clearly indicated and has since been segregated. The type series comprises 13 specimens, but only 10 are in AMNH.

Waddycephalus superbus Riley and Self, 1981: 246–248, figs. 2b, 4b.

Holotype: AMNH 409.

Paratypes: AMNH 179 (2 specimens), 409.1 (6 specimens).

Host: *Austrelaps superbus* (Squamata).

Locality: Green Bay, Tasmania (AMNH 409, 409.1), Cape Barren Islands, Bass Straits (AMNH 179), Australia.

Collector: Unknown (AMNH 409, 409.1),

University of Adelaide, 1912 (AMNH 179).

Remarks: ex Self. AMNH 409 was fig. 4b and a specimen from AMNH 179 was fig. 2b in Riley and Self (1981). Four specimens were cited as being in AMNH 179, only 2 are present.

PHYLUM GASTROTRICHA

CLASS CHAETONOTOIDEA

Chorododasys antennatus Rieger, Ruppert, Rieger, and Schoepfer-Sterrer, 1974: 219–237, figs. 1–8, 9a–c, 10.

Holotype: AMNH 967.

Locality: Off Beaufort, North Carolina, United States, Atlantic Ocean.

Collector: B. Coull, May 1970.

Remarks: ex Rieger et al. This species is placed in *Xenodasys* Swedmark, 1967 (Kisielewski, 1987: 856).

Heteroxenotrichula arcassonensis Ruppert, 1979: 41, figs. 23, 24.

Holotype: AMNH 95.

Paratype: AMNH 96 (1 specimen).

Locality: Cap Ferret Beach, 44°35'42"N, 01°35'09"W, Arcachon, France, Atlantic Ocean.

Collector: E. E. Ruppert, Nov. 3–18, 1971.

Remarks: ex Ruppert.

Heteroxenotrichula squamosa Wilke, 1954: 522–524, figs. 17, 18, 19a–c.

Neotype: AMNH 92.

Locality: Bogue Inlet Beach, 34°38'30"N, 77°05'23"W, Emerald Isle, North Carolina, United States, Atlantic Ocean.

Collector: E. E. Ruppert, Aug. 2–14, 1971.

Remarks: ex Ruppert. The neotype was designated by Ruppert (1979: 38). The slide data indicate that this specimen came from Plage d'Eyrac, France, in Dec. 1971, contradictory to the published data given above.

Heteroxenotrichula transatlantica Ruppert, 1979: 33–36, figs. 15–17.

Holotype: AMNH 90.

Locality: Bogue Inlet Beach, 34°38'30"N, 77°05'23"W, North Carolina, United States, Atlantic Ocean.

Collector: E. E. Ruppert, Sept. 1974.

Remarks: ex Ruppert.

Heteroxenotrichula wilkei Ruppert, 1979: 41, figs. 21, 22.

Holotype: AMNH 93.

Paratype: AMNH 94 (1 specimen).

Locality: Cap Ferret Beach, 44°35'42"N, 01°35'09"W (AMNH 93) and Plage d'Eyrac, 44°40'N, 01°30'06"W (AMNH 94), Arcachon, France, Atlantic Ocean.

Collector: E. E. Ruppert, Nov. 3–18, 1971 (AMNH 93) and Dec. 1971 (AMNH 94).

Remarks: ex Ruppert.

Xenotrichula affinis Remane, 1934: 477, fig. 4.

Neotype: AMNH 91.

Locality: Plage d'Eyrac, 44°40'N, 01°30'06"W, Arcachon, France, Atlantic Ocean.

Collector: E. E. Ruppert, Dec. 1971.

Remarks: ex Ruppert. The neotype was designated by Ruppert (1979: 36). This species is placed in *Heteroxenotrichula* Wilke, 1954 (Ruppert, 1979: 36).

Xenotrichula carolinensis Ruppert, 1979: 22, figs. 10–12.

Holotype: AMNH 86.

Paratype: AMNH 87 (1 specimen).

Locality: Bogue Inlet Beach, 34°38'30"N, 77°05'23"W, Emerald Isle, North Carolina, United States, Atlantic Ocean.

Collector: E. E. Ruppert, Aug. 2–14, 1971 (AMNH 87) and Aug. 1974 (AMNH 86).

Remarks: ex Ruppert. This species is a synonym of *Xenotrichula intermedia* Remane, 1934 (Balsamo et al., 1992: 496).

Xenotrichula lineata Schrom, 1972: 307–310, fig. 9.

Neotype: AMNH 88.

Locality: Bogue Inlet Beach, 34°38'30"N, 77°05'23"W, Emerald Isle, North Carolina, United States, Atlantic Ocean.

Collector: E. E. Ruppert, Aug. 2–14, 1971.

Remarks: ex Ruppert. The neotype was designated by Ruppert (1979: 27) as the species was originally described from living material (no specimens deposited as types).

Xenotrichula pygmaea Remane, 1934: 477, fig. 5.

Neotype: AMNH 97.

Locality: Bogue Inlet Beach, 34°38'30"N, 77°05'23"W, Emerald Isle, North Carolina, United States, Atlantic Ocean.

Collector: E. E. Ruppert, July 1974.

Remarks: ex Ruppert. The neotype was designated by Ruppert (1979: 45). This species is placed in *Heteroxenotrichula* Wilke, 1954 (Ruppert, 1979: 45).


Xenotrichula subterranea Remane, 1934: 475–477, fig. 3.

Neotype: AMNH 89.


Locality: Cap Ferret Beach, 44°35'42"N, 01°35'09"W, near Arcachon, France, Atlantic Ocean.

Collector: E. E. Ruppert, Nov. 19, 1971.


Remarks: ex Ruppert. The neotype was designated by Ruppert (1979: 31). This species is placed in *Heteroxenotrichula* Wilke, 1954 (Ruppert, 1979: 31).



Figs. 1-3. 1. *Leptoplana inquieta* Heath and McGregor, 1912 (neotype—AMNH 369). 2. *Planocera burchami* Heath and McGregor, 1912 (neotype—AMNH 382). 3. *Stylochoplana longastyletta* Freeman, 1933 (neotype—AMNH 370). Scale bars = 1 mm.



Figs. 4–6. 4. *Polystoma floridanum* Stunkard, 1924 (lectotype—AMNH 1464). Scale bar = 0.1 mm. 5. *Polystoma multifalx* Stunkard, 1923 (lectotype—AMNH 883.1). Scale bar = 1 mm. 6. *Acanthochasmus diploporus* Stunkard, 1931 (lectotype—AMNH 875). Scale bar = 0.1 mm.



Figs. 7–12. 7. *Amphistomum bicaudatum* Poirier, 1909 (lectotype—AMNH 526). Scale bar = 1 mm. 8. *Ascocotyle (Parascocotyle) diminuta* Stunkard and Haviland, 1924 (lectotype—AMNH 137). Scale bar = 0.1 mm. 9. *Haematotrema parvum* Stunkard, 1923 (lectotype—AMNH 133). Scale bar = 0.1 mm. 10. *Heterochinostomum magnovatum* Stunkard and Haviland, 1924 (lectotype—AMNH 138). Scale bar = 0.1 mm. 11. *Plagitura parva* Stunkard, 1933 (lectotype—AMNH 178). Scale bar = 0.1 mm. 12. *Pterygotomaschalos attenuatus* Stunkard, 1923 (lectotype—AMNH 656). Scale bar = 0.1 mm.



Figs. 13–16. 13. *Spirorchis picta* Stunkard, 1923 (lectotype—AMNH 131). 14. *Spirorchis scripta* Stunkard, 1923 (lectotype—AMNH 128). 15. *Sterrhurus branchialis* Stunkard and Nigrelli, 1934 (lectotype—AMNH 884). 16. *Zeugorchis longicirrus* Odlaug, 1938 (lectotype—AMNH 265). Scale bars = 0.1 mm.

REFERENCES

- Ali, J. H., J. Riley, and J. T. Self
 1982a. A description of a new species of *Raillietiella* (Pentastomida: Cephalobaenida) from Egyptian lizards with a reassessment of the taxonomic status of *Raillietiella geckonis* (Diesing, 1850) Sambon, 1910 and *Raillietiella affinis* Bovien, 1927. *Syst. Parasitol.* 4(2): 169–180, 4 figs.
 1982b. Amphibians as definitive hosts for pentastomids: *Raillietiella bufonis* n. sp. from *Bufo lemur* in Puerto Rico and a reassessment of *Raillietiella indica* Gerdolst, 1921. *Ibid.* 4(3): 279–284, 1 fig.
 1984a. A revision of the taxonomy of pentastomid parasites (genus *Raillietiella* Sambon, 1910) from American snakes and amphisbaenians. *Ibid.* 6(2): 87–97, 2 figs.
 1984b. Further observations of blunt-hooked raillietiellids (Pentastomida: Cephalobaenida) from lizards with descriptions of three new species. *Ibid.* 6(2): 147–160, 4 figs.
 1985. A review of the taxonomy and systematics of the pentastomid genus *Raillietiella* Sambon, 1910 with a description of a new species. *Ibid.* 7(2): 111–123, 3 figs.
- Alvey, C. H.
 1933. *Sphyrnura oligorchis* n. sp. from *Necturus maculosus*. *J. Parasitol.* 20(3): 140.
 1936. The morphology and development of the monogenetic trematode *Sphyrnura oligorchis* (Alvey 1933) and description of *Sphyrnura polyorchis* n. sp. *Parasitology* 28(2): 229–253, pl. 10, 11.
- Balsamo, M., M. A. Todaro, and P. Tongiorgi
 1992. Marine gastrotrichs from the Tuscan Archipelago (Tyrrhenian Sea): II. Chaetonotida, with description of three new species. *Boll. Zool.* 59(4): 487–498, 10 figs.
- Barker, F. D.
 1915. Parasites of the American muskrat (*Fiber zibethicus*). *J. Parasitol.* 1(4): 184–197, 2 pls., 4 text figs.
- Baugh, S. C.
 1956. Contributions to our knowledge of digenetic trematodes II. *Proc. Natl. Acad. Sci. India Sect. B* 26(5): 295–313, 4 figs.
- Bisset, S. A.
 1977. *Notocotylus tadornae* n. sp. and *Notocotylus gippyensis* (Beverly-Burton, 1958) (Trematoda: Notocotylidae) from waterfowl in New Zealand: morphology, life history and systematic relations. *J. Helminthol.* 51(4): 365–372, 6 figs.
- Boyko, C. B.
 1994. Catalog of Recent type specimens in the Department of Invertebrates, American Museum of Natural History. I. Microinvertebrates (Phyla Sarcomastigophora, Gnathostomula, Gastrotricha, Rotifera, and Tardigrada). *Am. Mus. Novitates* 3106: 44 pp.
- Boyko, C. B., and W. E. Sage III
 1996. Catalog of Recent type specimens in the Department of Invertebrates, American Museum of Natural History. II. Mollusca part 1 (Class Aplacophora, Polyplacophora, Gastropoda (Subclass Opisthobranchia), Bivalvia, and Scaphopoda. *Am. Mus. Novitates* 3170: 50 pp.
- Bray, R. A., and D. I. Gibson
 1990. The Lepocreadiidae (Digenea) of fishes of the north-east Atlantic: review of the genera *Opechona* Looss, 1907 and *Prodistomum* Linton, 1910. *Syst. Parasitol.* 15(3): 159–202, 10 figs.
- Brooks, D. R.
 1980. Revision of the Acanthostomidae Poche, 1926 (Digenea: Cryptogonimidae). *J. Linn. Soc. London Zool.* 70(4): 313–382, 28 figs.
- Brooks, D. R., and B. Holcman
 1993. Revised classification and phylogenetic hypothesis for the Acanthostomidae Looss, 1899 (Digenea: Opisthorchiiformes: Cryptogonimidae). *Proc. Biol. Soc. Washington* 106(2): 207–220, 9 figs.
- Brooks, D. R., and M. A. Mayes
 1975. Platyhelminths of Nebraska turtles with descriptions of two new species of spirorchids (Trematoda: Spirorchidae). *J. Parasitol.* 61(3): 403–406, 2 figs.
- Brusca, R. C., and G. J. Brusca
 1990. Invertebrates. Sunderland, MA: Sinauer Associates. 922 pp.
- Byrd, E. E.
 1939. Studies on the blood flukes of the family Spirorchidae. Part II. Revision of the family and description of new species. *J. Tennessee Acad. Sci.* 14(1): 116–161, 12 figs.
- Byrd, E. E., and J. F. Denton
 1938. New trematodes of the subfamily Reniferinae, with a discussion of the systematics of the genera and species assigned to the subfamily group. *J. Parasitol.* 24(5): 379–401, 17 figs.

- Cannon, L. R. G.
1987. Two new rhabdocoel turbellarians, *Umagilla pacifica* sp. n. and *U. karlingi* sp. n. (Umagillidae), endosymbiotic with holothurians (Echinodermata) from the Great Barrier Reef; and a discussion of sclerotic structures in the female system of the Umagillidae. *Zool. Scr.* 16(4): 297–303, 3 figs.
- Carpenter, J. H.
1971. Systematics and ecology of cave planarians of the United States. *Diss. Abstr.* *Int. B. Sci. Eng.* 32(2): 1283–1284.
- Chandler, A. C.
1923. Three new trematodes from *Amphiuma means*. *Proc. U.S. Natl. Mus.* 63(3): 1–7, 5 figs.
1937. A new trematode, *Hirudinella beebei*, from the stomach of a Bermuda fish, *Acanthocybium petus*. *Trans. Am. Microsc. Soc.* 56(3): 348–354, 2 figs.
- Cohn, L.
1902. Zwei neue Distomen. *Centrl. Bakter., Parasiten., Infektion, Originale* 32(10): 877–882, 5 figs.
- Crezée, M.
1975. Monograph of the Solenofilomorphidae (Turbellaria: Acoela). *Int. Rev. Gesamten Hydrobiol.* 60(6): 769–845, 24 figs.
- Curino, A. C., and N. J. Cazzaniga
1993. A new species of freshwater planaria from Chile (Platyhelminthes: Tricladida), with a nomenclatural note on *Girardia festae* (Borelli, 1898). *Proc. Biol. Soc. Washington* 106(4): 633–644, 9 figs.
- Dean, H. K.
1977. Two new Kalyptorhynchia (Turbellaria) from the coast of northern New England, USA. *Zool. Scr.* 6(4): 265–274, 12 figs.
1980. *Parathylacorhynchus reidi* gen. et sp. n., a schizorhynchid (Turbellaria, Kalyptorhynchia) from the coast of Maine, USA. *Ibid.* 9(1): 5–9, 5 figs.
- de Núñez, M. O.
1993. Life-history studies of heterophyid trematodes in the neotropical region: *Ascocotyle (Phagicola) diminuta* (Stunkard and Haviland, 1924) and *A. (P.) angrense* Travassos, 1916. *Syst. Parasitol.* 24(3): 191–199, 12 figs.
- Doe, D. A.
1974. Two new *Proschizorhynchus* species from the coast of Massachusetts, USA (Turbellaria, Kalyptorhynchia). *Zool. Scr.* 3(3): 101–110, 17 figs.
- Doe, D. A., and R. M. Rieger
1977. A new species of the genus *Retronectes* (Turbellaria, Catenulida) from the coast of North Carolina, U.S.A. *Mikrofauna Meeresboden* 66: 10 pp., 2 figs.
- Dollfus, R. P.
1948. Sur deux monostomes (Cyclocoelidae) pourvus d'une ventrose ventrale. Observations sur la classification des Cyclocoeloidea Albert Henry 1923, liste de leurs notes, répartition géographique. *Ann. Parasitol. Hum. Comp.* 23(3/4): 129–199, 14 figs.
- Doss, M. A.
1968. Index-catalogue of medical and veterinary zoology. Subjects: Trematoda and trematode diseases. Part 9: Checklist of generic, specific, and subspecific names A-M. Washington, D.C.: U.S. Gov. Printing Office. 382 pp.
- Eduardo, S. L.
1986. The taxonomy of the family Paramphistomidae Fischöder, 1901 with special reference to the morphology of species occurring in ruminants. VIII. The genera *Stephanopharyx* Fischöder, 1901 and *Balanorchis* Fischöder, 1901. *Syst. Parasitol.* 8(1): 57–69, figs. 163–170.
- Euzet, L., and J. P. Trilles
1961. Présence de *Declidophora minor* (Olsson 1868) en Méditerranée (Monogenea Polyopisthocotylea). *Comm. Int. Explor. Sci. Mer Méditerranée, Rapp. P. V.* 16(2): 291–295, 4 figs.
- Faubel, A.
1983. The Polycladidia, Turbellaria proposal and establishment of a new system[.] Part I. The Acotylea. *Mitt. Hamburgischen Zool. Mus. Inst.* 80: 17–121, 39 figs.
1984. The Polycladidia, Turbellaria proposal and establishment of a new system[.] Part II. The Cotylea. *Ibid.* 81: 189–259, 15 figs.
- Feinberg, H. S.
1970. A Catalogue of Type Specimens in the Department of Living Invertebrates The American Museum of Natural History, New York, New York, U.S.A. The Phyla Platyhelminthes and Gnathostomulida. New York: Dep. of Living Invertebrates, *Am. Mus. Nat. Hist.* 66 pp.
- Fleming, L. C., and M. D. B. Burt
1978. Revision of the turbellarian genus *Ecotocotyla* (Seriata, Monocelididae) associated with the crabs *Chionoecetes op-*

- ilio* and *Hyas araneus*. J. Fish. Res. Board Can. 35(9): 1223–1233, 5 figs.
- Freeman, D.
1933. The polyclads of the San Juan region of Puget Sound. Trans. Am. Microsc. Soc. 52(2): 107–146, 40 figs.
- Freitas, J. F. T.
1963. *Reisão da família Mesocoeliidae* Dollfus, 1933 (Trematoda). Mem. Inst. Oswaldo Cruz Rio de Janeiro 61(2): 177–312, 307 figs.
- Gibson, D. I., and R. A. Bray
1977. The Azygiidae, Hirudinellidae, Ptychogonimidae, Sclerodistomidae and Syncoeliidae of fishes from the north-east Atlantic. Bull. Brit. Mus. (Nat. Hist.), Zool. ser. 32(6): 167–245, 19 figs.
- Goto, S., and Y. Ozaki
1930. Brief notes on new trematodes III. Japanese J. Zool. 3(1): 73–82, 7 figs.
- Gupta, R.
1963. On *Stephanoprora nigerica* sp. nov., with a brief review of the genus *Stephanoprora* Odhner 1902 (Trematoda: Echinostomatidae). Zool. Anz. 170(3/4): 117–130, 3 figs.
- Heath, H., and E. A. McGregor
1912. New polyclads from Monterey Bay, California. Proc. Acad. Nat. Sci. Philadelphia 64(8): 455–488, pls. 12–18 (44 figs.), 12 text figs.
- Holl, F. J.
1930. *Phyllodistomum enterocolpium*, a new trematode from *Diemictylus pyrrhogaster* Boie. Annot. Zool. Japonenses 12(2): 449–451, 3 figs.
- Holleman, J. T., and C. Hand
1962. A new species, genus, and family of marine flatworms (Turbellaria: Tricladida, Maricola) commensal with mollusks. Veliger 5(1): 20–22, 3 figs.
- Hyman, L. H.
1931. Studies on the morphology, taxonomy, and distribution of North American triclad Turbellaria. V. Description of two new species. Trans. Am. Microsc. Soc. 50(4): 336–343, 8 figs.
1934. Report on triclad Turbellaria from Indian Tibet. Mem. Connecticut Acad. Arts Sci. 10(2): 5–12, 2 pls.
1935. Studies on the morphology, taxonomy, and distribution of North American triclad Turbellaria. VI. A new dendrocoelid from Montana, *Dendrocoelopsis vaginatus* n. sp. Trans. Am. Microsc. Soc. 54(4): 338–345, 5 figs.
1937a. Studies on the morphology, taxonomy, and distribution of North American triclad Turbellaria. VII. The two species confused under the name *Phagocata gracilis*, the validity of the generic name *Phagocata* Leidy 1847, and its priority over *Fonticola* Komarék 1926. Ibid. 56(3): 298–310, 6 figs.
1937b. Studies on the morphology, taxonomy, and distribution of North American triclad Turbellaria. VIII. Some cave planarians of the United States. Ibid. 56(4): 457–477, 21 figs.
1938a. Land planarians from Yucatan. Carnegie Inst. Washington Publ. 491: 23–32, 9 figs.
1938b. Faunal notes. Bull. Mount Desert Island Biol. Lab. 1938: 24–25.
1938c. North American Rhabdocoela and Alloeocoela. III. *Mesotoma* [sic] *arctica*, n. sp., from northern Canada. Am. Mus. Novitates 1005: 8 pp., 6 figs.
1939a. New species of flatworms from North, Central, and South America. Proc. U.S. Natl. Mus. 86(3055): 419–439, figs. 47–51.
1939b. Land planarians from the Hawaiian Islands. Arch. Zool. Exp. Gén. 80, Notes Rev. 3(12): 116–124, 9 figs.
1939c. Some polyclads of the New England coast, especially of the Woods Hole region. Biol. Bull. (Woods Hole) 76(2): 127–152, 18 figs.
1939d. North American Rhabdocoela and Alloeocoela–IV. *Mesostoma macropenis*, new species, from Douglas Lake, Michigan. Am. Midland Nat. 21(3): 646–650, 4 figs.
1939e. North American triclad Turbellaria. X. Additional species of cave planarians. Trans. Am. Microsc. Soc. 58(3): 276–284, 9 figs.
1939f. North American Rhabdocoela and Alloeocoela. V. Two new mesostomine rhabdocoels. Am. Midland Nat. 22(3): 629–636, 8 figs.
1939g. Acoel and polyclad Turbellaria from Bermuda and the Sargassum. Bull. Bingham Oceanogr. Collect. Yale Univ. 7(1): 36 pp., 38 figs.
1941. Terrestrial flatworms from the Canal Zone, Panama. Am. Mus. Novitates 1105: 11 pp., 17 figs.
1943. Endemic and exotic land planarians in the United States with a discussion of necessary changes of names in the Rhynchodemidae. Ibid. 1241: 21 pp., 25 figs.
1944. Marine Turbellaria from the Atlantic coast of North America. Ibid. 1266: 15 pp., 16 figs.

1945. North American triclad Turbellaria, XI. New, chiefly cavernicolous, planarians. *Am. Midland Nat.* 34(2): 475-484, 8 figs.
1950. A new Hawaiian polyclad, *Stylocho-plana inquilina*, with commensal habits. *Occas. Pap. Bernice P. Bishop Mus.* 20(4): 55-58, 1 fig.
- 1953a. The polyclad flatworms of the Pacific coast of North America. *Bull. Am. Mus. Nat. Hist.* 100(2): 269-392, 161 figs.
- 1953b. Some polyclad flatworms from the Galapagos Islands. *Allan Hancock Pacific Exped.* 15(2): 183-210, 14 figs.
- 1954a. The polyclad genus *Pseudoceros*, with special reference to the Indo-Pacific region. *Pacific Sci.* 8(2): 219-225, 2 figs.
- 1954b. Some land planarians of the United States and Europe, with remarks on nomenclature. *Am. Mus. Novitates* 1667: 21 pp., 7 figs.
- 1954c. North American triclad Turbellaria, XIII: three new cave planarians. *Proc. U.S. Natl. Mus.* 103(3333): 563-573, figs. 65-68.
- 1954d. A new marine triclad from the coast of California. *Am. Mus. Novitates* 1679: 5 pp., 3 figs.
- 1955a. The polyclad flatworms of the Pacific coast of North America: additions and corrections. *Ibid.* 1704: 11 pp., 8 figs.
- 1955b. Descriptions and records of fresh-water Turbellaria from the United States. *Ibid.* 1714: 36 pp., 22 figs.
- 1955c. Some polyclad flatworms from the West Indies and Florida. *Proc. U.S. Natl. Mus.* 104(3341): 115-150, figs. 81-89.
- 1955d. Some polyclad flatworms from Polynesia and Micronesia. *Ibid.* 105(3352): 65-82, figs. 1-5.
- 1955e. Miscellaneous marine and terrestrial flatworms from South America. *Am. Mus. Novitates* 1742: 33 pp., 42 figs.
- 1955f. A further study of the polyclad flatworms of the West Indian region. *Bull. Mar. Sci. Gulf Caribbean* 5(4): 259-268, 8 figs.
1956. North American triclad Turbellaria. 15. Three new species. *Am. Mus. Novitates* 1808: 14 pp., 17 figs.
- 1957a. North American Rhabdocoela and Alloeocoela. 6. A further study of *Mesostoma*. *Ibid.* 1829: 15 pp., 17 figs.
- 1957b. A few turbellarians from Trinidad and the Canal Zone, with corrective remarks. *Ibid.* 1862: 8 pp., 6 figs.
- 1959a. A further study of Micronesian polyclad flatworms. *Proc. U.S. Natl. Mus.* 108(3410): 543-597, 17 figs.
- 1959b. Some Australian polyclads (Turbellaria). *Rec. Australian Mus.* 25(1): 1-17, 21 figs.
- 1959c. On two fresh-water planarians from Chile. *Am. Mus. Novitates* 1932: 11 pp., 12 figs.
- 1959d. Some Turbellaria from the coast of California. *Ibid.* 1943: 17 pp., 19 figs.
1960. New and known umagillid rhabdocoels from echinoderms. *Ibid.* 1984: 14 pp., 13 figs.
1962. Some land planarians from Caribbean countries. *Ibid.* 2110: 25 pp., 28 figs.
- 1963a. North American triclad Turbellaria. 16. Fresh-water planarians from the vicinity of Portland, Oregon. *Ibid.* 2123: 5 pp., 3 figs.
- 1963b. Notes on a didymozoid trematode from the Bahama Islands. *Bull. Mar. Sci. Gulf Caribbean* 13(2): 193-196, 1 fig.
1964. North American Rhabdocoela and Alloeocoela 7. A new seriate alloeocoel, with corrective remarks on alloeocoels. *Trans. Am. Microsc. Soc.* 83(2): 248-251, 5 figs.
- [ICZN] International Commission on Zoological Nomenclature
1985. International Code of Zoological Nomenclature (3rd ed.). *Int. Trust for Zool. Nomencl., etc.* 338 pp.
- Janssen, C. N., and A. Faubel
1992. Revision of the subfamily Mesostominae, Rhabdocoela, Platyhelminthes. *Mitt. Hamburgischen Zool. Mus. Inst.* 89: 7-47, 15 figs.
- Jokiel, P. L., and S. J. Townsley
1975. Biology of the polyclad *Prosthlostomum* (*Prosthlostomum*) sp., a new coral parasite from Hawaii. *Pacific Sci.* 28(4): 361-373, 7 figs.
- Kawakatsu, M.
1986. Land planarians from West Malaysia (Turbellaria; Tricladida; Terricola). *Bull. Biogeogr. Soc. Japan* 41(1): 1-15, 7 figs.
- Kenk, R.
1974. Index of the genera and species of the freshwater triclads (Turbellaria) of the world. *Smithson. Contrib. Zool.* 183: 90 pp.
1989. Revised list of the North American freshwater planarians (Platyhelminthes: Tricladida: Paludicola). *Ibid.* 476: 10 pp.
- Kisielewski, J.
1987. New records of marine Gastrotricha from the French coasts of Manche and Atlantic I. Macrodasysida, with descriptions of seven new species. *Bull. Mus. Natl. Hist. Nat., Paris, 4^e sér.* 9, section A, 4: 837-877, 1 pl., 10 text figs.

- Kolasa, J.
 1984. Predation on mosquitoes by juveniles of *Mesostoma* spp. (Turbellaria). Freshwater Invertebr. Biol. 3(1): 42–47, 2 figs.
 1987. Population growth in some *Mesostoma* species (Turbellaria) predatory on mosquitoes. Freshwater Biol. 18(2): 205–212, 4 figs.
- Kolasa, J., and S. S. Schwartz
 1988. Two new *Mesostoma* species (Turbellaria, Rhabdocoela) from Australia. Zool. Scr. 17(4): 329–335, 11 figs.
- Lamothe-A., R., and G. Ponciano-R.
 1986. Revision de la subfamilia Acanthostominae Nicoll, 1914 y establecimiento de dos nuevos generos. An. Inst. Biol. Univ. Nac. Autón. Mexico 56(1985), Ser. Zool. 2: 301–322, 8 figs.
- MacCallum, G. A.
 1917. A new species of trematode (*Cladorchis gigas*) parasitic in elephants. Bull. Am. Mus. Nat. Hist. 37(36): 865–871, pls. 108–110, 1 text fig.
 1921. Studies in Helminthology. Zoopathologica 1(6): 137–284, figs. 69–151.
- MacDonald, C. A., and D. R. Brooks
 1989. Revision and phylogenetic analysis of the North American species of *Telorchis* Luehe, 1899 (Cercomeria: Trematoda: Digenea: Telorchidae). Can. J. Zool. 67(9): 2301–2320, 26 figs.
- McAllister, C. T., S. E. Trauth, and L. W. Hinck
 1991. *Sphyrnura euryceae* (Monogenea) on *Eurycea* spp. (Amphibia: Caudata), from northcentral Arkansas. J. Helminthol. Soc. Washington 58(1): 137–140, 4 figs.
- McConnaughey, B. H.
 1941. Two new Mesozoa from California, *Dicyemene californica* and *Dicyemene brevicephala* (Dicyemidae). J. Parasitol. 27(1): 63–69, 10 figs.
 1949a. *Dicyema sullivanii*, a new mesozoan from lower California. Ibid. 35(2): 122–124, 7 figs.
 1949b. Mesozoa of the family Dicyemidae from California. Univ. California Publ. Zool. 55(1): 1–33, 7 pls.
- Newman, L. J., and L. R. G. Cannon
 1994. *Pseudoceros* and *Pseudobiceros* (Platyhelminthes, Polycladida, Pseudocerotidae) from eastern Australia and Papua New Guinea. Mem. Queensland Mus. 37(1): 205–266, 51 figs.
- Nigrelli, R. F.
 1938. Parasites of the swordfish, *Xiphias gladius* Linnaeus. Am. Mus. Novitates 996: 16 pp., 5 figs.
- Nigrelli, R. F., and H. W. Stunkard
 1947. Studies on the genus *Hirudinella*, giant trematodes of scombriform fishes. Zoologica 31(4): 185–196, 26 figs.
- Odlaug, T. O.
 1938. *Zeugorchis longicirrus*, a new trematode from *Natrix sipedon*. Trans. Am. Microsc. Soc. 57(2): 173–177, 1 fig.
- Ogren, R. E.
 1987. Description of a new three-lined land planarian of the genus *Bipalium* (Turbellaria: Tricladida) from Pennsylvania, U.S.A. Trans. Am. Microsc. Soc. 106(1): 21–30, 8 figs.
 1989. Identification features of the two-lined land planarian *Rhynchodemus sylvaticus*, with evidence that *Rhynchodemus americanus* is conspecific. Ibid. 108(1): 40–44, 3 figs.
- Ogren, R. E., and J. T. Darlington
 1991. *Geoplana arkalabamensis* n. sp., a land planarian from the southern United States (Turbellaria: Tricladida: Geoplanidae). Trans. Am. Microsc. Soc. 110(3): 226–236, 14 figs.
- Ogren, R. E., and M. Kawakatsu
 1987. Index to the species of the genus *Bipalium* (Turbellaria, Tricladida, Terricola). Bull. Fuji Women's Coll. ser. II 25: 79–119.
 1988. Index to the species of the family Rhynchodemidae (Turbellaria, Tricladida, Terricola). Part I: Rhynchodeminae. Ibid. 26: 39–91.
 1990. Index to the species of the family Geoplanidae (Turbellaria, Tricladida, Terricola). Part I: Geoplaninae. Ibid. 28: 79–166.
 1991. Index to the species of the family Geoplanidae (Turbellaria, Tricladida, Terricola). Part II: Caenoplaninae and Pelmatoplaninae. Ibid. 29: 25–102.
- Olsson, P.
 1868. Entozoa, iakttagna hos Skandinaviska hafsfiskar. Lunds Universitets Årsskrift. Part II. Afdelningen for Matematik och Naturvetenskap 4(8): 64 pp., pls. 3–5.
 1876. Bidrag till Skandinaviens helminthfauna. I. Kongl. Svenska Vetenskaps-Akademiens Handlingar 14(1): 35 pp., 4 pls.
 1893. Bidrag till Skandinaviens helminthfauna. II. Ibid. 25(12): 41 pp., 5 pls.
- Overstreet, R. M., J. T. Self, and K. A. Vliet
 1985. The pentastomid *Sebekia mississippiensis* sp. n. in the American alligator and other hosts. Proc. Helminthol. Soc. Washington 52(2): 266–277, 27 figs.
- Paul, A. A.
 1935. *Polystoma integerrimum nearcticum* n.

- subsp. from the urinary bladder, genital ducts, kidneys and gills of *Hyla versicolor* Le Conte. J. Parasitol. 21(6): 442.
1938. Life history studies of North American fresh-water polystomes. Ibid. 24(6): 489-510, 3 pls.
- Poirier, M. J.
1909. Trématodes parasites de l'éléphant d'Afrique. C. R. 37^{me} Session Assoc. Franc. L'Avance. Sci. France: 580-582, 5 figs.
- Poulter, J. L.
1974. A new species of the genus *Pericelis*, a polyclad flatworm from Hawaii. In N. W. Riser and M. P. Morse (eds.), Biology of the Turbellaria. New York: McGraw-Hill. pp. 93-107, 10 figs.
1976. Hawaiian polyclad flatworms: Prosthlostomids. Pacific Sci. 29(4): 317-339, 22 figs.
- Price, E. W.
1931. Four new species of trematode worms from the muskrat, *Ondatra zibethica*, with a key to the trematode parasites of the muskrat. Proc. U.S. Natl. Mus. 79(4): 13 pp., 4 figs.
1932. On the genus *Phagicola* Faust, 1920. J. Parasitol. 19(1): 88-89.
1939. North American monogenetic trematodes. IV. The family Polystomatidae (Polystomatoidea). Proc. Helminthol. Soc. Washington 6(2): 80-92, 1 fig.
- Prudhoe, S.
1985. A Monograph on polyclad Turbellaria. Oxford: Br. Mus. (Nat. Hist.)/Oxford Univ. Press. 259 pp., 140 figs.
1989. Polyclad turbellarians recorded from African waters. Bull. Br. Mus. Nat. Hist. (Zool.) 55(1): 47-96, 45 figs.
- Remane, A.
1934. Die gastrotrichen des Küstengrundwassers von Schilksee. Schr. Naturwiss. Ver. Schleswig-Holstein 20(2): 473-478, 6 figs.
- Rieger, R. M.
1971a. *Bradynectes sterreri* gen. nov., spec. nov., eine neue psammobionte Macrostomide (Turbellaria). Zool. Jahrb. Abt. Syst. Oekol. Georg. Tiere 98(2): 205-235, 14 figs.
- 1971b. Die turbellarienfamilie Dolichomacrostomidae nov. fam. (Macrostomida). I. Teil Vorbemerkungen und Karlingiinae nov. subfam. 1. Ibid. 98(2): 236-314, 23 figs.
1974. A new class of Turbellaria-Typhloplanoida with a proboscis and its relationship to Kalyptorhynchia. In N. W. Riser and M. P. Morse (eds.), Biology of the Turbellaria. New York: McGraw-Hill. pp. 23-62, 17 figs.
1977. The relationship of character variability and morphological complexity in copulatory structures of Turbellaria-Macrostomida and -Haplopharyngida. Mikrofauna Meeresboden 61: 197-216, 11 figs.
1986. Asexual reproduction and the turbellarian archetype. Hydrobiologia 132: 32-45, 6 figs.
- Rieger, R. M., E. Ruppert, G. E. Rieger, and C. Schoepfer-Sterrer
1974. On the fine structure of *Chorododasys antennatus* sp. n. Zool. Scr. 3(5/6): 219-237, 11 figs.
- Rieger, R. M., and W. Sterrer
1975. New spicular skeletons in Turbellaria, and the occurrence of spicules in marine meiofauna. Z. Zool. Syst. Evolutionsforsch. 13(3): 207-248, 24 figs.
- Rieger, R. M., and S. Tyler
1974. A new glandular sensory organ in interstitial Macrostomida (Turbellaria) I. ultrastructure. Mikrofauna Meeresboden 42: 138-175, 15 figs.
- Riley, J., and J. T. Self
1979. On the systematics of the pentastomid genus *Porocephalus* (Humbolt, 1811) with descriptions of two new species. Syst. Parasitol. 1(1): 25-42, 18 figs.
1980. On the systematics and life-cycle of the pentastomid genus *Kiricephalus* Sambon, 1922 with descriptions of three new species. Ibid. 1(2): 127-140, 10 figs.
1981. A redescription of *Waddycephalus tertiusculus* (Baird, 1862) Sambon, 1922 and a revision of the taxonomy of the genus *Waddycephalus* (Sambon, 1922), pentastomid parasites of Asian, Australian and Indonesian snakes, with descriptions of eight new species. Ibid. 3(4): 243-257, 12 figs.
1982. A revision of the pentastomid genus *Parasambonia* Stunkard & Gandal, 1968: a new generic character, a description of the male and a new species. Ibid. 4(2): 125-133, 7 figs.
- Riser, N. W.
1948. *Amphilina bipunctata* n. sp., a North American cestodarian. J. Parasitol. 34(6.1): 479-485, 1 pl.
1981. New England Coelogynoporidae. Hydrobiologia 84: 139-145, 2 figs.
1987. *Nemertinoides elongatus* gen. n., sp. n. (Turbellaria: Nemertodermatida) from coarse sand beaches of the western North Atlantic. Proc. Helminthol. Soc. Washington 54(1): 60-67, 10 figs.

- Ruppert, E. E.
1979. Morphology and systematics of the Xenotrichulidae (Gastrotricha, Chaetontotida). *Mikrofauna Meeresboden* 76: 56 pp., 28 figs.
- Salley, E. J., J. R. Lichtenfels, and J. H. Shaw
1978. Checklist of types in the U.S. National Parasite Collection. Index-catalogue of medical and veterinary zoology. Spec. Publ. 4. Washington, D.C.: U.S. Gov. Printing Office. 233 pp.
- Schmidt, G. D.
1986. CRC handbook of tapeworm identification. Boca Raton, FL: CRC Press. 675 pp., 513 figs.
- Schrom, H.
1972. Nordadriatische gastrotrichen. *Helgoländer Wiss. Meeresunters.* 23(3): 286–351, 24 figs.
- Self, J. T., and A. A. Rego
1985. Reassessments and revisions of certain genera and species of the family Sebekidae (Pentastomida) including description of *Sebekia microhamus* n. sp. *Syst. Parasitol.* 7(1): 33–41, 5 figs.
- Sey, O.
1991. CRC handbook of amphistomes. Boca Raton, FL: CRC Press. 480 pp., 488 figs.
- Sinitzyn, D. F.
1906. (Distomes of fish and frogs in the vicinity of Warsaw.) [Russian.] *Varshavskbihiia Universitetskbihiia Izvhiest-bihiaa* 1: 97–207, 55(+) figs.
- Skrjabin, K. I.
1953. Trematody zhivotnykh i cheloveka. Vol. 8. Moscow: Izd. Akad. Nauk SSSR. 618 pp., 157 figs.
1955. Trematody zhivotnykh i cheloveka. Vol. 10. Moscow: Izd. Akad. Nauk SSSR. 653 pp., 175 figs.
1962. Trematody zhivotnykh i cheloveka. Vol. 20. Moscow: Izd. Akad. Nauk SSSR. 563 pp., 294 figs.
- Sluys, R.
1989. A monograph of the marine triclads. Rotterdam: Balkema. 463 pp., 322 figs., 12 pls.
1992. Synopsis of the freshwater triclads of the Caribbean (Platyhelminthes, Tricladida, Paludicola). *Stud. Nat. Hist. Caribbean Region* 71: 1–23, 8 figs.
- Smith, D. G.
1991. On some larger turbellarian worms (Platyhelminthes) living in temporary fresh waters of southern New England. *Hydrobiologia* 220(3): 255–265, 12 figs.
- Smith, E. H.
1960. On a new polyclad commensal of pro-sobranchs. *An. Acad. Brasileira Cienc.* 32(3/4): 385–390, 4 figs.
1961. A new commensal polyclad from Panama. *Veliger* 4(2): 69–70, pl. 16.
- Spurr, S. J.
1983. A new species of the genus *Paramalostomum* (Turbellaria: Macrostomida) from New England. *Trans. Am. Microsc. Soc.* 102(4): 333–348, 11 figs.
- Sterrer, W., and R. Rieger
1974. Retronectidae—a new cosmopolitan marine family of Catenulida (Turbellaria). In N. W. Riser and M. P. Morse (eds.), *Biology of the Turbellaria*. New York: McGraw-Hill. pp. 63–92, 15 figs.
- Stunkard, H. W.
1916. On the anatomy and relationships of some North American trematodes. *J. Parasitol.* 3(1): 21–27.
1917. Studies on North American Polystomidae, Aspidogastridae, and Paramphistomidae. *Illinois Biol. Monogr.* 3(3): 114 pp., 79 figs.
1922. Two new genera of North American blood flukes. *Am. Mus. Novitates* 39: 8 pp., 3 figs.
1923a. A new genus of trematodes from the eastern painted turtle. *Anat. Rec.* 24(6): 373–374.
1923b. Studies on North American blood flukes. *Bull. Am. Mus. Nat. Hist.* 48(7): 165–221, 61 figs.
1923c. Observations on the development of *Polystoma multifalx*, n. sp., from the pharynx of *Chrysemys floridiana* [sic]. *Anat. Rec.* 26(5): 357.
1923d. A new trematode from the snapping turtle, *Chelydra serpentina*. *Ibid.* 26(5): 358.
1924a. On some trematodes from Florida turtles. *Trans. Am. Microsc. Soc.* 43(2): 97–117, 11 figs.
1924b. A new trematode, *Oculotrema hippopotami* n. g., n. sp., from the eye of the hippopotamus. *Parasitology* 16(4): 436–440, pl. 16.
1925. A new blood fluke, *Unicaecum ruszkowskii* n. g., n. sp.; a contribution to the relationship of the blood-infesting trematodes. *Anat. Rec.* 31(4): 317.
1926. A new trematode, *Vasatrema amydae* n. g., n. sp., from the vascular system of the soft-shelled turtle, *Amyda*. *Ibid.* 34(3): 165.
1927. Sur *L'Unicaecum ruszkowskii*, trematode sanguicole des tortues d'eau douce de l'Amerique du Nord. *Ann. Parasitol. Hum. Comp.* 5(2): 117–126, 3 figs.

1928. Observations nouvelles sur les trématodes sanguicoles du genre *Vasotrema* (Spirorchidae) avec description des deux espèces nouvelles. *Ibid.* 6(3): 303–320, 9 figs.
- 1929a. The parasitic worms collected by the American Museum of Natural History Expedition to the Belgian Congo, 1909–1914. *Bull. Am. Mus. Nat. Hist.* 58(6): 233–289, 37 figs.
- 1929b. Further observations on the cercaria which occurs in the marine annelid, *Hydroides dianthus*. *J. Parasitol.* 16(2): 106.
1931. Further observations on the occurrence of anal openings in digenetic trematodes. *Z. Parasitenk.* 3(4): 713–725, 7 figs.
1932. Some larval trematodes from the coast in the region of Roscoff, Finistère. *Parasitology* 24(3): 321–343, 13 figs.
1933. *Plagitura parva* n. sp. from *Triturus viridescens*. *J. Parasitol.* 20(2): 134.
1935. A new trematode, *Probolitrema californiense*, from the coelom of the sting ray, *Myliobatus californicus*. *Ibid.* 21(5): 359–364, 1 fig.
1936. The morphology and life history of *Plagitura parva* Stunkard, 1933, (Trematoda). *Ibid.* 22(4): 354–374, 11 figs.
- 1938a. Parasitic flatworms from Yucatan. *Carnegie Inst. Washington Publ.* 491: 33–50, 8 figs.
- 1938b. *Oochoristica parvula* n. nom. for *Oochoristica parva* Stunkard, 1938, preoccupied. *J. Parasitol.* 24(6): 554.
1943. A new trematode, *Dictyangium chelydrae* (Microscaphiidae = Angiodictyidae), from the snapping turtle, *Chelydra serpentina*. *Ibid.* 29(2): 143–150, 6 figs.
1951. Observations on the morphology and life-history of *Microphallus limuli* n. sp. (Trematoda: Microphallidae). *Biol. Bull. (Woods Hole)* 101(3): 307–318, 4 figs.
1959. The morphology and life-history of the digenetic trematode, *Asymphylogora amnicolae* n. sp.; the possible significance of progenesis for the phylogeny of the digenea. *Ibid.* 117(3): 562–581, 7 figs.
1960. Further studies on the trematode genus *Himasthla* with descriptions of *H. mcintoshi* n. sp., *H. piscicola* n. sp., and stages in the life-history of *H. compacta* n. sp. *Ibid.* 119(3): 529–549, 24 figs.
1962. *Caballerocotyla klawei* sp. n., a monogenetic trematode from the nasal capsule of *Neothunnus macropterus*. *J. Parasitol.* 48(6): 883–890, 7 figs.
- 1965a. A digenetic trematode, *Botulus cablei*, n. sp., from the stomach of the lancetfish, *Alepisaurus borealis* Gill, taken in the South Pacific. *Biol. Bull. (Woods Hole)* 128(3): 488–492, 4 figs.
- 1965b. *Paratriotaenia oedipomidatis* gen. et sp. n. (Cestoda), from a marmoset. *J. Parasitol.* 51(4): 545–551, 5 figs.
- 1965c. Type species of *Paratriotaenia*. *Ibid.* 51(6): 996.
- 1966a. The morphology and life history of the digenetic trematode, *Himasthla littorinae* sp. n. (Echinostomatidae). *Ibid.* 52(2): 367–372, 4 figs.
- 1966b. The morphology and life-history of *Notocotylus atlanticus* n. sp., a digenetic trematode of eider ducks, *Somateria mollissima*, and the designation, *Notocotylus duboisi* nom. nov., for *Notocotylus imbricatus* (Looss, 1893) Szidat, 1935. *Biol. Bull. (Woods Hole)* 131(3): 501–515, 8 figs.
1967. The morphology, life-history, and systematic relations of the digenetic trematode, *Uniserialis breviserialis* sp. nov., (Notocotylidae), a parasite of the bursa Fabricius of birds. *Ibid.* 132(2): 266–276, 7 figs.
1971. Renicolid trematodes (Digenea) from the renal tubules of birds. *Ann. Parasitol.* 46(1): 109–118.
1979. *Auritelorchis* nom. nov. for *Paratelorchis* Stunkard & Franz, 1977, preoccupied. *Trans. Am. Microsc. Soc.* 98(1): 142.
1980. Successive hosts and developmental stages in the life history of *Neopechona cablei* sp. n. (Trematoda: Lepocreadiidae). *J. Parasitol.* 66(4): 636–641, 10 figs.
- Stunkard, H. W., and C. H. Alvey
1929. A new liver fluke, *Zalophotrema hepaticum*, from the California sea lion, *Zalophus californianus*. *J. Parasitol.* 16(2): 106–107.
1930. The morphology of *Zalophotrema hepaticum*, with a review of the trematode family Fasciolidae. *Parasitology* 22(3): 326–333, pl. 33.
- Stunkard, H. W., and F. W. Dunihue
1931. Notes on trematodes from a Long Island duck with description of a new species. *Biol. Bull. (Woods Hole)* 60(2): 179–186, 4 figs.
- Stunkard, H. W., and R. Franz
1977. *Paratelorchis dollfusi* n. g., n. sp., a digenetic trematode from the striped swamp-snake, *Regina alleni*: systematic

- and taxonomic considerations. Trans. Am. Microsc. Soc. 96(3): 383–389, 1 fig.
- Stunkard, H. W., and C. P. Gandal
1968. The pentastomes, *Waddycephalus teretiusculus* (Baird, 1862) Sambon, 1922 and *Parasambonia bridgesi* n. gen., n. sp., from the lungs of the Australian snake, *Pseudechis porphyriacus*. Zoolologica 53(2): 49–56, 6 figs.
- Stunkard, H. W., and C. B. Haviland
1924. Trematodes from the rat. Am. Mus. Novitates 126: 10 pp., 5 figs.
- Stunkard, H. W., and W. F. Lynch
1944. A new anoplocephaline cestode, *Oochoristica anniellae*, from the California limbless lizard. Trans. Am. Microsc. Soc. 63(2): 165–169, 1 pl.
- Stunkard, H. W., and R. F. Nigrelli
1934. Observations on the genus *Sterrhurus* Looss, with a description of *Sterrhurus branchialis* sp. nov. (Trematoda, Hem- iuridae). Biol. Bull. (Woods Hole) 67(3): 534–543, 2 figs.
- Stunkard, H. W., and J. R. Uzmann
1955. The killifish *Fundulus heteroclitus*, second intermediate host of the trematode, *Ascocotyle (Phagicola) diminuta*. Biol. Bull. (Woods Hole) 109(3): 475–483, 4 figs.
- Van Cleave, H. J.
1921. Notes on two genera of ectoparasitic trematodes from fresh-water fishes. J. Parasitol. 8(1): 33–39, 2 figs.
- Wardle, R. A., and J. A. McLeod
1952. The zoology of tapeworms. Minneapolis: Univ. of Minnesota Press. 780 pp., 419 figs.
- Wilke, U.
1954. Mediterranean gastrotrichen. Zool. Jahrb. Abt. Syst. Oekol. Geogr. Tiere 82(6): 497–550, 29 figs.
- Yamaguti, S.
1968. Monogenetic trematodes of Hawaiian fishes. Honolulu: Univ. of Hawaii Press. 287 pp., 147 figs.
1971. Synopsis of digenetic trematodes of vertebrates. 2 vols. Tokyo: Keigaku Publishing Co. 1074 pp., 349 pls.

INDEX, BY PHYLUM AND CLASS

PLATYHELMINTHES

TURBELLARIA

Acanthiella, 4
Acanthozoon, 4
Acerotisa, 4, 18
adventitium, *Bipalium*, 5
Afronta, 4
alba, *Acerotisa*, 18
albopapillosus, *Acanthozoon*, 4
album, *Plagiostomum*, 20
albus, *Oligocladus*, 18
Alleena, 4
Alloiplana, 26
Amaga, 10
americanus, *Rhynchodemus*, 23
Amphiscolops, 5
andina, *Geoplana*, 10
andrewsi, *Mesostoma*, 14
angustus, *Desmorhynchus*, 7
Anoplodium, 5
aphalla, *Geoplana*, 10
appia, *Mesostoma*, 14
appinum, *Mesostoma*, 14
arctica, *Mesostoma*, 14
arcticum, *Mesostoma*, 14
arcticum, *Nesion*, 16
arimana, *Dugesia*, 8
arkalabamensis, *Geoplana*, 10

Armatoplana, 25
asymmetrica, *Monosolenia*, 15
atypica, *Retronektes*, 22
aurantiaca, *Afronta*, 4
Australopacifica, 10
bajae, *Pseudoceros*, 21
bathycola, *Bertiliella*, 5
bellostriatus, *Prostheceraeus*, 21
bermudensis, *Amphiscolops*, 5
bermudensis, *Paramyozonaria*, 18
Bertiliella, 5
bichaeta, *Myomacrostromum*, 16
bimarginatus, *Pseudoceros*, 22
Bipalium, 5
bipolaris, *Florianella*, 9
Bothromesostoma, 15
Bradynectes, 5
brasiliensis, *Latocestus*, 13
buchanani, *Speophila*, 23
burchami, *Planocera*, 3, 20
Caenoplana, 10
californica, *Acerotisa*, 4
californica, *Eurylepta*, 9
californica, *Hoploplana*, 11
californicum, *Mesostoma*, 14
californicum, *Thysanozoon*, 26
californicus, *Stylochus*, 25
canadensis, *Pseudoceros*, 22
canai, *Girardia*, 10

carolinensis, *Dalyella viridis*, 7
cavernicola, *Phagocata*, 19
celeris, *Notoplana*, 17
celeris, *Pucelis*, 17
chaetonotoides, *Acanthiella*, 4
chloranota, *Stylochoplana*, 17
Cirrifera, 6
clio, *Retronektes*, 22
Coelogygnopora, 6
coerulea, *Caenoplana*, 10
Comprostatum, 6
contamanensis, *Geoplana*, 10
corallophilus, *Pseudoceros*, 22
coronum, *Paromalostomum*, 18
costaricense, *Bipalium*, 5
costaricensis, *Bipalium*, 5
Crassicollum, 6
crassula, *Myopea*, 16
Cryptobioceros, 21, 22
Cryptocelis, 6
curvipenis, *Mesostoma*, 14
Cystiplana, 7
Dalyella, 7
delmaris, *Macginitiella*, 13
Dendrocoelopsis, 7
Desmorhynchus, 7
diabolis, *Dugesia*, 8
Diplandros, 7
Diporodemus, 7, 8

- Discocelis*, 8
dorotocephala, *Girardia*, 8, 9
Dugesia, 8
durhami, *Monocelis*, 15
Ectocotyla, 8
ehrenbergi, *Mesostoma*, 14
elongatus, *Nemertinoidea*, 16
Enchiridium, 8
Endeavouria, 10
Endocincta, 8, 9
epichitonius, *Nexilis*, 16
erato, *Paracatenula*, 18
erotica, *Coelogynopora*, 6
ernesti, *Marcusia*, 14
Eukytorhynchus, 12
Eulatocestus, 13
Euplanaria, 9
Eurylepta, 4, 9, 18
Euryleptodes, 9
festai, *Girardia*, 8
flavotuberculatum, *Thysanozoon*, 26
Florianella, 9
floridana, *Gnesioceros*, 11
franciscanus, *Stylocheilus*, 25
Freemania, 17, 18
"funilia", *Solenofilomorpha*, 23
funilis, *Solenofilomorpha*, 23
Fusantrum, 9
galapagensis, *Latocestus*, 13
Geocentrophora, 9
Geoplana, 10
georgiana, *Sphalloplana*, 24
Girardia, 8–10
"glandularis", *Macrocotyla*, 14
glandulosa, *Macrocotyla*, 13
glandulosa, *Syndemis*, 25
Gnesioceros, 11
gracilis, *Phagocata*, 19
griseus, *Pseudoceros*, 22
guaymensis, *Solenofilomorpha*, 23
hewatti, *Indistylus*, 11
hirudo, *Ectocotyla*, 8
hoffmasteri, *Speophila*, 23
Hoploplana, 11
hubrichti, *Speophila*, 23
hymanae, *Coelogynopora*, 6
hymanae, *Pericelis*, 19
Imogene, 11, 25
indigenus, *Diporodemus*, 7
Indistylus, 11
inquieta, *Leptoplanea*, 13
inquieta, *Notoplanea*, 17
inquilina, *Notoplanea*, 17
inquilina, *Stylocheilus*, 24
insularis, *Comprostatum*, 6
insularis, *Discocelis*, 8
insularis, *Euryleptodes*, 9
jenneri, *Myozonaria*, 16
juxtaforcipis, *Coelogynopora*, 6
kansensis, *Sphalloplana*, 24
karlingi, *Kytorhynchella*, 12
katoi, *Prosthiostomum* (*Lurymare*), 21
Kenkia, 11, 12, 14
kewense, *Bipalium*, 5
Koinostylus, 20, 22
Kytorhynchella, 12
Kytorhynchoides, 3, 12
Kytorhynchus, 12
laevis, *Mexistylus*, 15
lata, *Gnesioceros sargassicola*, 11
latafaucium, *Myopea*, 16
latocelis, *Prosthiostomum*, 21
Latocestus, 5, 13
Latoplanea, 15
leoparda, *Eurylepta*, 9
Leptoplanea, 13
Leptostylus, 13
levis, *Latoplanea*, 15
limnoriae, *Leptoplanea*, 13
litoricola, *Freemania*, 18
litoricola, *Pucelis*, 18
longastyletta, *Stylocheilus*, 24
longiductum, *Anoplodium*, 5
longipenis, *Stylocheilus*, 24
Longiprostatum, 13
luracola, *Hoploplana*, 11
Lurymare, 21
Macginitella, 13
Macroatrium, 13
Macrocotyla, 13, 14
macropenis, *Mesostoma*, 14
macroprostatum, *Mesostoma*, 15
macrostylus, *Kytorhynchus*, 12
magnum, *Mesostoma*, 15
Marcusia, 14
meixneri, *Kytorhynchella*, 12
Mesostoma, 14, 15
mexicana, *Alleena*, 4
mexicana, *Ommatoplanea*, 15
mexicana, *Phaenocelis*, 19
mexicana, *Pucelis*, 19
mexicanus, *Pseudoceros*, 22
Mexistylus, 15
microbursalis, *Euplanaria*, 9
Microplana, 15
microstylus, *Kytorhynchus*, 12
mohri, *Sphalloplana*, 24
Monocelis, 15
monopharyngea, *Phagocata gracilis*, 19
Monosolenia, 15
montiporae, *Prosthiostomum* (*Prosthiostomum*), 21
morgani, *Phagocata morgani*, 19
multicelis, *Acerotisa*, 4
multicelis, *Prosthiostomum*, 21
multitesticulata, *Ectocotyla*, 8
musculare, *Crassicollum*, 6
Myomacrostromum, 16
Myopea, 16
Myozonaria, 16
nahantensis, *Proschizorhynchus*, 20
natans, *Notoplanea*, 17
natans, *Pucelis*, 17
Nemertinoidea, 16
Nesion, 16
Nexilis, 16, 17
Notogynaphallia, 10
Notocomplana, 17–19
Notoplanea, 3, 17, 24
novacambrensis, *Leptostylus*, 13
oahuensis, *Rhynchodemus*, 23
occidentalis, *Cryptocelis*, 6
oculatus, *Kytorhynchus*, 12
oculifera, *Imogene*, 11
Oligocladus, 18
Ommatoplanea, 15
oregonensis, *Phagocata*, 19
ostreophagus, *Pseudostylus*, 22
pacifica, *Procerodes*, 20
paguri, *Ectocotyla*, 8
panamensis, *Geoplana*, 10
papillatus, *Proschizorhynchus*, 20
Paracatenula, 18
Paramyozonaria, 18
Parathylacorhynchus, 18
Paromalostomum, 18
Pasipha, 10
pennsylvanicum, *Bipalium*, 5
Pericelis, 14, 19
Phaenocelis, 6, 19
Phaenoplanea, 25
Phagocata, 19
philadelphica, *Euplanaria*, 9
Plagiostomum, 20
Planocera, 3, 20
platygasticum, *Mesostoma*, 14
Pleioplanea, 13
plenus, *Diporodemus*, 7
poaceaglandis, *Coelogynopora*, 6
Polycelis, 20
Polypharyngea, 24
Praestheceraeus, 21
pricei, *Speophila*, 23
Probursa, 20
Procerodes, 20
Proschizorhynchus, 20
Prostheceraeus, 21
Prosthiostomum, 21
Pseudobiceros, 21, 22
Pseudoceros, 21, 22
Pseudogeoplana, 10
Pseudostylus, 22
Pucelis, 17–19
punctata, *Endocincta*, 8

punctatum, *Enchiridium*, 8
purpurea, *Phaenocelis*, 6
ramosum, *Anoplodium*, 5
reishi, *Stylochoplana*, 25
Retronectes, 22, 23
rhammiphorum, *Fusantrum*, 9
rhynchida, *Kenkia*, 11
Rhynchodemus, 7, 23
rickettsi, *Longiprostatum*, 13
riedi, *Parathylacorhynchus*, 18
riedli, *Kytorkhynchella*, 12
rubra, *Cystiplana*, 7
rufocephalata, *Microplana*, 15
sanchezi, *Dugesia*, 8
sanguinea, *Notoplana*, 17
sanguinea, *Pucelis*, 17
sanjuaniana, *Notoplana*, 17
sanjuaniana, *Pucelis*, 17
sargassicola, *Gnesioceros*, 11
segnis, *Notoplana*, 3, 17, 19
septemlineata, *Geoplana*, 10
setosum, *Macroatrium*, 13
simplex, *Paramyozonaria*, 18
singularis, *Diplandros*, 7
Solenofilomorpha, 23
Speophila, 23, 24
Sphalloplana, 23, 24
sterreri, *Bradynectes*, 5
stylifera, *Zyganotroplana*, 26
Stylochocestus, 11
Stylochoplana, 13, 17, 24, 25
Stylochus, 25
Stylostomum, 4
subpallida, *Geoplana*, 10
subterranea, *Phagocata*, 19
sylvaticus, *Rhynchodemus*, 23
Syndesmis, 25
Syndisyrix, 25
texanus, *Pseudoceros*, 22
Thalamoplana, 8
thalia, *Retronectes*, 22
Thysanozoon, 4, 26
tibetica, *Polycelis*, 20
titicacana, *Dugesia*, 8
tripartitus, *Stylochus*, 25
Triplana, 25
tropica, *Geocentrophora*, 9
tuberculata, *Ommatoplana*, 15
tuberculatus, *Mexistylolochus*, 15
ucayalensis, *Geoplana*, 10
unichaeta, *Myomacrostromum*, 16
urania, *Paracatenula*, 18
usaguia, *Hoploplana*, 11
vaga, *Geoplana*, 10
vaginatus, *Dendrocoelopsis*, 7
velata, *Phagocata*, 19
veneris, *Probursa*, 20
"venus", *Comprostatum*, 6
vernale, *Mesostoma*, 15

verrilli, *Gnesioceros*, 11
virginiana, *Sphalloplana*, 24
viridis, *Dalyella*, 7
viridis, *Stylochoplana*, 25
woodworthi, *Phagocata*, 19
xanthoderma, *Cirrifera*, 6
yucatanii, *Diporodemus*, 7
zebra, *Prostheceraeus*, 21
Zyganotroplana, 26

MONOGENEA

Caballerocotyla, 26
coronatum, *Polystomoides*, 27
Diclidophora, 26
fairporti, *Gyrodactylus*, 26
floridanum, *Polystoma*, 26
Gyrodactylus, 26
hippopotami, *Oculotrema*, 26
klawei, *Caballerocotyla*, 26
minor, *Diclidophora*, 26
minus, *Octobothrium*, 26
multifalx, *Polystoma*, 27
Nasicola, 26
nearcticum, *Polystoma integerri-*
mum, 26
Neopolystoma, 26, 27
Octobothrium, 26
Oculotrema, 26
oligorchis, *Sphyrnura*, 27
opacum, *Polystoma*, 27
orbiculare, *Neopolystoma*, 26
orbiculare, *Polystoma*, 27
oris, *Polystomoides*, 27
osleri, *Sphyrnura*, 27
polyorchis, *Sphyrnura*, 27
Polystoma, 26, 27
Polystomoides, 27
Sphyrnura, 27

TREMATODA

Acanthochasmus, 28
Acanthostomum, 28
Allassostoma, 28
Allassostomoides, 28
Allassostoma, 28
Allassostomoides, 28
amnicolae, *Asymphyllodora*, 29
Amphistomum, 28
amphiumae, *Cephalogonimus*, 30
amydae, *Vasatrema*, 35, 36
Anenterotrema, 29
Ascocotyle, 29
Asymphyllodora, 29
atlanticus, *Notocotylus*, 33
attenuatum, *Vasotrema*, 35
attenuatus, *Pterygotomaschalos*,
 34
Auridistomum, 34

Auritellorchis, 33
auritum, *Anenterotrema*, 29
beebei, *Hirudinella*, 32
bicaudata, *Brumptia*, 29, 30
bicaudatum, *Amphistomum*, 28
bilobatum, *Lobatozoum*, 32
Blairium, 28
Botulus, 29
branchialis, *Sterrhurus*, 35
brevicaecum, *Mesocoelium*, 32
breviserialis, *Uniserialis*, 35
Brumptia, 29, 30
cablei, *Botulus*, 29
cablei, *Neopechona*, 32
californiense, *Probolitrema*, 34
Catoptroides, 33
Cephalogonimus, 30
ceratosa, *Zygocotyle*, 36
Cercaria, 30
chelydrae, *Auridistomum*, 34
chelydrae, *Dictyangium*, 30
Cladorchis, 30
compacta, *Himasthla*, 31
compactus, *Cephalogonimus*, 30
compactus, *Stephanopharynx*, 35
copulans, *Liolope*, 32
coronarium, *Proctocaeum*, 28
Cyclocoelum, 30
cygnoides, *Gorgodera*, 30
Dasymetra, 36
Dictyangium, 30, 31
diminuta, *Ascocotyle (Parascoco-*
tyle), 29
diminuta, *Phagicola*, 29
diploporus, *Acanthochasmus*, 28
Distoma, 31
dollfusi, *Paratelorchis*, 33
Echinochasmus, 31
elegans, *Spirorchis*, 34
enterocolpium, *Phyllodistomum*, 33
gigas, *Cladorchis*, 30
Gorgodera, 30
gracilis, *Hapalorhynchus*, 31
haematobium, *Henotosoma*, 31
Haematotrema, 31
Hapalorhynchus, 31
Henotosoma, 31
hepaticum, *Zalophotrema*, 36
Heterochinostomum, 31
Himasthla, 31, 32
Hirudinella, 32
japonicum, *Mesocoelium*, 32
Lecithochirium, 35
limuli, *Microphallus*, 32
Liolope, 32
littorinae, *Himasthla*, 32
Lobatozoum, 32
longicirrus, *Zeugorchis*, 36
loossi, *Cercaria*, 30

loossi, *Gorgodera*, 30
lunata, *Zygocotyle*, 36
magnovatum, *Heterochinostomum*, 31
magnovatus, *Echinochasmus*, 31
magnum, *Alassostoma*, 28
Mesocoelium, 32
Microphallus, 32
microporus, *Botulus*, 29
minimum, *Acanthostomum*, 28
Neopechona, 32
Notocotylus, 33, 35
Opechona, 33
Parabascoides, 33
Paramonostomum, 33
Parascotyle, 29
Paratelorchis, 33
parva, *Plagitura*, 33
parvum, *Alassostoma*, 28
parvum, *Haematotremas*, 31
parvum, *Paramonostomum*, 33
parvus, *Alassostomoides*, 28
parvus, *Spirorchis*, 31
patellare, *Phyllodistomum* (*Ca-*
toptroides), 33
pearsei, *Mesocoelium*, 32
Phagicola, 29
phasidi, *Cyclocoelum*, 30
Phyllodistomum, 33
picta, *Spirorchis*, 34
Plagitura, 33
Probolitremas, 34
Proctocaecum, 28
pseudoechinatum, *Distoma*, 31
Pterygotomaschalos, 34
Renicola, 30
robustum, *Vasotremas*, 35
roscovita, *Cercaria*, 30
ruszkowskii, *Unicaecum*, 35
scripta, *Spirorchis*, 34
secundus, *Stephanopharynx*, 28,
 34
singulare, *Anenterotremas*, 29
Spirorchis, 31, 34
Stephanopharynx, 28, 34, 35
Stephanoprora, 31
Sterrhurus, 35
Stunkardiella, 28
Telorchis, 33
Unicaecum, 35

Uniserialis, 35
Vasatremas, 35, 36
Vasotremas, 35
ventricosa, *Hirudinella*, 32
Wardius, 36
yucatanensis, *Parabascoides*, 33
Zalaphotremas, 36
Zeugorchis, 36
zibethicus, *Wardius*, 36
Zygocotyle, 36

CESTODA

Amphilina, 36
annellae, *Oochoristica*, 36
Aporhynchus, 37
bipunctata, *Amphilina*, 36
crassus, *Triaenophorus*, 37
japonica, *Amphilina*, 36
lamontaeae, *Nybelinia*, 36
norvegicum, *Tetrabothrium*, 37
norvegicus, *Aporhynchus*, 37
Nybelinia, 36
oedipomidatis, *Paratriotaenia*, 37
Oochoristica, 36, 37
Paratriotaenia, 37
parva, *Oochoristica*, 37
parvula, *Oochoristica*, 37
robustus, *Triaenophorus*, 37
Tetrabothrium, 37
Triaenophorus, 37

RHOMBOZOA

abelis, *Dicyemenea*, 38
adminicula, *Conocyema*, 37, 38
apollyoni, *Dicyema*, 38
balamuthi, *Dicyema*, 37
brevicephala, *Dicyemenea*, 38
californica, *Dicyemenea*, 38, 39
Conocyema, 37, 38
Dicyema, 37, 38
Dicyemenea, 38, 39
granularis, *Dicyemenea*, 38
sullivani, *Dicyema*, 38

PENTASTOMIDA

PENTASTOMATA

aegypti, *Raillietiella*, 40
basiliscus, *Porocephalus*, 39

bridgesi, *Parasambonia*, 39
bufonis, *Raillietiella*, 40
calligaster, *Waddycephalus*, 41
cartagensis, *Raillietiella*, 40
clelii, *Kiricephalus*, 39
crotali, *Raillietiella*, 40
gabonensis, *Kiricephalus*, 39
Kiricephalus, 39
komodoensis, *Waddycephalus*, 41
longicauda, *Waddycephalus*, 41
microhamus, *Sebekia*, 40
minor, *Parasambonia*, 39
mississippiensis, *Sebekia*, 40
monarchus, *Raillietiella*, 40
Parasambonia, 39
Porocephalus, 39
porphyriacus, *Waddycephalus*, 41
punctulatus, *Waddycephalus*, 42
Raillietiella, 40
scincoides, *Raillietiella*, 40
scutata, *Waddycephalus*, 42
Sebekia, 40
superbus, *Waddycephalus*, 42
tortugensis, *Porocephalus*, 39
Waddycephalus, 41, 42

GASTROTRICHA

CHAETONOTOIDEA

affinis, *Xenotrichula*, 43
antennatus, *Chorododasys*, 42
arcassonensis, *Heteroxenotrichu-*
la, 42
carolinensis, *Xenotrichula*, 43
Chorododasys, 42
Heteroxenotrichula, 42, 43
intermedia, *Xenotrichula*, 43
lineata, *Xenotrichula*, 43
pygmaea, *Xenotrichula*, 43
squamosa, *Heteroxenotrichula*, 42
subterranea, *Xenotrichula*, 43
transatlantica, *Heteroxenotrichu-*
la, 42
wilkei, *Heteroxenotrichula*, 43
Xenodasys, 42
Xenotrichula, 43

Recent issues of the *Novitates* may be purchased from the Museum. Lists of back issues of the *Novitates*, *Bulletin*, and *Anthropological Papers* published during the last five years are available free of charge. Address orders to: American Museum of Natural History Library, Department D, Central Park West at 79th St., New York, N.Y. 10024. TEL: (212) 769-5545. FAX: (212) 769-5009. E-MAIL: scipubs@amnh.org