

2942 -- 3646

BIRDS

Faradje

July 8. 1911.

2942. { Purple Heron. ♂. t. n. e.  
Ardea p. purpurea. Hirundo griseopilea  
July 9. Swallow. ♀. o.n.e. A flock of these diminutive  
2943. s. 2572. swallows, numbering perhaps 20, was feeding  
low over a papyrus swamp.

July 10.

2944. Erithacus gubernator  
Small Shrike with rufous back and light gray crown.  
♂ Stomach contents: a green mantis.  
2945. Young of preceding. ♂ immature (in barred plumage)  
Stomach contents: insects.

Were sitting on the small trees about an abandoned  
village site.

2946. A dull gray-brown Finch with whitish superciliary  
line & streaks on crown. ♂. t. e. Bill pale buff,  
*Polioptila gallica elongans*  
iris rather dark brown, feet light brownish.

Was sitting perched amid brush, giving a  
low, but sweet and varied song. One or two had  
been seen a few days previously near Faradje,  
and a few more were later noticed about Aba.

2947. *Phyllastrephus senegalensis orientalis*  
A green & gray Bulbul with slightly rufous tail. ♀.  
o.n.e. Stomach contained a few insects and some  
small seeds as from fruit.

Shot in a patch of forest about midway between  
Faradje and Aba, where it seemed to be common.

In

Inches



Aba  
July 11.

It is a bird of the underbrush, resembling No 2871 in habits and notes, tho the latter are less musical.

2948. Goatsucker (*macrodiphteryx vexillarius*) ♂ teats slightly enlarged
2949. " " " ♂ teats very slightly enlarged.

Between 2 o'clock and daybreak this morning, with the aid of a bright moon, I counted 28 goatsuckers, all of the present species, males predominating. We were marching from Mordjan (Maripili) to Djaberet, <sup>(or Djalini)</sup> on the road to Aba, and the majority of the nightjars were frightened from the ground by our approach. In the early evening, on the contrary, they fly about actively, but I have not seen them sit on the ground.

(At Aba a few days later, Alimasi put up a long plumed male in the afternoon, and it lit on a bough of a small tree.)

Had one taken the same walk in February he would undoubtedly have found the route occupied by *m. macrodiphteryx* and *Scotornis*. Of the two species of the interesting genus *macodipteryx*, *m. vexillarius*,

Aba

July 11. 1911.

of which the males here acquire new plumes in June and July, might be called the rainy season member; and *M. macrodipterus*, which is most active in January and February, its racquet-tipped primaries molting in during November and December, is in a way a bird of the dry season, tho I do not wish to suggest that either of them migrates. <sup>Probably some</sup> ~~whether any~~ <sup>also</sup> such difference <sup>exists</sup> exists in the time of breeding ~~I am in doubt.~~ ~~for~~ *M. macrodipterus* lays in March and April, ~~and the young of~~ *M. vexillarius* has been discovered in May. <sup>nearly</sup> Both species are silent.

*S. cotorris* is much more noisy and conspicuous in the dry season; its loud clicking voice, often slowing down to a "k-k-k-k-k-k-k-k-k-k---" and even changing to a low quavering note, is heard continually at least from October till April or May, but not at present tho the birds are still occasionally seen. Eggs have been found from March to June.

(In April 1912, we heard *S. cotorris* calling in a papaya swamp at mid-day in cloudy weather. They stopped calling entirely in late May.)

Aba

July 12, 1911.

2950. Black Widow bird. *Colius pectoralis concolor*. ♂. testes slightly enlarged. Stomach contents: small seeds. (molting into nuptial plumage)

For the past three weeks we have been seeing molting individuals of this species. Some have already acquired complete black plumage. They seem to drop all the old rectrices at once.

2951. *Piparia rufifrons*. Dusky Swallow with buff throat. ♂. testes much enlarged. Iris dark brown; bill black; feet brownish.

One of a pair of swallows alighting in a small niche in a high vertical cliff just above a horizontal crack in the rock where there were two cup-shaped nests of mud. One of the nests was taken down the following day, and found to be old and empty; the other looked newer but could not be reached.

July 13.

*Indicator indicator*

2952. Honey Guide. ♂. testes slightly enlarged. Shot in trees on top of a rocky hill near village of Ibu.

2953. *Hirundo pusilla*. Streak-breasted Swallow ♂. t.m.e. Iris very dark brown, bill and feet blackish.

2954. Same as preceding. ♂. t.m.e.; skull not quite completely ossified.

2 w

Aba  
July 13, 1911.

A half dozen pairs or more were flying about the rocky hills near Aba village, but tho we found many signs of old nests beneath rocky ledges, no occupied nests were located. Perhaps it is not the season, for it was in June (the dry season there) that we saw the nest in Boma. To date specimens were secured from a party of 6 or 8 perching on the dry twigs of a small tree jutting out of a steep wall of rock. Tho never collected till now, this swallow has been observed several times since we left Boma, notably at Kinshasa, Mala, and Dunge.

- Piparia rufipectus*  
2955. Dusky Swallow with buff throat. ♂ testis slightly enlarged.  
2956. Mate of 2955. ovary slightly enlarged

This is a true rock swallow, found only about the hills. Its flight is rather slow, the birds are silent, and often alight on the rocks, often in very steep places.

2957. A small thrush (*Ruticilla*) with rufous tail. ♂ testes slightly enlarged. Iris dark brown, bill and feet black. Stomach contents: insects.

Bears a strong resemblance to the female

100

Aba

July 13, 1911.

of the European Redstart, which visits the rule in the dry season. This specimen was perched on a stump in cultivated land close to the foot of the hills.

2958. *Chamaezidae sp.* Thrush, black, with white wing coverts and rufous rump and belly. ♂. t. n. e. Iris dark brown, bill & feet black. Stomach contents: fruit and pieces of a couple of small insects.

A bird of the hills, alights on rocks and trees alike, and sings continually. Its posture is often very horizontal, it stands high on its legs, slowly raising and lowering the tail. Its song is sweet and varied, and tho' loud has perhaps something finch-like about it, but recalls also the songs of the black thrushes with white wing-coverts (such as Nos 2976 +

For a bird of its size it is wary.

2959. Small brown Lark with streaked breast. ♂ testes *Helicorhynchus modestus* very small; shell not quite completely ossified. Stomach contents: seeds and remains of very small insects.

Four of these larks were feeding on the stones of a bare, low hill; and when pursued alighted

Aba

July 13. 1911.

on top of stumps in a nearby plantation.

In arising from the earth they sometimes utter a faint call note.

July 14.

2960. White-crested Plantain-eater (*Turacus*). ♂. t. n. e.  
*Leucophrys*

Stomach contents: fruit. Very common at Aba.

Several times tos. I heard the call of the red-crested Musophaga coming up from the woods in the valleys.

2961. Red & black Weaver (*Pyromelaena*) <sup>frammiceps</sup> ♂. testes slightly enlarged (molting into nuptial plumage) A common bird throughout the whole region. First noticed molting about June 20 (near niangara); by July 4 (at Paradje) some were in full plumage.

2962. <sup>Alouesa</sup> *merinus auxilis*. A small brown Fly catcher. ♀ immature. Iris dark brown, bill dusky with base of mandible pale buff, feet dusky. Sitting in a tree on top of a rocky eminence near village of Ibu.

2963. Gray Goatsucker. ♂. t. n. e. Iris dark brown, *Caprimulgus trimaculatus tristigma*. rim of eyelids buff, bill brown with blackish tip. feet dark brownish.

This is the goatsucker whose low weird voice is heard on the hills at dusk. It utters a sort of whistled { "coo-hoo" or "coo-hoo-hoo"  
"whow-whow" "whow-whow-whow"  
and repeats it again and again. Less often

2 m/

P 430  
Aba

July 14, 1911.

there are four syllables. During the day they rest in sheltered positions on stones. This morning was cloudy, and two were heard calling, flushed and one of them shot.

2964. <sup>Dendrocoptes pileatus</sup> Greenish Woodpecker with large red crown patch and spotted breast. ♂. testes slightly enlarged. Iris dark red; bill bluish gray with dusky culmen and tip; feet greenish. Stomach contents: some white worm-like insect larvae and pieces of small insects.
2965. Sunbird. ♀ immaturae.
2966. <sup>Cinnyris bouvieri</sup> ~~Nyctangor dybowskii (Ousta.)~~ Small gray Weaver with red back and rump, and sides spotted with white. ♂. t.w.e. Iris red, rim of eyelids slightly swollen and purplish pink in color; bill black; feet dusky brownish. Stomach contents: tiny seeds.
2967. <sup>Erythrocercus dybowski</sup> Same as preceding. ♀. ovary slightly enlarged. Iris red, rim of eyelids gray; bill black; feet dusky brownish. Stomach contents: tiny seeds.

These little weavers were found in the grass and tilled land at the base of the hills, associating with two other related species (represented in our collection by nos. 1982 and 2251.) I have never seen them elsewhere.  
(seen hopping or walking at base of a hill, near Vankurekhovenville. Aug 3.11)

uba

july 15. 1911.

2968. Ground Hornbill (Bucorvus) ♂ [Immature] albusinicus.  
very small. Iris dark brown, skin of orbits and  
throat bluish; bill & feet black. Stomach contents:  
pieces of beetles and large ants, seeds, and  
other vegetable material.

One of three that were shown me by a native  
this morning, as they sat in a tree in a  
swamp. Their long legs give them a most  
unusual aspect for hornbills, but their  
flight is more typical of the family, buoyant,  
and with the usual alternation of flapping  
and swooping. Two of them had the skin  
of the throat blue, the other one red. They  
went to feed in some cultivated fields, and  
then in newly cleared land, but were very  
wary. At a little distance the flight seems  
noiseless, this in rising from the ground they beat their wings loudly.  
The following day we saw another, with  
red throat, in a plantation; and a day  
later heard ~~the~~ call at dusk, a low, deep,  
"hoo" repeated at short intervals.

2969. Warbler, <sup>Hylis flavogaster</sup> with steel blue upper parts, white  
patches on wing coverts, and buff breast. ♂. t.n.e.  
Iris very dark brown, maxilla blackish, mandible

Aba.

July 15. 1911.

bluish gray, feet dark bluish gray. Stomach contents: tiny beetles. Was feeding amid foliage of a small tree in a plantation.

July 16.

2970. Small Wood 1400 spec. ♀. ovary slightly enlarged.  
*Scotopelia aterrimus* *minimus*? Iris dark brown; bill gray with base black, and corners of mouth yellow; feet black. Stomach contents: small insects.

2971. Creeper. ♀ immature. Iris dark brown; maxilla *Salpornis salvadoris* *minimus* dusky brown, mandible buffy gray. feet dark, bluish gray.

Came flying across road in the undulating style of a woodpecker, lit on a small dead tree, and started to climb upwards with all the ease of a nuthatch. The only one I have seen.  
(One seen near Farage Aug 12)

2972. Warbler. ♀ immature. *Hylis flavigaster*.

3. 2969. *Eunectes guineensis*

2973. Small Red-backed Shrike. ♂ imm. Not uncommon about Aba.

*Campephaga phoenicea*

2974. Red-shouldered Cuckoo Shrike. ♂. testes slightly enlarged. Stomach contained a small green caterpillar.

July 17.

2975. Small Lark. ♀. o. w. e. Collected on top of a hill; but they feed also in cultivated fields close by.

In

Aba  
July 17

297

297

29

July  
29

29

29

Aba

July 17. 1911.

115-23 July

*Buetholaea clericalis*

2976. Small Black Chat with white forehead and wing patches.

♂. t. n. e. Common about Aba, mostly near gardens and other cleared land, but sometimes on hills as well.

*Riparia rufigula*

2977. Dusky Swallow with buff throat. ♂. testes somewhat enlarged.

*Indicator indicator*

2978. Honey Guide (*I. sparrmanni*) ♂ immature. (in plumage like that of female, save for a few scattered black feathers on throat.)

*Indicator minor*

2979. Small Honey Guide. ♀. ovary somewhat enlarged.

Iris dark brown, orbits dark gray; bill blackish with base of mandible gray, feet greenish.

These two honey guides were collected about a huge rock close to the frontier on the road to Ye. Of the larger species there were three or four; one adult male flying wildly about and calling excitedly. There were two of the smaller ones, hopping <sup>silently</sup> about the boughs of trees a little way up the side of the rock.

July 18.

2980. Yellow-breasted Bunting. ♂. testes slightly enlarged.

*Emberiza sahariensis*

Stomach contents: insect remains. Common

at Aba.

- 2980 a. *Phoenicurus familiaris* ♀. o. n. e. S.C. small insects & a few <sup>flies</sup>.

2981. Warbler with gray back, rusty crown, and pale buffy

*Cisticola petrophila*.

Aba.

July 18. 1911.

breast. ♂ testes somewhat enlarged. Irides light brown; maxilla dusky brown, mandible brownish gray, feet pale buff. Stomach contents: tiny insects.

On areas of smooth stone, where grass cannot grow, small accumulations of soil often support spiky plants like pineapple tops. (*Alocos*) bearing red flowers on a central stalk, as well as clumps of a light green vine, pliable but

X. *Sarcostemma* tough, branching at right angles and bearing <sup>not sometimes a terminal tuft of small white flowers.</sup> <sup>\*  
masses of lichen  
and in some places  
cactus-like euphorbias</sup> <sup>Asclepiadaceae.</sup> no leaves. There are also other plants peculiar to such spots but less striking in appearance; and altogether they form small patches of low dense vegetation, where hop and skulk these warblers, which

occasionally venture out on to the flat rock. (Planted in a patch at Van Buren's house. Song - a musical trilling effort, something like a *Lippia's song*.)

2982. Gray Catbird. ♂. testes slightly enlarged. *Coprinolagus trinacrolutes* (tristis) <sup>nest</sup>. Calling at dawn, was followed up and shot.

2983. Large gray Cuckoo Shrike ♂. testes very small. Iris dark brown, bill & feet black.

Stomach contents: a green caterpillar and finely divided insect remains.

2984. Thrush. ♂.

2958.

July 19. 2985. Falcon (*Aesalon ruficollis*\*) ovary slightly

1 n

Aba

July 19, 1911.

enlarged. Crop and stomach contained flesh and bones of a weaver bird, identified by its beak.

While on a hill near Ibu's village we saw a pair of these falcons fly by. In another place I saw one sitting at the edge of a pool of water on a rock as tho bathing. The present specimen was shot from a tree.

2986. *Red-rumped Woodpecker*. ♂. t. n.e. Stomach contained beetle larvae, and pieces of adult beetles.

2987. *Small Black Chat*. ♀ immature. Stomach contents: small insects.

2988. *Yellow-breasted Bunting*. ♂ ♀ immature. Stomach contents: pieces of a small grasshopper and other insects.

(Dajin - Morgan) July 26. *Chlorophoneus auripennis* *beetles*

2989. *Yellow- & green Shrike* with orange breast. ♀. ovary slightly enlarged. Stomach contained a caterpillar and remains of insects.

2990. *Woodpecker*. ♀. ovary somewhat enlarged. Iris dark red, bill gray, feet greenish. Stomach contained insect larvae.

2991. *Young of preceding*. ♀ immature

2992. "Fly catcher-Shrike". ♀ o. n.e. Iris dark brown; *Nilagus affinis* mandilla black, mandible bluish with tip black;

Aba - Faradjie  
July 20, 1911.

- feet blue gray. Stomach contents: insects.  
2993. *Pycnonotus cinnamomeus* Small Black Chat ♂ imm. Stomach contents: insects.
2994. Sunbird ♂ t. e.; skull not quite completely ossified  
*Cinnyris splendidus* Stomach contained tiny insects, but to judge from the sweet flower-like odor there were flower juices present as well.

The present specimen was secured near Madafili, the village of Maripili, half way between Aba & Faradjie. It occurs at Aba, as does also the white-breasted sunbird, neither of them having been seen so far at Faradjie.

2995. *Enicocetes albicauda* Small Red-backed Shrike. ♂ immature.

(Mandjari) Stomach contained remains of insects.  
July 21.

2996. *Ploceus superciliosus* (Shell.) <sup>Scamandracula</sup> Red & Black Weaver (*Pyromelana*) ♂. testes slightly enlarged. Stomach contained small seeds and other tender vegetable material, as well as pieces of a few insects.

2997. <sup>Ploceus</sup> *weaver*, with yellow front and breast, black cheeks & throat, brown back wings & tail. ♂. testes slightly enlarged. Iris dark brown; maxilla black, mandible light bluish; feet pinkish buff, claws dark brown. Stomach contents: seeds and insect remains.

Paradje

July 21, 1912.

*Placens superciliosus* (Shell.)

2998 Same as #2997. ♂. testes very small. Stomach contents: remains of small insects. Iris dark brown, maxilla brownish black, mandible light bluish gray. feet pinkish buff, claws brownish.

A common species of the tall grass, from Niangara to Aba.

2999. Shrike. *Laniarius major* ♂. ovary slightly enlarged. (Maxilla abnormal)

3000. Kingfisher. *Gentianula viridis* ♀. o. n. e. (*Haleyon sivegarinis*? *cyanurus*)

3001. Long-tailed Fly catcher. *Tyrannus swinhonis* ♀. o. n. e.

July 22. 3002. Rufous crowned Swallow. ♀. ovary slightly enlarged.

July 23. 3003. " " " ♂. testes .. "

3004. " " " ♀. ovary .. "

3005. " " " ♂. immature.

These swallows were shot as they rested on the roofs of houses in the post. Nos 3002 & 3 were mates, no. 3004 came later, accompanied by three immature birds, of which no. 3005 was one.

The species has been observed also about the post of Dungu.

(The steel-blue swallow with white throat spot occurs along the ribali above Dungu.)

Faradje.

July 25, 1911.

3006. Thrush (*Cossyphax*?) ♀. <sup>verticalis</sup> ovary somewhat enlarged.

Stomach contents: pieces of small beetles.

A common bird, but shy. frequenting not only the woods near water but any patch of dense brush available, this specimen having been secured in a hedge of lemon and guava trees in the post. The pleasant song, often loud, but again very low as the at a great distance, is most frequently heard in the morning and late afternoon. It often resembles strongly, perhaps imitates, the notes of *Turdus pelios*, and I have heard the calls of several other birds reproduced with amazing skill, including those of the hawk *Asturinula*, a cuckoo (*Cuculus*), and a goatsucker (the *Caprimulgus* we collected at Niangara). (Aug 13. Has since been heard to mimic the small streak breasted Nalyon, as well as the *Sobolo* (*Chrysococcyx maculatus*)).

3007. Screech Owl (*Aluco*) ♂ t. n.e. Iris dark brown, *Tyto alba affinis*. Ann of eyelid dark brown, beak pale pinkish gray. claws and scales on feet dusky brownish.

One was seen to fly out of a crevice in rocks near Aba, and today's specimen was

Faradje  
July 25, 1911

(Sunnuk)  
July 25, 1911

30

30

In 1912 we  
had them  
in October &  
of May

# Paradjie

July 25. 1911.

among the rubber trees (manihot) of the post,  
being mobbed by small birds.

(Second day on road to Vankerkhovenville)  
July 29.

3008.  
3.3696.

Large gray Hawk, with wide crest. ♀. left ovary  
*Circusetus cinereus*. enlarged. right ovary much smaller. Iris dull yellow;  
bare basal half of bill yellow, tip black; feet yellow claws  
black. Stomach contents: lizard scales and  
pieces of beetles. Shot from a tall tree in a strip  
of woods along a stream.

3009. Turtle Dove. ♂. testes enlarged.

This dove is common at Paradjie, Aba, and  
Vankerkhovenville, and was noticed up to a  
distance of 6 or 7 hours march from Dungu, on  
the road to Paradjie. Since a couple of weeks

In 1912 we first  
hear them call  
in latter part  
of May.

it has been calling <sup>commonly</sup> the male uttering a  
or 6 short, series of which the next is highest pitched, the final  
series of 5, low cooing notes, either from a perch,

or as he walks about on the ground after a  
female, lowering his head and swelling the

upper breast exactly as do domestic pigeons.

The flight is accompanied by a slight whistling of wings, audible only at short  
distances.

The other turtle dove, with a black crescent  
on the back of its neck, that was so common  
and noisy from Niangara to Paradjie in  
the dry season, is now entirely absent.

*Gida decolorata* (Linn.)  
dark cinnamon brown above, with  
white spots on the wings and tail,  
white below, with a black crescent  
on the back of the neck.

3010. Tiny Warbler. ♀. ovary somewhat enlarged.

With one or two others, was hopping about the branches of small-leaved acacias.

- Vanderkloovenville. (Native name Nzoro, from river that flows by it.)  
July 30.

3011. Large streaked brown Shrike with long tail. ♂  
*Corvinella corvina*  
t.n.e. Iris dark brown, rim of eyelids and bill yellow; feet dark green. Stomach contents: insect remains.

3012. Same as preceding. ♀. ovary slightly enlarged.  
Iris dark brown, rim of eyelids and bill yellow; feet dusky greenish. Stom. cont.: insect remains
3013. Same as preceding. ♀ immature. Stomach contents: insect remains.

Members of a scattered flock of about 10, resting on stumps and trees about a native plantation. Their call is loud and grating, and they did not allow a close approach.

3014. Black & white Widow bird, <sup>*Vidua senegal*</sup> ♂. t. e.; but skull not quite completely ossified. Crop contained small seeds.

Vanderklovenville. (= Nzoro)

July 31. 1911.

*Colius passer concolor*

3015. Black Widow-Bird. ♂. testes slightly enlarged
3016. Small green-backed Cuckoo. ♀. ovary much enlarged  
*Chrysocolaptes cassini*  
 an egg in oviduct. Stomach contained remains of caterpillars and insects.
3017. Black Weaver with yellow wing coverts and rump.  
*Euplectes cassinii*.  
 ♂. t. e. Common at V-k-h-ville & on road to Faradje.
3018. Gallinule. ♂. t. n. e. Iris rather dark brown.  
*Porphyrio alleni*. Bill and feet brown with a decided tint of red, particularly on the tibia and the base of mandible  
 Stomach contents: small seeds.

Was feeding among a few grasses along the shore of the Nzoro at the post. After being wounded it swam on the water, and then (according to white officials, as well as natives, a mistake has been made in the names of the rivers here. The Nzoro is the river at the post, the Kitabi is crooked a half-hour or so to the N.) promptly dived rivers here. The Nzoro is the river at the post, the Kitabi is crooked a half-hour or so to the N.)

- Aug. 1.  
 3019. Black-and-white Kingfisher. (*Ceryle*) ♂. ovary slightly enlarged. Stomach contents: bones & scales of fish.

3020. Small grey Slareola with white malar mark.  
*G. nucifrons*.  
 ♂. testes slightly enlarged.

- Alcedo lugens  
 3021. Gray Flycatcher. ♂. testes slightly enlarged.  
 Common along the wooded bank of the Kibali, at Dungu as well as here, perching on sticks and branches close to the water.

During this visit to Nzoro I met Hippolyte de Methelin de Pajigny.

1m

Vanderklovenville. (= Nzoro)

Aug 1. 1911.

3023. *Hirundo nigrita*, Steel Blue Swallow with white throat patch. ♀. t. n.e.

One of a pair alighting on a snag sticking out of the Kibali.

The rufous-crowned swallow also occurs at Vanderklovenville, and it too, as I have noticed also at Faradje, has a liking for water, often alighting on rocks in the rivers.

3023. Brown Boatswain (*Caprimulgus* <sup>nigriscapularis</sup>) ♂. t. n.e.

Iris dark brown, rim of eyelids light brownish, bill brown with blackish tip, feet brown.

Shot in woods on the bank of the Kibali, after being flushed from the ground.

3024. Red-throated Sunbird, <sup>*Chalcomitra senegalensis* acuta</sup> ♂ immature.

Aug 2.

3025. Black Weaver with mape and upper breast orange red.

*Pyromelana avorsa*

♂. t. e. This dark brown, bill black, feet dark brown. Stomach contained small pieces of insects and vegetable material.

3026. Small yellow-throated Bee-eater. ♂. t. n.e.  
*Melithreptus pusillus*

3027. Black Weaver with yellow wing-coverts & rump.

*Agelaioides badius* <sup>*cessorinus*</sup> ♂. t. e.

The birds themselves sometimes sit with the rump feathers overlying the secondaries.

1 m

Vanherckhovenville (= Nzoro)  
Aug 2. 1911

so as to exhibit the beautiful golden yellow patch.

3028. Red-<sup>v</sup>-black Weaver (*Ploceus* <sup>taeniatus</sup> *taeniatus*) ♂ t.e. Molting into nuptial plumage. Common, of course, at Vkhville

Aug 3.

3029. *Hypothymis ocularis*.

- Yellow Weaver with black line thru eye. ♀. t.e. but skull not completely ossified. Stomach contents: insects. Rather common, its peculiar "chirring" note being heard much more often than its author is seen. Their nests are usually hung, from the ends singly. of the boughs of acacias, at an elevation of 12 or 15 feet, but the great majority are empty.

*Crateropex* sp.

3030. Large Brown Bullbul with "scaly" plumage on throat. ♀ ovary slightly enlarged. Iris yellowish white; bill black. Stomach contents: insect remains.

In its habits and loud harsh notes it resembles the related grayer species (s. no. 2919), which frequents high grass and scrub on dry ground, while the present species is more partial to the neighborhood of water, haunting for example the wooded banks of the Ribali, and sometimes scolding passing canoes from concealment.

Vankerekhovenville. (At mt Saima, near village of Saduma)

Aug. 4. 1911.

3031. *onycotodus harrabi* Large Wedge-tailed Starling. ♂. testes slightly enlarged.  
Iris orange red.
3032. Red-&-black Weaver (*Pyromelana*). ♂ imm.; t.e. (in  
full nuptial pl.) Stomach contents: seeds & insect remains
3033. *Spermophilus refugia* Dusky Swallow with buff throat. ♀. ovary somewhat  
enlarged.
3034. Same as preceding. ♀. o.e.

No. 3033 had attached her nest to the side  
of a cliff, where it was securely protected from  
above by an arched projection of rock, and could  
only be approached from below to within 11 feet.  
Externally it was a thin, frail, half-cup shell  
composed of pellets of mud, but had a thick  
padded lining, consisting mainly of feathers,  
but with thin dry strips of vegetable material below.  
The feathers were principally from fruit pigeons  
but there were to be noticed a few small plumes of  
goatshakers, resplendent starlings, and of the  
swallow itself. Her two eggs, just about to  
hatch, were white, rather thickly spotted with  
brown, sometimes slightly rufous, these spots lying  
thickest in a circle about the large end.

Close beside this nest was an old abandoned  
one; and a couple of feet away, in a similar

Vankerekhovenville  
Aug. 4. 11.

3035  
Aug. 5.

Aug. 6  
3037

3038

3039  
3291

Vankerehovenville

I believe the hill we climbed was the same as that visited by Boyd Alexander. It was in much the same line as the Saima proper, but not quite so high. It was, however, to be preferred, for <sup>in coming from Ngolo</sup> tho' a few steep outcrops of rocks could be made out on the eastern side of Mt Saima, it was for the most part covered with high grass, scrub, and small trees; and I had already found at Aba, as well as close to the post of Vkhville that the typical hill-birds and mammals preferred bare rocky situations, even tho' not nearly so high.

We lodged in the village of a chief named Gaduma, and so had quite a long walk before reaching the base of the hill. This path was bordered by high, dense, cane-like elephant grass, and passed 2 small <sup>groups of little</sup> hamlets. Then one began to climb up the rocks. In one place there was still a large patch of cane to be passed a very disagreeable feature, that I had been spared at Aba. It was on these rocky slopes the the fruit-pigeons and white-crested Turacos were eating the fruit of wild fig trees, but I believe that Alexander, like myself, did not care to collect the pigeons. I looked at them, of course, with the glasses, and noticed no difference from the green pigeons of Taradje Vinago calva, of which I had already only too many examples.

The top of the hill was divided into several parts by gullies, the real summit, to

the south, being fairly bare and level, but other parts having ledges, huge blocks of granite, as well as trees and brush, so that I found the shyer birds difficult to approach.

Unfortunately I could only ascend the hill twice, and my second trip was marred by the effects of some canned lobster, eaten with the *Pie nieberding*, who happened to be also at Saduma's. A third visit was paid to the lower rocks, for the white-rumped swifts.

The first morning we went up in a dense fog, but when this had cleared the view from the summit was very beautiful, as it always was in these regions if one could only get up a little above the level of the monotonous scrub. Off toward Nyoro and ~~Korobi~~ were great numbers of green hills, while closer by, low down were low rocky eminences, and groups of native huts standing in the midst of their plantations. Turning in the other direction one could follow the winding <sup>wooded</sup> ~~wooded~~ border of the Kibali, which has a few more hills on its left bank.

The aloes and other rock-loving plants were much the same as at Aba. I am surprised that Boys Alexander does not mention the baboons, of which we found a band, very wary, on each visit, nor the *Hippotesci*.

Gaima is of course one of the principal

features of the landscape when viewed from the higher places near Bangura's, and still looks pretty large from the hill at Piaga's.

Long after my visit I was told by Matari and others that there was a cavern, or caverns, somewhere in the vicinity, which served as a refuge for the manu in their conflicts with the troops of the Congo State.

In Sept 1912 monsignor Van Schoote told me that he had been to the top of the Gaima, starting from Saduma's, the natives having previously cut him a path. He took observations with an aneroid, he said, besides measuring "hyprometriquement". Saduma's village was found to be at 662 meters above sea-level, and the top of the hill 1096 m., making the relative height 434 meters, which <sup>seemed</sup> ~~really seems~~ to me greatly exaggerated. (But he may well have been right.) Alexander gives the height as 400 ft. only, but I should not object to another hundred feet. The highest hills near Aba I estimate at 350 <sup>Pungu N. of "Baramba" at 300-350 ft.</sup> feet, and Namambula, NW of Pawa, at 600. The highest hills seemed to me those between Mararamana and Bafwabofla. The road only goes over 5 or 6 low rises <sup>up</sup> at the range and the dense forest prevents one from seeing much, yet as viewed from Mararamana they appear at least 650 ft high. All this is of course estimated, from comparisons of large trees growing on the slopes, from shot-gun ranges, and from looking about while climbing.

The hills at Baramba, Aba, and Vankeerchhouw are all granitic. Namambula has a ridge of hematite across its top, and is probably composed <sup>inside</sup> of the same material, boulders of which are to be seen <sup>near</sup> Mararamana.

Vanderklovenville:

Aug 4. 1911.

position, the mud shell of a new nest had just been completed, one of its builders coming with a feather as we watched.

[A streak-breasted swallow was seen near the post of V.-K.-ville, as well as near Lorraine, where I actually shot one but lost it in the cane-grass.]

3035. Ibis. ♂. testes slightly enlarged.

Aug 5. Hagedesie bearded *Sarcops calvus*

3036. Rail (*Corethrura*). ♂ testes slightly enlarged.

Shot while walking on the stones in a little wood-bordered mountain brook. ♀ one sits very still to await them. They are not very shy; a female was observed at very close quarters a few minutes after, and as she was too close to fire, I thought to frighten her further away. At the first stir, however, she darted back into the woods. To judge from their notes, they must be ~~extremely~~ common here.

Aug 6. 3037. Small Streaked Kingfisher. ♀. o.n.e. (*Halcyon chelicuti*)

3038. Large Wedge-tailed Starling. ♂. testes somewhat enlarged. *Onychognathus fuscatus*. Stomach contents: a green fruit as large as an olive and with a large stone. Shot on the same large dead tree as no. 3031. The call, heard at a distance resembles slightly the peep of a pullet.

3039. <sup>5-2958.</sup> Thrush, black, with white wing coverts and rufous rump & belly. ♂. t.n.e. Stomach contents: fruit

Vanderklovenville (= Nzono)

Aug. 6. 1911.

of a wild fig tree (*Ficus*). Rather common on the rocky hills, at it was at Aba.

3040. Red-e.-black Weaver (*Pyromelaena*) ♂. t.e. *flavimaculata*.

3041. Warbler, <sup>*Euphopygia* sp.</sup> with gray back, white superciliary line & wing bars, rufous rump, and white belly. ♂. t.e. Iris dark brown, bill black, corners of mouth pale yellow, feet bluish gray. Stomach contents: insects.

Rather common, especially about the high cane-like grass known here as Baka (Matete in Kingwana). The song, given by the male from the top of a small tree, consists of half a dozen short notes, and is repeated at short intervals, but is not particularly musical.

3042. Thrush (*Cossypha*?). ♀. o.e. Stomach contents: 3 hard stones from fruit, remains of insects and a tiny millipede.

3043. Black Swift with white throat & rump. ♀. ovary <sup>reticulata</sup> <sub>*micropterus caffer streubeli*</sub> slightly enlarged.

3044. Same as preceding. ♀. ovary slightly enlarged.

There were 2 pairs of these swifts flying about some huge isolated boulders at the base of the hills.

There was an overhanging ledge, which bore on its under surface a couple of old nests of streak-breasted

Vanderklovenville (= Nzoro)

Aug. 6. 1911

swallows [but I believe that a narrow horizontal crack was more probably their retreat. (One of these swifts was noticed at Abo).]

3045. Coly. ♂ testes enlarged.  
*Colius nigerfthalmus*

3046. " " "

3047. " ♀. ovary "

3048. " ♀ " somewhat enlarged.

Aug. 7. 3049. .. ♂ testes much enlarged.

Stomachs of all 5 contained earth.

These colies were captured by native in a small cave on the steep bank of a wooded ravine. Its entrance opened between the roots of a tall tree, and was large enough for a man to crawl into on hands and knees. The inner chamber, directly beneath the tree, measured



fairly level floor  
b. The earth was  
c. Here the colies,  
eat the earth, and  
inside entrance is framed  
to size, and at the  
piece of cloth, that  
ng the exit. A  
birds enter, and  
practised in the

day time, the birds do not come there to sleep.

Vanderklovenville (= Nzoro)

Aug. 6. 1911

swallows [but I believe that a narrow horizontal crack was more probably their retreat. (One of these swifts was noticed at the entrance.)]

3045. Coly. ♂ testes enlarged.  
*Colius n. longipennis*

3046. " " " "

3047. " ♀ ovary "

3048. " ♀ " somewhat enlarged.

Aug. 7. 3049. " ♂ testes much enlarged.

} Stomachs of all 5 contained earth.

These colies were captured by native in a small cave on the steep bank of a wooded ravine. Its entrance opened between the roots of a tall tree, and was large enough for a man to crawl into it on hands and knees. The inner chamber, directly beneath the tree, measured approximately 10 feet x 6, with a fairly level floor and domed roof, some 5 feet high. The earth was a stiff red clay, free of large stones. Here the colies, as well as other birds, come to eat the earth, and are captured as follows: The entrance is framed with green branches to reduce its size, and at the top of this door is attached a piece of cloth, that can be lowered suddenly, closing the exit. A naked black sits inside. The birds enter, and he shuts the opening. This is practised in the day time, the birds do not come there to sleep.

Vanderklovenville (= Nzoro)

Aug. 7. 1911.

by pigeons is said to exist up on the mountain.

I asked why they did not fear the man inside, but they replied that the darkness prevents them from seeing him.

(For the eating of earth by doves, see no. 3308.)

*Ptilocichla turdina*

3050. <sup>s. 2871.</sup> A brown-backed Thrush-like bird, with white underparts,

spotted on the breast with brown. ♂. testes somewhat enlarged. Iris bright brown; mandible and tip of maxilla light blue-gray, rest of maxilla blackish. feet pale flesh color. Stomach contents: a caterpillar and 2 winged termites

3051. mate of preceding ♀. o. e. Stomach contained insect remains, esophagus 2 winged termites.

Secured in damp woods along a brook. They call loudly very early in the morning, but are seldom heard at other times of day.

Aug. 8.

3052. Sunbird with green back, gray belly, and metallic green head. <sup>Cinnyris ruficauda versicolor pluvialis</sup> ♂. testes much enlarged.

3053. <sup>say Thryophilus sp.</sup> Warbler ♂. testes somewhat enlarged.

3054. " ♂. " slightly " . Stomach contained pieces of a small millipede.

Vanderklovenville (= Nzoro)

No. 3055. (Note - Niangara, June 9. 1913) 2 young of this hawk (*Asturimula*), with tails not fully grown, were brought by natives about a week ago, and are being kept alive by an agent. Their plumage is light gray, with bars below, and black throat-stripe, just like adults, and they utter continually the low whining note.

three, syllables.

*Yungip cuo beletie*

3061. Small Gray Woodpecker. ♀. o. n. e.  
3062. Small Green-backed Cuckoo. ♂ testes somewhat enlarged  
*Chrysococcyx klaasi*  
Stomach contents: remains of caterpillars and a bug.  
Often heard calling. The larger yellow-breasted  
"Gliotocoli" has never been heard at Faradje.  
(One afterwards heard at Faradje - in October)  
Between Niangara and Dungu, in June, the  
notes of only two reached my ears.
3063. *Sitagra taenioptera*.  
Yellow Weavers, with black face and throat. ♂.  
3064. t. e. These weavers were common at Vanderklovenville,  
and in some papyrus on a little island in the Nele, <sup>on Aug. 9.</sup> Two of  
their nests were found, containing each 2 eggs. One nest and  
its set of eggs has been preserved (labelled no. 3063). Today,  
on the road to Faradje, more nests were found in  
papyrus bordering a stream. The eggs, of which 4 sets  
of 2 each were found, were all incubated, and so only  
one egg and a nest were preserved (labelled no. 3064).

Vanderklovenville (= Nzoro)

Aug 10. 1911

3060. Mate of no 3059. ♀. o. n. e. Stomach contents: a small green caterpillar, a beetle, and other insects.

These birds search the foliage of trees, often acacias; and are apparently rather common about Vanderklovenville and Faradje. The long-drawn sibilant notes, which seem to serve as a song, are of two, or sometimes three, syllables.

(*Youngicus beobletus*)

3061. Small Gray Woodpecker. ♀. o. n. e.

3062. Small Green-backed Cuckoo. ♂ testes somewhat enlarged  
*Chrysococcyx klassi*  
Stomach contents: remains of caterpillars and a bug.  
Often heard calling. The larger yellow-breasted

"Goliathocole" has never been heard at Faradje.  
(One afterwards heard at Faradje - in October.)

Between Niangara and Dungu, in June, the notes of only two reached my ears.

3063. *Sitagra taenioptera*. A Yellow weavers with black face and throat. ♂.

3064. t. e. These weavers were common at Vanderklovenville,  
and in some papyrus on a little island in the Nelle, two of  
their nests were found, containing each 2 eggs. One nest and  
its set of eggs has been preserved (labelled no. 3063). Today,  
on the road to Faradje, more nests were found in  
papyrus bordering a stream. The eggs, of which 4 sets  
of 2 each were found, were all incubated, and so only  
one egg and a nest were preserved (labelled no. 3064).

Faradje

Aug 12. 1911.

3071. *Nesocharis capistrata*  
Weaver. ♀. a much enlarged.  
3 2 3 3 4.

Was hopping about the branches of a tree in such a fashion as to suggest strongly a titmouse, especially when its colors are taken into consideration. Indeed

the first one I saw at Dungu I took for a titmouse.

3072. *Amblyospiza* (colored like *A. capitalis*, but larger. W. 92.5 T. 70.)  
white-fronted Weaver (*Amblyospiza*) ♂. t.e.

Near the post of Faradje I have never met with this weaver. On the road to Dungu, however, a few were seen about papyrus swamps, and No 3072 was shot some 10 miles south of Faradje, where there is much of the <sup>tall</sup> candle-like grass known here as Baka. About Medje, it will be remembered, their nests were always placed in the same kind of grass.

Aug 11. 3073. *F. intermedia* Partridge. ♀. ovary slightly enlarged. Crop contained termites (not winged) and pulp of fruit.

Aug 13. *Hirundo smithi*

3074. Rufous-crowned Swallow. ♀. ovary slightly enlarged.

Aug 14. 3075. Mate of No. 3074. ♂. testes slightly enlarged.

3076. Sparrow. ♂. t.e.; but skull not quite completely ossified. Passerinae. Not numerous in the vicinity of Faradje.

Aug 15.

3077. Goatsucker (*m. vexillarius*) ♀. ovary somewhat

Faradjic

Aug. 17. 1911.

of nests that contain no eggs. These empty ones, however, are usually of poorer construction, often partly transparent, while all the nests are placed at a height of 8 to 10 feet above the water. Those suspended in papyrus-tops have generally an outer network of fibers from papyrus, but internally are composed of grass. It is almost unnecessary to describe their form, for it is that of a typical weaver's nest, the entrance pointing downward, without prolonged spout.

As far as I have seen, the set is invariably two, and while the eggs in different nests differ from brown to light green (spotted with brownish), those in the same nest are alike. Two nests and their complement of eggs were collected today.

Aug 18. *Calamocichla* sp.

3085. "Papyrus Bird": ♂ t.e. Stomach contents: small insects.

3086. " " " ♂ t.e.

3087. " " " ♀. ovary slightly enlarged (Mate of no. 3086)

Very common at Faradjic, heard also near Aba.

3088. *Crotophaga* sp.  
S. 3085. Large brown Bullbul with "scaly" plumage on throat. ♂

testes somewhat enlarged. Stomach contents: fruit and pieces of a small millipede.

Faradje

Aug 19. 1911.

the north about the middle of July.

*Serpophaga viridis*

3094. Paradise Flycatcher. ♀. o.m.e. (molting). Iris dark brown, bill blue with black tip, rim of eyelids and feet blue.

3095. Large yellowish Weaver. ♂. t.e. Crop and stomach contained maize and ants. Nesting in numbers on trees along the Dungu. During last dry season, these trees were completely deserted, and bore only old ~~abandoned~~ nests, whereas, in the *Sturi* [With this number, (3095) is written above the bracket] I believe this species nests all year round. [This species, also a single egg from a colony in a large tree on an island in the Dungu, were sent to the post office Sept 20-30, 1911.]

3096. Red-wattled Flycatcher. ♀. o.m.e. The female of this fly catcher in flight sometimes beats her wings noisily like the male.

3097. Cossypha. ♂. testes somewhat enlarged. Stomach contents: insect remains.

3098. Rufous-crowned Swallow. ♂. testes somewhat enlarged.

3099. " " " ♂ slightly "

- Aug 20. " " " ♂ " "

3100. " " " ♂ " "

3101. Heron. ♂. testes very small. Stomach contents: *Ardea p. purpurea* fish.

- Aug 21. 3102. Small Red-backed Shrike. ♂. t.n.e. St. cont.; insects.

3103. Swallow. ♂. t.n.e.

3104. Paradise Flycatcher. ♀. o.m.e.

Paradye

Aug 25. 1911.

*F. icterorhynchus*

3115. Partridge ♂ t.e.

*Sitoger taenioptera*

Yellow Weavers with black face & throat. ♂ t.e.

3117.

3118. Black - & white Widow-bird. ♂ t.e. skull not quite completely ossified. *Vidua serena*

3119. Same as preceding, ♀. very much enlarged; skull not completely ossified. Crop filled with small seeds.

Aug 26.

3120

*Malacoptilus poliocephalus*

Large yellow & green Shrike, with gray crown &

cheeks. ♂ immature. Stomach contained a small millipede and a bee.

Aug 27.

3121.

Hawk (*Polyboroides*) ♂. testes not enlarged. Irides *Gymnogyps typicus*. blackish brown. cheeks yellow, cere and base of mandible whitish, horny part of beak black, feet bright yellow. claws black. Stomach contained 3 young colies (*C. leucophthalmus*, *migricollis*), which must have been taken from the nest, for their primaries, etc., were still enclosed in sheaths.

Shot along the river above the post, where it alighted in leafy trees, seeking concealment in the foliage. Its tarsal joint could be bent a little way backwards. The supraocular shield was very small. This strange long-limbed bird of prey was observed 3 times in the vicinity of Medje, once

1/n

Paradise

Aug 28. 1911.

3124. *Cinnyris cupreus* <sup>verticalis</sup> Bronzed Sunbird. ♂ immature. Testes much enlarged.

Aug 29. 3125. *Turdus (Cossypha) verticalis* Thrush. ♂. testes somewhat enlarged.

Stomach contents: a blue berry, and remains of small beetles.

3126. *Turdus (Cossypha) verticalis* ♂ immature. Stomach contained insect remains and seeds (of fruit?).

3127. *Hypsipetes ocularis* Yellowish Weaver, with black throat & line thru eye. ♂ immature, testes slightly enlarged. Stomach contained seeds & insect remains.

3128. *Indicator indicator* Warren Guide. ♂. t.n.e.

Nos. 3122+3. *Struthio camelus semitorquata*. This pigeon has at least four different types of notes:

1) Most characteristic, loud cooing, repeated over and over. "coo coo, cu-cu coo coo"

2) A shorter performance, also oft repeated.

"c-c-c-coo, coo coo"  
Hoarse, low. louder

3) A low, hoarse protracted cooing sound, sometimes repeated, usually uttered when two are together - probably connected with courting.

4) A hoarse "Kääaaaa".

3131. Small widow bird (*Vidua*) ♀ o.e.; but shall not completely ossified.

3132. *Pipit (macronyx)*. ♂. testes somewhat enlarged. Stom. cont.; insect remains.

Paradise

Aug 28. 1911.

3124. *Cinnyris cupreus* <sup>reticulatus</sup> Bronzed Sunbird. ♂ immature. Testes much enlarged.
- Aug 29.  
3125. Thrush (*Cossypha*) ♂. testes somewhat enlarged.  
Stomach contents: a blue berry, and remains of small beetles.
3126. Thrush (*Cossypha*) ♂ immature. Stomach contained insect remains and seeds (of fruit?).
3127. *Hypsipetes ocularis*. Yellowish Weaver, with black throat & line thru eye. ♂ immature, testes slightly enlarged. Stomach contained seeds & insect remains.  
Indicator indicator
3128. Honey Guide. ♂. t.n.e.  
While we were in a clump of trees along a swampy brook near the post this bird came in and began to call loudly, keeping always about a tree where I had noticed a hive of bees a few minutes before.
3129. *Emilia lepida*. Green and gray warbler with chestnut throat and shoulder patches. ♀ im. Stomach contents: insect remains.
3130. Same as preceding. ♀. ovary somewhat enlarged. Irides brownish red. Stomach contents: insect remains.
3131. Small widow bird (*Vidua*) ♀ o. e.; but shell not completely ossified. <sup>perena</sup>
3132. Pipit (*macronyx*). ♂. testes somewhat enlarged. Stom. cont.; insect remains.

Faradjie

Aug. 30. 1911

3138. Goatsucker. ♂. t. n.e. Stomach contents: a dung-  
*Coprimulgus natalensis chadensis*  
beetle & many winged termites.

Aug. 31.

3139. Goatsucker ♂. t. n.e. Iris rather dark brown,  
*Coprimulgus natalensis chadensis*  
rim of eyelids light brown, bill brown with blackish  
tip, feet buff. Stomach contents: winged termites

3140. Goatsucker. ♂. t. n.e. Stomach contents: many  
*Coprimulgus natalensis chadensis*  
winged termites, 2 beetles, & 2 bugs.

3141. Goatsucker. ♀. ovary slightly enlarged. Stomach  
*C. n. chadensis*  
contents: winged termites and a beetle.

These goatsuckers were shot at dusk while  
flying in numbers about a termite hill whence  
the winged individuals were taking flight. Mingled  
with them were also long-tailed goatsuckers  
(*Scotornis*). With the exception of a young bird  
(no 2855) the species is new to our collection.

3142. *Estrilda sulfurea*  
Weaver. ♂. testes somewhat enlarged.

3143. " " " ♀. o.n.e.

members of a flock about a dozen in number,  
feeding on the ground amid weeds near the post.

The second time the species has been observed at

Faradjie.

- Sept. 2. 3144. Pipit (*Macronyx*) ♂. croesus.  
testes somewhat enlarged.  
Stomach contents: a small millipede & insects.

Paradise  
Sept. 4 1910

contents: 3 berries and pieces of an insect.

3151. Dark Slaty Flycatcher. ♂ t. n. e. *Melaenornis pammelaina*  
*Colius passerinoides*

3152. Black Widow bird. ♂ t. e. Stom. cont.; seeds.

3153. " " " ♂. " " " "

With the evident purpose of displaying their charms to the females, at this season the males make long slow flights over the grass, well-sustained and direct, but not as a rule more than a few yards above the tops of the vegetation. During these performances the tail is spread, but owing to its peculiar folded condition, is expanded in a vertical, instead of horizontal plane. The appearance of the



bird during such a flight I have tried to express by a silhouette, all the more appropriate for its black plumage. When speed is their aim, they are not encumbered by their long tail, but shut it tightly, and proceed with great rapidity.

3154. Thrush (*Cosyphus*) ♂. testes somewhat enlarged.  
verticilio

Stomach contents: remains of small insects.

3155. Thrush (*Cosyphus*) ♀. ovary slightly enlarged.  
Stomach contents: fruit and a small hairy caterpillar.

Paradise

Sept. 4. 1911.

and still faster as they die away. Meanwhile the male, who is making them, sits on a stalk of papyrus; but as he finishes (or sometimes just before finishing) he flutters off into the air (but keeping always amid the papyrus), where with hovering flight and tail pointing directly downwards he gives 4 or 5 loud wing beats, and then makes for a new perch, often to recommence his "song" at once. When not engaged in singing they usually keep well concealed in the tangle of old dry papyrus, saw-grass, and other aquatic vegetation that commonly fills a papyrus swamp waist-high, or even shoulder high. From this concealment they occasionally utter a single chüt, or cluck, or more rarely a single loud clap of the wings.

*Coliaceus malacca*

3159. Black Weaver, with yellow wing coverts & back ♂.

(in brown plumage) t. n. e. but skull completely ossified in spite of its apparently immature plumage. Stomach contents: small seeds. The adult males, in black plumage, while perching, often show plainly the patch of downy white feathers in the middle of the breast.

Faradje

Sept. 4, 1911.

contained tiny seeds.

*Dicrurus aemulus divaricatus*

3167. Drongs & Shrike. ♀. o.m.e. Stom. cont.; insects.

*Lyrus algutinus*

3168. Red - Black Barbet. ♀. ovary somewhat enlarged

Stomach contents: red peppers (chewed in pieces)

3169. Snipe (*Gallinago*) ♂. t. n.e.

*mellotis*

3170. Sandpiper (*Actitis*) ♂. t. n.e.

*Recurvirostra glauca*

3171. Sandpiper (*Actitis*) ♂. t. n.e.

*limosa*

3172. Bronzed Sunbird. ♂. t. e. *Cinnyris cupreus*

*Indicator indicator*

Sept. 8.

3173. Honey Guide, ♀. o.m.e. Uttered notes like those  
of a male.

*Melanopteryx nigerrimus* (Vieill.)

3174. Black Weaver (*Ploceus nigerrimus*) ♂. t. e.

Stomach contents: seeds & insects.

Uncommon at Faradje.

*Emberiza affinis*

3175. Rufous-backed Bunting. ♂ immature.

Stom. cont.; remains of insects. Iris dark  
brown; maxilla dark gray, mandible blue  
gray; feet light pinkish gray. (note Oct 1912 - This  
(note Oct 1912 - This  
was the only example  
seen at Faradje)  
another ab. 2.18)

3176. Small Black Chat. ♀. o.m.e. skull not quite  
completely ossified. Stom. cont., small insects

3177. Large Warbler, with dusky olive back, and  
*Bradypterus carpalis* white breast heavily spotted with dusky. ♂.

t. e. Iris dark brown; bill mostly black, but  
bluish gray beneath mandible, feet brownish.

Garadje  
Sept 8. 1911.



grous and grass.  
thin shells,  
now, as well as  
a larger spots  
the larger end.  
kind of attaching  
d one of last  
side-binding  
as well as  
resent seasons,  
as the nest  
2 - This species is  
seen in the high grass  
"or "matate" of Chaco  
Cerroville, Aug. 11, and

as the nest  
2 - This species is  
seen in the high grass  
ba" or "matete". Thus I  
nowville, Aug. 11, and

e. Motacilla alba

1. t. u. e

Apparently absent in the middle of the  
~~spring~~  
~~dog~~ season; this is the first I have seen for  
many months, perhaps a migrant from the  
north.

3185. Blackish Sunbird, *Chalcomitra senegalensis* ad.  
with large red breast patch. St. ne.

Faradje  
Sept 8. 1911.

had been gathered from the papyrus and grass. It contained 3 eggs, with very thin shells, white, finely speckled with brown, these larger spots being most abundant about the larger end. This must be the usual method of attaching the nest, for I found an old one of last year that still retained its side-binding about a dry papyrus stem, as well as an abandoned nest of the present season,

supported by two stalks just as the nest  
I have described. (note - Nov. 1912 - This species is  
occasionally observed in the high grass  
known as "lala" or "matate". Thus,  
have heard it on the ridge of a hill at Vankerkloofsville, Aug. 11, and  
near Faraday, Nov. 1912.)

3182. Black - & - white Wagtail. ♂. t. e. Motacilla alba  
Sept 9.  
3183. Flycatcher. ♂ immature. Say's Bell's myiotheretes  
3184. Bee-eater (merops nubicus). ♂. t. m. e.

Apparently absent in the middle of the rainy season; this is the first I have seen for many months, perhaps a migrant from the north.

3185. Blackish Sunbird with large red breast patch. St. ne  
Chalcomitra senegalensis ad

Faradje  
Sept. 10. 1911.

*Dicrurus adsimilis divaricatus*

3194. Drongo. ♂ t. n.e. When a couple of these drongos make up their minds to sing they put forth only a most disagreeable mixture of creaking, metallic notes { <sup>at Sungai, in Apr. 1912, 9</sup> <sub>and Drongos imitate other birds</sub>

3195. Spotted Papyrus Bird. ♂ t. e.  
*Bradypterus celebensis*

3196. Little Papyrus Bird. ♂. stomach contents: insects  
*Bradypterus abyssinicus*. Both the species represented by nos 3195 & 3196 have ossified tendons in the legs just above the tarsal joint; but in the gray papyrus bird, (Colomoscelis nictior) like no. 3181, etc., this hardening of the tendons is lacking.

3197. Paradise Flycatcher. ♂ t. n.e.  
*Sericophorus nigriceps*

3198. Brownish Weaver with red head. ♂ t. e. Iris dark brown, bill brownish, feet buff.

3199. Guinea Fowl ♀. o.n.e. Crop contained a great deal of dwarf millet ("malu") <sup>(Cleusine)</sup>, not yet ripe, some pieces of tubers, and several large black ants, of the termite-hunting kind.

3200. Green & yellow Weaver with black cheeks. ♂ immaturae, testes slightly enlarged.

Sept. 11.

3201. Black Weaver <sup>*Pycnonotus anzorgei*</sup>, with red crown and upper breast, and upper back yellow. ♂ t. e. Stom. cont.: greenish seeds.

Paradise

Sept 12. 1911.

woods, but never in the dryer, open places, nor about the post.

*Hypotheca chalybeata?*

- 3212 5 small Blue-Black Weaver ♀ ovary somewhat enlarged  
" " " " " skull not completely ossified
- 3213 " " " " " ♀ " "
- 3214 *Quelea cardinalis* Small Brownish Weaver with red head ♂ testes enlarged
- 3215 *Quelea cardinalis* " " " ♂ testes somewhat enlarged
- 3216 *Quelea cardinalis* " " " ♂ " "

Rather common in the high grass, of which they eat the seeds.

- 3217 *Ploceus superciliosus (Shell)*  
S. 3222  
3227 yellow breasted Weaver, with dark green back, & cheeks  
throat black. ♂



His nest was built in a very conspicuous position, on an almost leafless bush, at a height of six feet. Roughly oval in shape, it had a small round entrance on one side, high up, and was woven mostly of fibers stript from some grass or other plant that was covered with short hairs. But the softness thus given to the structure was evidently not enough, for the bottom of its interior was lined with a material closely resembling thistle-down, making the softest of beds for the reception of the three

Faradjî

Sept 16. 1911.

yellow on inner border; culmen dark brown, rest of bill as well as lores and feet greenish yellow. Stomach contents: a minnow.

Shot at 5 PM, while flying up a brook from the river. Twice before (yesterday evening and this morning) it had been seen proceeding in the same direction.

3224. *Dohrniciella apicalis* Brown Warbler with long and wide tail. ♂. t. e. Iris grayish brown; maxilla black, mandible bluish gray, feet brown. Stomach contents: insects.  
There are red fleshy tendons just above tarsal joint, as in nos 3195 & 6.

A bird of damp meadows, where the grass is not very high. From time to time ~~with drooping tail,~~ they fly high in the air, making perhaps a slight noise with the wings, and then coming down to rest on an exposed stalk of grass, or a small tree. From such perches they utter a rather weak, "cheep, cheep, cheep, cheep, cheep, . . . ."

Sept 17.

3225. Warbler ♂. t. n. e. Iris dark brown, bill dusky brown, *Sarcophagus arcturovianus* slightly greenish along edges of both mandibles, lower mandible pinkish at base, feet light grayish brown. Shot in the rank grass along a creek.

Garadje  
Sept 21. 1911.

*Dicrurus adsimilis davaricatus*

3237. Drongo. ♂ t.n.e. Was eating a large moth, holding it beneath his foot and pulling at it with his beak. He was even seen to take a short flight, grasping the moth with one foot.

3238. Gray Plantain-eater (*Schizorhynchus*). ♂ t.n.e. Stomach contents: fruit & pieces of leaves.

- Sept 22. 3239. Snipe (*Gallinago*) ♂ t.n.e.

3240. Ruff. ♂ t.n.e. *Philomachus pugnax*.

- Sept 24. 3241. Snipe (*Gallinago*) ♂ t.n.e.

3242. Ruff. ♂ t.n.e. *Philomachus pugnax*.

3243. Long-tailed Cootsucker (*Scotornis*) ♂ t.n.e.

Stomach contained remains of coleoptera, including

2 small dung-beetles. (This species is just beginning to alight on roads at dusk.)


3244. Short-tailed Warbler. *Sylvietta brachyura* ♀. ovary somewhat enlarged

3245. ~~Argos montanus~~ <sup>Argos montanus</sup> Gray-backed Weaver, with breast brown, spotted with white. ♀. ovary somewhat enlarged. On a bough, 7 feet above a small pool of water, in some swampy woods, was a large mass of old dry leaves, almost certainly a deserted concial's nest, with an entrance at one side. On a lining of matted hair and a few snake scales <sup>apparently</sup> procured from the excrement of some carnivore, this weaver had laid her 6 pure white eggs.

Paradij.

Sept. 30. 1911.

3253. Least Bittern. ♀ ovary somewhat enlarged.  
*Ixobrychus exilis*.  
 Stomach contents: a tiny fish & 2 large spiders.  
 Shot at sundown, as it flew up from a cleared area in a papyrus swamp, where it had been noticed the day before, at the same hour.
3254. Black Titmouse with white wing patches. ♂ t. n. e.  
~~*Melaniparus niger leucomelas*~~  
 Iris dull yellow, bill black, feet very dark bluish gray.
3255. Same as preceding. ♂ t. n. e. skull not quite completely ossified.
3256. Sandpiper. ♂ t. n. e.  
~~*Acrocephalus schoenobaenus*~~
3257. Warbler. ♂ t. n. e. { Collected on same spot as  
 3258. "              ♀ t. n. e. } No 3225.  
~~*Cisticola e. erythrops*~~
3259. Buff-cheeked Warbler. ♂ testes slightly enlarged.
3260. Bronzed Sunbird. ♀. o. e. *Cinnyris cupreus*.



Her penile nest, with lateral door, was composed of strips of dry grass and similar vegetable material, and lined with soft vegetable down, being suspended from a <sup>Bauhinia</sup> bush, about a yard from the ground. It contained two eggs, of very

Paradise

Oct 40 1911.

3268. *Andropadus* sp.

Green & gray Bulbul. ♂ t.e.

3269. Crested Wood-shrike (*Prionops*) ♂ t.n.e. Stomach concavata  
contents: insect remains. Bill black, corners of mouth and circumocular wattle yellow, iris light gray, feet reddish orange, claws gray.

A bird of forest, sometimes coming down low so that the snap of its beak is audible while feeding, but usually frequenting the higher trees in flocks of 6 to 10, which are difficult to follow when alarmed. I

have heard them give low hoarse notes.

Our Log guide speaking in Bangala, called them "ndege nakozi".

3270. Green & gray Bulbul with rufous tail. ♀ im.

3271. ♂ t.e. (a forest bird, *Apalis* sp., *Apalis* *millepunctata*, *Beda cyanea*, *Leucosoma mariae*)

3272. ♀ immature. (" ", olive & yellow, with rufous tail)

3273. Wagtail, with gray back, pale yellowish breast, yellow tail coverts, and very long rectrices. ♀ im.  
*Macroura cinerea*

Shot along a brook in forest.

(The common yellow-breasted wagtail, with gray crown, was first seen here this season on Sept 27.)

3274. Drongo. ♂ t.n.e. Stomach contents: 3 small caterpillars, & insects (beetles, a bug, etc)  
*Apaloderma n. marina*.

Paradje

Oct 5. 1911.

rhythm, sounding in fact as tho the two birds were calling at once, and having a little of the rolling quality of Cryptoleola.

3281. Large white-rumped Swift (*Chaetura*) ♀. o. n.e.  
A forest-loving species, not to be seen near the post.

3282. Owl (*Bubo lacteus*) ♂. t. n.e.

One of a pair roosting in a patch of forest. They were not at all shy, tho the trees were so high that they were in little danger from my shot-gun.

- Oct. 6.
3283. Small Black Hornbill ♀. ovary slightly enlarged.  
*Lophoceros fasciatus*. Stomach filled with pieces of beetles & bugs.  
Three were seen at Aba on July 11, and two today, about half way between Aba & Paradje.
3284. Woodpecker. ♂. t. n.e. Stom. cont.; insect larva.
3285. " ♀. o. slightly enlarged. Stom. cont. insect larva  
(The large woodpecker of the forest.)
3286. Barbet (*Priotelus*) ♀. ovary slightly enlarged.  
Stomach contents: fruit.
3287. Tiny Kingfisher (*Ospidina picta*) ♂. testes slightly enlarged. Stomach contents: 3 "pill-bugs" (Centopoda) while perching, this bird has a peculiar habit of raising and lowering its head from time to time, with a short abrupt movement.

Paradise

Oct. 8. 1911.

Stomach

*Selincous modestus*

3298. Lark. ♂. testes slightly enlarged. Stomach contents tiny hard seeds.

3299. Mate of preceding. ♀. ovary slightly enlarged. Stomach contents: tiny hard seeds.

This pair of larks was collected on a small granite hill, about 5 hrs. E. of Paradise. No other typical hill birds seen there.

3300. Black Weaver with yellow wing coverts & rump.  
*Euplectes cressonii*. ♂. t. e. Stomach contents: hard seeds.

- Cinnys. 3301. Tiny Sunbird. ♂. t. n. e.

Oct. 9. *Cinnyris* ~~see butiri~~ *butiri*. *Hybris aquatorialis*

3302. Red-<sup>v</sup>-black Barbet (*Pogonornis rhynchotus*) ♂. testes slightly enlarged. Stomach contents: red peppers.

3303. Same as preceding. ♀. ovary slightly enlarged. Stomach contents: red peppers.

3304. Black Barbet with red face. ♂. testes slightly enlarged. Stomach contents: fruit.

Oct 10. *Gymnomyia verticalis* ~~veidleri~~ *leucostoma*

3305. Sunbird. ♂. testes slightly enlarged.

3306. White-breasted Sunbird. ♀. imm. *Anthreptes longirostris leucostoma*

3307. Rufous-winged Shrike with black cap. ♂. *Artizelous minutus*

- testes slightly enlarged. Stomach contents: grasshoppers & beetles.

3308. Turtle Dove. ♂. t. e. Very common; still singing. Often gather in flocks and eat earth on a spot in the post where a mægaine containing salt is said to have stood. Goats, and a mule, have been observed licking the ground at the same place. The doves also came (now)

of turtle dove  
19 of them dove once  
12 were in a single  
exchange. Cases  
of all contained  
earth.

Faradje

Oct. 12. 1911.

istic "laughing" note, it sometimes utters a crow-like "caw". (In our experience plantain-eaters are noisy all year round, without reference to season)

Oct 13. 3317. Red-&-black Barbet, <sup>(Pomachynchus)</sup> *Lybius aguaticus*. Stomach contents: red peppers.

Numerous about the post, pecking large holes in ripe papayas, <sup>(of which they eat the seeds as well)</sup> in company with colies, besides eating the small native peppers, which are so hot that if I simply bite into one my mouth burns for a quarter of an hour. The fruit of wild fig trees are also devoured; and many other fruits as well, I believe. <sup>(note: with regard to red peppers being eaten by birds, see no. 8123, a pigeon which had swallowed 75!)</sup>

They often produce a loud, hoarse, and prolonged "clicking" note, which may be compared with that of the nightjar *Scotornis*. These sounds, especially that of the nightjar, resemble nothing so much as a rapid succession of electric sparks.

Oct 14. 3318. *Hirundo nebulosa*. Large brown-rumped Swallow. ♀. o.n.e.  
At this season, they roost in company

## Paradjie

Oct. 15. 1911.

3327. <sup>*Hippolais turcica ocularis*</sup> Yellow Weaver with black throat & line thru eye.

♂. testes slightly enlarged.

3328. Weaver. ♀ imm. *Vidua serena*

Oct 16.

3329. Coucal. ♂. testes slightly enlarged. Stomach contents: 2 large caterpillars & 1 small one, remains of beetles and a bug.

The voice of this coucal, heard practically throughout the year, is in full a series of 15 to 20, <sup>loud</sup> cooing notes, first descending, and then remounting again:

"too

too

too

too too

too too too

too too too

too too too

too too too

too

too

too

too, too, too, too

Sometimes 2 birds will call in unison, one starting a little after the other, or the "toos" may be fewer longer, more slowly repeated, and in one key. A very different harsh call is also occasionally heard.

## *Phyllostethus clavigula*

3330. Bulbul. ♀. ovary somewhat enlarged. Stomach contents: fruit including red peppers.

*Platysteira cyanocephala*

3331. Red-wattled Fly catcher. ♀. o.n.e.

*Cecropis abyssinica gubernator*

3332. <sup>*Coracina gurneyi*</sup> Strike. ♀. o.n.e. Stomach contents: insect remains.

*Lathrotriccus tristis*

3333. Black Barbet with red face. ♂. t.n.e.

O

Paradje

Oct. 20. 1911.

3344. Black & white Kingfisher (*Ceryle*) ♂ testes

slightly enlarged. Stomach contents: fish.

3345. Goatsucker. ♂. testes slightly enlarged. Stomach  
*Caprimulgus natalensis chadensis*  
contents: some winged termites & pieces of beetles.

3346. Goatsucker. ♂. testes slightly enlarged. Stomach  
*C. natalensis chadensis*  
contents: many winged termites & 2 beetles.

Nos 3345 & 6, together with a long-tailed Nightjar (*Scotornis*) were capturing escaping termites on the wing, at dusk. Usually this species is perfectly silent, but recently I have heard a male utter, in flight, a short "kik."

Oct 21.

3347. Black Weaver with yellow back & wing coverts. ♂.  
t. e. *Coliusposseri macrourus*

Males in breeding plumage often exhibit their downy white breast-patch while perching to display their charms to the female they adopt an ungraceful jerky flight, spreading and swagging the tail up and down; but the voice is very weak. They rarely perch beside their mate, but the many rods away keep continual watch over her, for if she flies they immediately depart in the same

Faradje

Oct. 23, 1911.

*Campetherina permixta*

3351. <sup>♂</sup> Woodpecker. ♂. testes slightly enlarged. Stomach content

ants. Found in a swampy strip of woods along  
a brook; a common species to the south, in the  
forest region, but this is the first time I have

seen it at Faradje. Utters a "whining" note.

Later seen near midjan, midway between Faradje & Aba.

3352. Small Wood Hoopoe. ♀. ovary slightly enlarged.

*Scopelus aterrimus sinensis*

Stomach contained several berries and a few

pieces of insects. The note of this species may  
be written "whā, whā, whā, whā . . ."

and is uttered rather slowly, being heard  
generally in the morning.

*Camphagoides pheniceus*

3353. <sup>♂</sup> Barred Cuckoo Shrike. ♀. o.n.e. Stomach contents:

\$3070.

remains of insects.

*Dendropicos labeculatus*

3354. <sup>♂</sup> Small Brunish Woodpecker with streaked breast.

♀. o.n.e. Iris dark red, maxilla dusky, mandible mostly blue gray, feet green.

*Dendropicos leucotis*

3355. <sup>♂</sup> White-headed Barbet. ♀. ovary slightly enlarged.

Stomach contents - fruit.

Observed from Niangara to Aba, also at  
Vankerklovenville.

*Camaroptera griseoviridis*

3356. <sup>♂</sup> Green - gray Warbler. ♂. testes somewhat enlarged

*Neocharaxis cinctata*

3357. <sup>♂</sup> Weaver. ♂. testes somewhat enlarged. Stom. cont. seeds.

\$3071.

Paradise.  
Oct. 26. 1911.

species seems to be nearly, if not entirely, absent.

3366. Red-<sup>x</sup>-black Weaver (*Pyromelaena*) ♂. t. e. <sup>luminescens</sup>

3367 ✓ " " " " ♂. t. e.

3368 ✓ " " " " ♂. t. e.

3369. ✓ " " " " ♀. ovary somewhat enlarged.

A very common bird in the high grass. During the breeding season the males show their brilliant colors to great advantage in slow, horizontal flights, with flapping wings. At the same time they utter a "beady" note, consisting of fine chirps rapidly reiterated. This note is also given from a perch on a bush or grass-stalk, and resembles the sounds produced by 3 other related weavers, namely, the black widow bird, and the 2 yellow <sup>x</sup> black weavers represented by nos. 3334 & 3347.

3370 *Ragonoticta rara*. <sup>✓</sup> Weaver. ♀. o. e.; a completely formed egg - pure white - in oviduct. Crop contained small seeds.

3371. Flycatcher. ♀. im. *Cat's bella my angae*

3372 <sup>✓</sup> *Sitopra luteola* (Leht.) Small yellow Weaver with black face & throat.

♂. testes somewhat enlarged, but skull not

Paradise

Oct. 29. 1911.

fixed on the ground. To carrion they come in larger numbers. The corresponding in a way to the turkey buzzard of America, their powers of flight are vastly inferior.

Oct. 30.

3381. Kestrel. ♀. o. n. e. Stomach contents: a lizard's <sup>lizard</sup> tail. <sup>Cercosaura t. timorensis</sup> First kestrel seen this season. Here there is no kestrel during the rainy months as at Medje.

See 3386  
Oct. 31.

3382. Tiny Kingfisher (O. spidina). ♂. testes slightly enlarged. Stomach contents: a grasshopper. Rather noisy and active at present, as the mating.

3383. Harrier (Circus macrourus). ♀. ovaries not enlarged. Crop and stomach contained 3 lizards. Iris brown, cere greenish yellow, corners of mouth green, bill black with blue base, feet yellow, claws black.

3384. Kite (Ezanius caeruleus). ♀. ovaries not enlarged. Iris orange brown. Stomach contents: fur and bones of a rat.

Nov. 1: Platysteira cyanea myanmae

3385. Red-wattled Flycatcher. ♂. t. n. e..

Oct. 30.

3386. <sup>Procnias</sup> Small blue-breasted Weaver. ♂. testes slightly enlarged.

Faradje

Nov. 5. 1911.

brown. Shot in same places as no. 3391.  
*Sylvieta brachyura*

3396. Short-tailed Warbler. ♂. t. n.e.  
*Dryoscopella incana*

3397. Blue-gray Warbler with rufous patch on primaries.

♂ testes slightly enlarged. A characteristic bird of swampy woods, usually seen in pairs. Its notes are far from musical; they commonly resemble the words "three cheers," repeated monotonously, and sounding as tho the bird would like to whistle if he only knew how.

*Neothraupis capistrata*.

3398. Weaver. ♂. imm.

*Indicator indicator*

Nov. 6  
3399. Honey Guide. ♂. testes slightly enlarged. Stomach contained beeswax.

Early yesterday afternoon in a small wooded swamp close to the post, we came upon a male of this species, who at once started his excited chattering call, and soon flew off some distance, calling continually, and returning shortly to make sure we were following. We replied, according to the custom here, with low whistles, and were led out onto the higher ground, thru the tall

Paradise  
Nov. 6. 1911.

a tool to dig out the hive, and the birds, seeing that I stayed on the spot, retired noiselessly. A little while later I saw them again, for they were sitting with puffed-out breasts and open bills, uttering a low "chvee-r-r", which I had never before heard. They were apparently quarreling, for one shortly chased his ~~sister~~ off at top speed.

With the aid of burning grass and a long-handled native farming implement ("gita") my two boys soon had the hive unearthed, paying a penalty of only 6 stings; but, as we expected, the comb contained no honey, only a little orange-colored pollen (?), and many larvae. The hole was a chamber excavated by termites.

A shower of rain prevented me from seeing the return of the honey-guides, but we placed this reward in the forks of a tree, and returning an hour and a half later, found that they had been feasting. Early the next morning

Faradje

Nov. 6. 1911.

3400. Brown Fly catcher. ♀ imm. *Bradornis pallidus subalaris*
3401. Kite (*Elaeius*) ~~caeruleum~~ ♂ imm. Stom. cont.; fur of a rat.  
Nov. 8. *Pistilia phoenixoptera emini* <sup>3402.</sup> ~~33358.~~ <sup>AD</sup> leaver. ♀ o.e.; an egg (pure white) in oviduct  
Crop contained termites (wriggless).  
*Hirundo negevensis*
3403. Large brown-rumped Swallow. ♂. testes slightly enlarged
3404. " " " " " ♀ o.v.e.
3405. " " " " " ♀ immature.
3406. " " " " " ♀ immature  
(note trace of light spots on outer rectrices of no. 3406.)
- Nov. 9.
3407. Owl (*Strix*) ♀ ovary slightly enlarged. Stomach  
*Tyto alba affinis* contents: feathers and bones of a small bird.  
Shot, in the port, where I had been aware, for  
more than a month, from their excrement  
and pellets, that owls were accustomed to  
come at night. Their favorite perches were  
in a rubber tree with several stout horizontal  
boughs, and on some small beams that  
projected from the wall of a house, beneath  
the roof of the verandah. The first time I  
heard them make a noise was yesterday,  
when one uttered a hoarse "shshshsh--"

Faradje

Nov. 11. 1911.

*Lamprocolius chalcurus*

3412. Resplendent Starling. ♂ testes slightly enlarged  
Stomach contents: insects. Iris deep yellow, bill  
and feet black.

3413. Same as preceding ♀. ovary slightly enlarged  
Stomach contents: insects. ♀. *chalcurus*.

Nos. 3412 + 3 differ from nos. 3410 + 11 both in color  
and size. All four specimens were shot  
from flocks in a pasture, where they often  
alighted on the ground to feed. The larger  
species, in life, appears to have more brightly  
colored irides, and often makes considerable  
noise with its wings in flying. Their  
calls are somewhat alike.

Nov. 12.

3414. Gray-backed Wagtail. ♀. s. n. e. Evidently  
*Motacilla alba*  
a migrant from the north, for this is  
the only one I have seen since April last.

During the dry season it is rather common  
here.

Nov. 13.

3415. Resplendent Starling. ♂. t. n. e. Stomach  
contents: berries. *Lamprocolius chalcurus*  
S. 3412.

3416. Resplendent Starling. ♂ immature.  
S. 3412.

Faradje

Nov 14. 1911.

contents: berries and a large grasshopper.

3422. Black Barbet with red face. ♀ imm. o.n.e. Stomach  
contents: fruit & a large black ant.

- Nov. 15. 3423. Roller (Coraciidae). ♀. o.n.e. Stomach contained  
remains of a grasshopper, 2 dung beetles  
and other coleoptera. (Outer tail feathers not  
elongate.) Shot in a bushy pasture.

3424. Hoopoe. ♂. testes somewhat enlarged. Stomach  
(photo) Mupia somalensis.  
contents: pieces of a beetle's shell and an  
insect - larva resembling a large ant-lion.  
The hoopoe is here a bird of the dry season,  
the first one being observed this year on  
Nov. 9.

3425. Snipe (Gallinago). ♀ o.n.e. This particularly  
fond of swamps, they may also be seen  
from time to time in dry fields, as was  
the case with this specimen.

3426. Black Barbet with red face. ♂. testes slightly  
enlarged. Stomach contents: berries. Iris  
medium brown.

The characteristic note of this barbet is  
a dry "hic-cup, hic-cup, hic-cup,  
hic-cup, hic-cup-----", ~~after~~ introduced

Paradise

Nov. 16. 1911.

3434. Small Wood Pigeon. ♂. t.n.e. Iris dark brown, bill dark gray, corners of mouth yellow, feet blackish, soles green. Stomach contents: a small caterpillar and insects.

Rather common: climbs about in leafy trees and bushes. The notes which have already described as "whā, whā, whā—" are uttered by the males.

3435. Rhinoceros Bird. ♀. o.n.e. Iris bright orange, Buceros Africana. Outer portion of bill red, base of mandible and sides of maxilla at base bright yellow, feet dusky brownish. Stomach contents: several ticks, many short hairs, a small insect, etc.

Said to have been shot as it sat on a rhino's head, close to the ear. The rhino departed precipitately.

3436. Starling (Creatophora?) ♂. testes slightly enlarged. Stomach contents: berries. In this example the paired gular wattles are well developed, tho those of the grown are

Paradise

Nov. 17. 1911.

3444. Resplendent Starling. ♀. o.n.e. Stomach contents: small insects. *Lamprolius chalcurus*

3445. Black & white Kingfisher. ♀. ovary somewhat enlarged. *Ceryle rudis rudis*. Stomach contents: fish.

3446. Small Resplendent Starling. ♀. im. o.n.e.

Stomach contents: small insects. *Lamprolius calypatus chloropterus*

- Nov. 19. 3447. *Pardala phoenicptera emini*. ♂. testes slightly enlarged. Iris red. Crop contained grass seeds. S. 3358.

3448. *P. phoenicptera emini*. ♂. testes somewhat enlarged. " " Crop contained a termite.

3449. *P. phoenicptera emini*. ♀. ovary slightly enlarged. "

3450. Small black Weaver. ♂. t.e. skull not quite completely ossified. *Hypochera* sp?

3451. Green & yellow Shrike with orange breast band. ♂. t. n.e.

3452. Small yellow breasted Fly catcher. ♂. t. n.e.

3453. Fly catcher-Shrike. ♂. testes slightly enlarged; skull not quite completely ossified. Stomach contents: remains of beetles.

3454. Black Weaver with yellow wing-coverts & rump. ♂. testes much enlarged.

- Nov. 20. 3455. Purple Dove with black nuchal crescent. ♂. *Streptopelia vinacea*. t.e. Crop contained millet ("malu")

Absent during a large part of the rainy season. First heard calling this season on Sept. 4. Leaves about March.

Vankurchhovenville (= Nzoro))

Aug. 7. 1911.

On Aug 6, thirty colies and a pigeon were thus taken. Judged from their plumage and the color of the bill, 25<sup>of the colies</sup> were adult and 5 young. The pigeon was of the common<sup>large</sup> species with black nuchal mark. (*Streptopelia semitorquata*) On Aug 7, when I visited the place there were a dozen or more colies waiting outside, but only two were taken that day. Nos. 3045-9 I selected for the perfectness of their tails. Their stomachs contained nothing but red earth. I have taken a sample of this clay from inside the cave; and it has been suggested that it may contain salt, the to my tongue it has not a very saline taste.

Only a few yards away, in the same bank, is a large, but more open cavity, which the natives have covered with boughs so that the birds will go only to the one where they can be captured. Both, the natives claim, have been excavated by birds and animals; pigs and bushbucks being said to come there by night. They are not new, indeed the blacks claim that their fathers caught birds there, and that it is done continually. A similar hole, much frequented

Vankur  
Aug 7.

3050.  
22771.

3051

Aug

3052

3053  
1141

3054

Vanderkloovensville  
Aug 9. 1911.

(n. v.)

3055. Gray Hawk (Asturinula) ♂ t. n. e. Stomach contained remains of a rat and a small lizard.  
*Cornuella cornuta*
3056. Large streaked Brown Shrike with long tail ♀. Ovary slightly enlarged. Stomach contents: remains of black ants. (taken on the wing.)
3057. Same as preceding. ♂ immature. Stomach contents: remains of black ants.

Today's specimens were shot in the same field as those of July 30. There were 4 or 5 flying out from the trees at intervals to capture ants on the wing. In going to the village of Baduma we passed a field where there were 3 or 4 of the same shrikes, and while coming back, five days later, noticed half a dozen in the same place. On the other hand, in the many other clearings and plantations along our road we saw not a single example.

3058. Cuckoo. ♂. testis slightly enlarged. Stomach contained Cuculus jacksoni remains of caterpillars.
- Aug 10. 3059. Gray Warbler. ♂ t. n. e. Iris dark brown; maxilla black, mandible bluish gray with black tip; feet bluish. Stomach contained 3 small caterpillars and some insect remains.

Vander

3055

3061

3062

3063

3064

Faradje.  
Aug. 11. 1911.

3065. *Ostryphantes baglafecht* (Vieill.)  
Black masked Weaver. ♀. ovary slightly enlarged.

Iris light buffy yellow, bill black, feet pale pinkish  
buff.

3066. *Colius sp. macroura*  
Black Weaver with yellow wing-coverts & back. ♂. t.e.

Common everywhere in this region.

3067. *Pogonornis auricularis*  
Black Weaver, with back of head and a band  
on upper breast orange red, upper back yellow. ♂. t.e.

Quite a number seen around Vankerckhoven-  
ville and on road to Faradje. They perch and  
feed on the high grasses.

3068. *Sturnus sinensis*  
Shrike. ♂. t.n.e. Stomach contents: insects.

3069. *S. 2992*  
" Flycatcher-Shrike". ♂. t.n.e.; skull not completely  
ossified. Wings open

Evidently in search of insects these birds  
hop about in almost warbler-like style amid  
the boughs, not in the forest, but in the trees  
scattered about the more open country, especially  
in acacias.

Aug 12.  
3070. *Cuculus phoeniceus*  
An olive, gray, and yellow Cuckoo-shrike, whitish  
below, and barred throughout with black. ♂ immature.

Iris dark brown, bill & feet blackish. Stomach contents:  
insects. Sat on tops of low bushes in an open marsh,  
darting down from time to time after insects.

1 m

Faradje  
Aug 12. 1911.

3071. <sup>Aug 12. 1911.</sup>  
35354  
35355

3072.

Aug 11.  
3073

Aug 13.  
3074  
Aug 14.  
3075

3076

Aug 15.  
3077

Faradji.

Aug 18. 1911.

Faradji  
Aug 18

enlarged. Stomach contained about 20 small beetles.

3 roaches, and 2 leaf hoppers. Still flying aloft in numbers in the early evening.

Aug 8.  
3078.

Ground Hornbill (Bucorvus) Sex? Lores bluish gray, close to eyes dark blue, throat indigo blue, cheeks and lower portion of the throat rose madder, beak in front of eyes pinkish & gray.

Aug 17.  
3079.

Night Heron (N. leuconotus) ♂. testes slightly enlarged. Iris brown, lores and feet greenish yellow, streaked with light green beneath mandible. One of a pair found in the afternoon in trees on a small island in the River Dungu.

3080.

Buff-breasted Bee-eater with red throat. ♂. t. n.e.

3081.

Melitophagus b. frenatus ♂ t. n.e.

3082.

" " " " " ♀. o.n.e.

Common about the post & river.

Sarcops calvus

3083.

Paradise Flycatcher. ♂ immature. Bill dusky with light gray base, rim of eyelids dusky, iris dark brown, feet blue.

Sitapha taenioptera

3084.  
3063.

A Yellow Weaver with black face & throat. ♂. t.s.

Many of these weavers are nesting in the tall reeds and papyrus along the Dungu, but like many other members of their family they build a large number

1 m

Faradje

Aug 18. 1911.

3089. *Hypothymis ocularis*.  
Yellow Weaver with black line thru eye. ♀. o.m.e.

Stomach contents: insect remains.

3090. <sup>Dromocichla in canna</sup>  
Blue-gray Warbler with rufous patch on primaries.

♂. t.e. Stomach contents: small insects.

3091. <sup>Melanotis nigricans (Vieill.)</sup>  
Black Weaver. ♀. ovary somewhat enlarged. Stomach  
contents: insects.

With this number I have labelled a nest and  
a set of 2 eggs of this species, collected at Vankerck hov-  
enville on Aug. 8, as well as 2 odd eggs taken at  
the same spot, where there were a dozen or  
more nests attached to high stalks of grass.  
a very unusual situation. Each nest was  
hung between two of the higher flowering stems,  
at a height of 10 or 12 feet; three of them contained  
sets of 2 light blue eggs, and a fourth held 2 young.  
Many black males, and a couple of greenish females  
were observed.

Aug 19.

3092. Boatswain (*M. vexillarius*) ♂. testes slightly enlarged  
Stomach contents: principally small beetles, but  
also 3 small grasshoppers, 2 tiny leaf-hoppers, and  
a very small bug.

3093. Sandpiper (*Actitis hypoleucus*) ♀. o.m.e. Stomach  
contained maggots. The first migrants arrive from

Faradje

Aug 21. 1911.

*Heliolais erythroptera*

3105. Warbler. ♀ immature. Stomach contained the leg of a spider and a few insect remains.
3106. Pipit (*Macronyx*) ♀. imm. <sup>crocus</sup>. Stomach contents: insect remains.
3107. " " ♀ imm. " " "
3108. Small Bustard. ♂. testes slightly enlarged. Iris *Lissois melanaster*. light brown, rim of eyelids dull lemon yellow, top of maxilla blackish, rest of bill light yellowish gray, corners of mouth a little yellower, feet very light yellow. Stomach contained seeds & small beetles, and one large black dung beetle. (Body-skeleton also preserved). Alighted in an open spot in the grass, not far from the post. Has been seen near Niangara, at village of Denge.
3109. Gray Owl (*Bubo cinereus*). ♂ t.n.e. Stomach contained remains of many roaches and a spider.

Shot in a small clump of trees during daytime.

- Aug 22.  
3110. Slaty Gray Rail (Skeleton) ♂. testes somewhat enlarged. Skin saved for identification.
3111. Partridge. ♂ t.e. Crop filled with seeds and bluish berries.

Aug 23.  
3112. Fruit Pigeon. ♀ juv.

3113. Pipit (*Macronyx*). ♀ immature. St. cont. insects. <sup>crocus</sup>

Aug 24.  
3114. Bulbul with yellow under tail coverts. ♂. testes

slightly enlarged. Stomach contents: fruit. Notes are broken musical twitterings, and birds occasionally call with both wings raised abft. Seems to have no regular song.

Farey  
Aug 2

3115  
3116  
3117

3118  
3119

Aug 21  
3120

Aug 22  
3121

Aug 27. 1911.

clinging to the lower side of an oblique dead branch; and others were met with at Rungu and Paradje. They are often followed by 2 or 3 drongos, which try continually to peck at them, and attract one's attention by the rasping calls. Only one have I seen go down to the ground; usually they progress by repeated short flights, not allowing an approach to within gunshot; their pale plumage giving a false impression of size.

3122. Pigeon with black crescent on back of neck. ♂. testes enlarged. *S. semitorquata*  
Crop contained many round seeds, evidently of the kind so widely cultivated by the natives. At the present time these have been gathered in heaps, and left in the fields to rot and dry.
3123. Same as preceding. ♂. t.e. Crop contained 75(!) small red peppers, and 35 other small fruits.

The common large pigeon here; the big light gray wood pigeon *Columba unicincta* being found toward Dungu, and the smaller *Turturoena* <sup>in going toward the forest.</sup> being first met between Dungu and Niangara.



Paradje

Aug. 29. 1911

3133. Rail <sup>Sarothrura pulchra</sup> (Corethrura) ♂. testes slightly enlarged.

It is now certain that this rail does make the noise ascribed to it by natives, a sort of "gung-gung-gung-gung-gung-gung-gung...", as well as a shriller "kik-kik-kik-kik-kik...", uttered much more rapidly. The present individual produced both types of notes just before being shot, while walking along the muddy banks of a brook.

3134. Gray Warbler with rufous patch on primaries ♀.  
o.e. Stom. contents; small insects.

In almost every one of the little patches of woods in this vicinity, a pair of these warblers resides, finding their living by searching carefully the boughs of the trees.

Aug. 30. <sup>Indicator</sup> <sup>Indicator</sup> Parando River.

3135. Honey Guide, ♂. Led Mr. Lang to a bee-hive in a large tree. The nest, like that found yesterday at Paradje, contained no honey; but the following day 2 honey-guides were seen feeding on the comb.

Paradje.

3136. Widow-bird (<sup>area</sup> *Vidua*) ♂ t.e.; but skull not completely ossified.

3137. Rufous-crowned Swallow. ♂. testes somewhat enlarged.

Paradise

Sept 2. 1911.

3145. Pipit (*macronyx*) ♀ immature. Stom. cont.; insects <sup>croesus</sup>

3146. Coly. ♀. o.e. *Colius nigricollis leucophathalmus*

Between forking branches of a mimosa bush, about 5 feet from the ground. She had a nest, a bulky cup composed of shreds of bark, grass & weed stems, and a few dry leaves. When I first saw it, a couple of days ago, there were 2 eggs of a dirty whitish color; but one was broken since then. The remaining egg I have preserved (labelled no. 3146 a.)

On Sept 5, another nest was found in a mimosa bush, about 10 feet up, and containing 2 eggs. The nest & eggs have been preserved (no 3146 b)

At Aba on July 19, I took a coly's nest which I found in a small tree, also about 10 feet from the ground. It contained an egg and a newly hatched young coly. (no 3146 c)

The latter was preserved in alcohol, and shows natal down, barely 2.5 mm long, over whole upper surface, breast, and even abdomen, though

Sept 3. 3147. Widow-bird (*Vidua*), ♂. testes enlarged; skull not completely ossified.

3148. Widow-bird (*Vidua*). ♂. t.e.; skull not quite completely ossified.

Sept 4. 3149. Paradise Flycatcher. ♂ immature.

3150. Red-faced Barbet. ♂. testes slightly enlarged. Stomach

Paradjie

Sept. 4. 1913.

*Cisticola notatus malayoticus*

- 3156 Warbler, with grayish brown back & thick bill. ♂  
testes slightly enlarged Stomach contents: insect remains. A common bird in all the more open country, giving vent to his emotions, on the wing, by notes seemingly meant for a song. First he flies to a considerable height in the air, and then comes down obliquely, uttering a reiterated guttural <sup>strident or even</sup> gurgling sound. until, at the end he alights in a conspicuous position on some tree or bush. Similar notes may also be delivered from a perch.

3157. Small Brown Warbler, with dusky streaks on upper *Bradypterus abyssinicus* breast. ♂ testes somewhat enlarged. Iris dull brown; bill black, with base of mandible bluish gray, feet brownish. Stom. cont., remains of tiny insects
3158. Same as preceding. ♂ testes somewhat enlarged. *Bradypterus abyssinicus*. Stomach contents: remains of tiny insects

This same description would fit the song of the spotted Pycnonotus bird as well; but there the notes of the smaller species are much more prolonged than could be heard only when one is very close to or in the swamp. These species of the genus are loud enough to carry much greater distances, even several hundred yards.

A common inhabitant of papyrus swamps, where, during the rainy season, its notes are among the most characteristic sounds, tho heard less frequently during the hotter hours of the day. They consist of a sort of chirrup, repeated some 12 or 15 times, slowly at first, but then faster even separated by short intervals.

P-470  
Faradje

Sept 5. 1911.

3160. Bronzed Sunbird. ♀ juv.  *Cinnyris cupreus*  
*Sitagra taenioptera*.
3161. <sup>A</sup> Yellow Weaver with black face & throat. ♂ t. e.  
Stomach contents: seeds.
3162. Owl (*Scops capensis*) ♂ t. n. e. Iris yellow; bill dusky,  
but greenish beneath mandible, toes brownish,  
claws dusky.

Sept 6.

3163. Owl (*Scops capensis*) ♂ t. n. e. Stomach contained 2  
legs of a mantis, and a feather, perhaps from  
the owl itself.

No. 3162 was shot at dusk in some trees about  
a spring, where he had been heard calling for  
two evenings previous. No. 3163 was secured  
the following evening, only a few hundred  
yards away, where he was clinging to a  
tall grass stalk, in the open. The call  
which we have heard occasionally at dusk  
or during the night, is a single short note  
somewhat metallic, repeated at short intervals.

3164. Warbler. ♂. testes somewhat enlarged. Common.  
*Sitagra taenioptera*.
3165. <sup>A</sup> Yellow Weaver, with black face & throat. ♂ t. e.  
Stomach contents: seeds & insect remains.
3166. Small Blue-black Weaver. ♂ t. e. Crop & stomach  
hypochlora dialyptera but skull not completely ossified

Faradje  
Sept 8. 1911.

Stomach contents: insect remains.

Also a bird peculiar to the papyrus, where it goes sneaking about, uttering a low chip, or chuk, and sometimes, if one stands very still, coming very close. (Sings like the Little Papyrus Bird, but much louder; in fact this is the species more commonly heard.)

3178. *Linaria lepida* Green-<sup>+</sup> Gray Warbler with chestnut throat + shoulder-patches. ♂. testis slightly enlarged.

About Faradje this is a common bird in woods near water, as well as of papyrus swamps, with a conspicuous voice, producing loud and ringing notes, or even short reiterated whistles, but nothing really musical.

3179. *Tephrolepis viridis* Paradise Flycatcher. ♀ o.v.e. *Clinuspasianus coeruleo*  
3180. Black Widow Bird. ♂ immature (in brownish plumage) *Calamocichla violacea*  
3181. Gray Papyrus Bird. ♂ t.e.

This bird and his mate were very anxious about their nest, which was supported at a height of 11 feet in the papyrus, as shown by the sketch, being placed in the head of one of the rushes, and attached by its side to the stalk of another higher one. Its shape was that of a cup, while its materials

Faradje

Sept. 9. 1911.

*Anthreptes longirostris haemacephalus*

3187. white breasted Sunbird ♂. testes slightly enlarged.

3188. " " " ♀ o.m.e. Iris dark

brown; bill & feet dusky greenish. The first  
& have seen at Faradje in all our stay.

3189. Small Blue black <sup>Hypothrochlea</sup> W eaves ♂. t.e.; but skull not com-  
pletely ossified.

3190. " " " ♂ .., "

3191. " " " ♂ .., "

3192. " " " ♀. ovary somewhat

enlarged, but skull not completely ossified. Crops  
of all four contained tiny seeds.

Notwithstanding its lack of elongated tail  
feathers, this bird bears many resemblances  
to the little widow bird (*Vidua*). They both  
feed on the ground in small flocks, sometimes  
together; the females resemble each other  
considerably; and I have even seen a male  
of the black species hover in the air beside a  
perching female in exactly the same way as  
the widow birds do continually. The condition  
of the skull is likewise the same.

Sept 10.

*Pantholaea clericalis*

3193. Small Black Chat with white forehead & wing coverts.

♂ testes slightly enlarged. Stomach contents: insects

Faradje  
Sept. 11. 1911.

3202. Cong-tailed Goatsucker (Scotornis) ♂ t. n. e.

3203. " " " Scotornis ♂ " "

Shot as they alighted on the road before sunrise. After sunset, at the same spot, they fly about actively, but do not alight, nor have they begun as yet to give their loud "clicking" call.

3204. Spotted Papyrus bird. ♀. o. e.; an egg in oviduct,  
Bradypterus berpalus  
But shell not completely ossified.

3205. <sup>Hyphantornis ocularius</sup> Yellow Weaver with black throat and line thru eye. ♂ testes slightly enlarged.

Sept. 12.

Sitagra luteola (Licht.)

3206. Small yellow Weaver with black face & throat.  
♂ t. e. Iris bright brown, bill black, feet bluish gray, Crop contained seeds.

3207. Greenish Bee-eater with forked tail, yellow Dicrocerus hirundineus belgini throat and blue under tail-coverts and breast band. ♀. o. n. e. Iris red; bill black; feet dark gray.

3208. <sup>Acteropus sp.</sup> Two grayish Bulbuls. ♂. testes slightly enlarged

3209. Stomachs contained insects.

3210. Small Red-backed Shrike. ♂ testes slightly enlarged <sup>Eurystomus gularis</sup>

3211. Small Red-breasted Sunbird. ♂ t. e. Has

been observed several times recently in swampy

Paradip  
Sept. 12, 1911.

Paradip  
Sept 16

eggs, which were of an unusual light grayish blue color, without a spot. Their ownership was made certain by our seeing the female enter the nest twice.

3218. Heron (Ardea goliath) ♂. testes slightly enlarged.  
Stomach contents: fish bones. Iris yellow,  
orbita dark green; maxilla black, mandible  
blackish above, but light green beneath tip.  
and whitish below, posteriorly; feet black.

- Sept 15. 3219. Partridge <sup>G. interlynchus</sup> ♂ t.e. Crop contained ants, termites,  
a beetle, bits of grass, small fruit, and other  
vegetable matter.

3220. <sup>Lagonosticta rara</sup> Small Red-and-Black Weaver. ♂. testes somewhat  
enlarged. Crop contained small seeds & termites.  
The "song" of this species is a dry, musicless  
attempt at a trill, recalling the note of the  
chipper (Spizella socialis).

Sept 16.

3221. Small Green-backed Bee-eater ♀ o.m.e.  
<sup>Melittophagus pusillus</sup>
3222. Black-<sup>-</sup>-white Kingfisher (Ceryle) ♂ t.m.e.  
Stomach contents: fish bones.
3223. Least Bittern (Ardita) ♀. ovary somewhat  
enlarged <sup>Tayacai</sup> Iris dull orange red, shading to  
brown

Sept 17.  
3224

Paradjie

Sept. 17. 1911.

3226. *Ploceus superciliosus* (S. Bell) <sup>♂</sup>  
Weaver ♂. testes somewhat enlarged. Stomach contents  
seeds & tiny bits of stone.  
3227. *Ploceus superciliosus*.  
Same as preceding. ♀. o.e. Stom. cont.: seeds.

Sept 19.

3228. Owl (*Scope*) <sup>♂</sup> *cayensis* t.n.e. Stomach contained a few  
pieces of insects. Shot in same place as no 3162.

3229. Black--white Kingfisher (*Ceryle*) <sup>♂</sup> *radiata* t.n.e. Stomach  
contents: fish. Iris dark brown; bill & feet black.

3230. Wood Hoopoe <sup>♂</sup> t.n.e. Stomach contents: termites  
*Phoeniculus purpureus niloticus*  
& ants.

3231. Wood Hoopoe ♀. o. n.e. Stomach contents: termites  
*Phoeniculus p. niloticus*  
& stones of fruit.

3232. Partridge <sup>♂</sup> ovary slightly enlarged. Crop contained  
small seeds. The female sometimes mounts a tree  
while her mate is calling from aloft.

Sept 20. *Cisticola natalensis malgasci*

3233. Warbler. im. Stomach contents: small beetles, etc.

Sept 21.

3234. White-breasted Sunbird. ♂. testes very slightly enlarged,  
skull not completely ossified. Stomach contents: small  
spiders & a tiny beetle. *Antrocephalus longirostris haussarium*  
*Cisticola erythrops*

3235. Buff-cheeked Warbler. ♀. o.e. Very common here,  
in high grass and brush.

3236. Sandpiper. ♀. o.e. *Pisobia minute*.

Garadjie  
Sept 2

3237

Sept 22

3238

3240

Sept 24  
324

3241

3242

3243

3244

Faradjé

Sept 25. 1911.

3246. Hornbill (Bucorvus) ♀ o.w.e. Stomach contained a few insect remains. Naked skin of orbits & throat blue.

Sept 26.

3247. Goatsucker (M. vexillarius) ♀ ovary somewhat enlarged. Stomach contents: small beetles, including some dung beetles, a few bugs, and an earwig.

They no longer fly about high in the air; indeed since Sept 3rd I have not seen a single male.

3248. <sup>23138</sup> Goatsucker. ♂ testes slightly enlarged. Stomach contents: a few remains of beetles.

Sept 27.

3249. yellow-breasted Finch. ♀ o.e. Serinus mozambicus barbatus.  
3250. Goatsucker (Scotornis) ♂ testes slightly enlarged. Stomach contents: a few pieces of beetles.

3251. <sup>23248</sup> Goatsucker. ♂ [im.] t.n.e. Stomach contents: a dung beetle, a bug, & a green leaf hopper.

Sept 28.

3252. <sup>23251</sup> Goatsucker. ♂ testes slightly enlarged. C. natalensis chadensis. Members of this species, as well as of Scotornis, are now occasionally seen to alight on the ground at dusk.

- 3252 a. Cinnyris cupreus. ♀. ovary enlarged

Faradjé

Sept. 30

3253

3254

3255

3256

3257

3258

3259

3260

Paradje

Oct. 1, 1911.

light gray ground color, marbled with brown and spotted with dark purplish brown, a few scravls of the same dark color being also present. The female bird was caught in the nest at dusk.

3261. Gray Owl (Bubo cinereus) ♂ t. n. e.

Oct. 1. Plectroptera cyanocephala <sup>griseo-</sup>

3262. Red wattled Flycatcher. ♂ t. n. e. (Cnemoptera <sup>griseo-</sup> viridis)

3263. Green & gray Warbler. ♂ imm. Heard occasionally

Oct. 3. about the post. Common in the larger patches of woods

3264. Weaver. ♀. ovary slightly enlarged.

3.3214

3265. Creeper. ♂. testes slightly enlarged; but skull not completely ossified. Iris dark brown, bill dusky brown, grayish below, feet grayish brown. One of a pair that were climbing about on some large trees in a swamp.

Oct. 14 Nos 3266 to 3290 incl. were collected in and about a patch of forest some 3/4 hour S. of rest house Mordjan or Marapili, midway between ~~aba~~ <sup>BB</sup> & ~~Peru~~ <sup>Peru</sup>.

3266. Black-headed Oriole. ♀. ovary slightly enlarged.

3267. Golden Oriole. ♀. o. n. e. Iris red. Stomach

Oriolus auratus

contents: a caterpillar. Our first 2 specimens were collected at Paradje on Apr. 26. On May 1<sup>st</sup> they were numerous, but no more were noticed until Aug 17, when a single example was seen in a tree near the R. Dungu. On Sept 12, 3 more were observed, and now in early October, they are again common.

Faradji  
Oct. 5. 1911.

3275. Cuckoo. ♀. o.n.e. Stomach contents: a slug, *Centromerches a. intermedius*, a large grasshopper, and pieces of a smaller grasshopper and beetles.
3276. Shrike (*Laniarius*) ♀. o.n.e. *sueleureopeltis*
3277. Shrike (*Malacorhynchus*) ♀. inv. Iris orange. Stomach contents: a large bug. *polic. ecarinatus*
3278. Small Gray Woodpecker. ♂. t.n.e. Stomach contents: small white grubs. Iris dark red, bill dusky with gray base, feet dark grayish green.
3279. *Hypargos dybowskii*. Small gray Weaver, with red back, and white spots on sides. ♀. ovary somewhat enlarged. Crop contained seeds.
3280. Red-crested Plantain-eater (*Musophaga*) ♀. *rossae*. ovary slightly enlarged. Stomach contained fruit. Shot in a patch of forest about midway between Faradji and Aba, where I also heard the call of the great Plantain-eater (*Corythiols*). This *Musophaga* has never been noticed close to the post of Faradji, probably because there are no suitable woods; but between Dungu and Niangara they appear to be common. Its notes have some resemblance to those of *Durucus*, but without such regular

Faradje

Oct 6 1911.

*Cercus pectoralis*

3288. Light gray Cuckoo shrike. ♀. o. n. e. Stomach contents: 3 caterpillars (hairy).

Was hopping actively about in the boughs of a high forest tree.

3289. <sup>3270.</sup> *Phylloscopus scandens orientalis* Bullbul. ♂. t. n. e. Stom. cont.; 3 small caterpillars (hairy), and insect remains.

3290. Same as preceding. ♀. o. e. Stom. cont.; a small green caterpillar and insect remains.  
[nos 3266 to 3290 incl. were collected in and about a patch of forest some 3/4 hour S. of fruit house mudjan or marapili, midway between Atba & Faradje]

3291. Black Widow-bird. ♂. t. e. *Colius caeruleocephalus concolor*

3292. " " " ♀. o. e.

3293. Red- & black Weaver (*Pyromelana*) ♂. testes much enlarged.

3294. Weaver. ♂ im. t. n. e. Stom. cont.; seeds & small stones.

3295. Small Papyrus Bird ♂. t. e. Stom. cont.; tiny insects.  
*Bocagia glycinivora*

These birds have nearly quit singing altogether. The main season of song comprises the months of June, July, August & September.

3296. Small Brown Fly catcher. ♀. ovary slightly enlarged.

3297. Young of 3296. ♀ im.

Shot in a papyrus swamp, 5 hrs. E of Faradje.

P-490  
Paradjie

continually to another spot in the post where men were accustomed to urinate.

Oct 11. 1911.

3309. Coucal. ♀ juv. Stomach contents: remains of (*Centropus nigerinus*) <sup>large</sup> grasshoppers & beetles.

3310. Warbler ♂ testes slightly enlarged.  
*Phacellodomus sp.*

3311. Thrush. ♂ immature.

3312. Shrike (*Elephantulus*) ♂ immature. A very common bird here.

3313. Red- & black Weaver. ♂. testes slightly enlarged.  
*Pyrenestes ostrinus* A bird of the swamps, not very common.

- Oct 12. 3314. Shrike (*Malaconotus*) ♂. testes slightly enlarged.  
Stomach contents: a large dragon-fly.

3315. Shrike (*Malaconotus*) ♂ testes very small.  
(Skull completely ossified, inspite of gray bill) Iris deep yellow. Stomach contained several large paper-wasps.

The peculiar protracted whistle of this shrike, not loud, is often introduced by one or two abortive attempts, only audible at close range; and is heard most frequently in the early morning. The birds hope about in low trees and bushes, and seem fond of the rubber groves (manihot).

3316. Gray Plantain-eater (*Schizorhynchus*) ♂ t. n.e.  
*Schizorhynchus zonurus* Stomach contents: fruit. Besides its character-

Paradje.

Oct. 14. 1911.

amid the high grass and bushes, for often while watching for goatsuckers at dusk, I have seen flocks of 70 to 100, easily identified by their nasal calls, come down and fly anxiously around, with much whish of wings, before settling down for the night.

*Sitagra taenioptera*.

3319. <sup>A</sup> Weaver. ♂ imm. t.m.e. Iris light gray.

*Yungipicus obsoletus*

3320. Small Gray Woodpecker. ♀ o.m.e.

3321. White-eye. ♂. testes much enlarged. *Zosterops superciliosus*

*Calamonastes niloticus*

3322. Papyrus bird. ♀ juv. Secured, of course, in a papyrus swamp. My attention was drawn to it by its plaintive note, quite distinct from the voice of an adult.

*Euphonia lata incana*

3323. Blue-gray Warbler, with rufous patch on primaries. ♂. testes slightly enlarged; skull not quite completely ossified.

3324. Shrike (*Pelecanoides*). ♀. o.m.e.

*Pachycephala flavigularis*

3325. Bulbul. ♂. testes slightly enlarged. Stomach contained small fruits, the intestine the seeds of red peppers.

3326. Swallow (*Psalidoprocne*) <sup>magnirostrum</sup> ♂ immature.

Paradje

Oct. 15. 1911.

3327. <sup>A</sup>

3328.

3329.

3330.

3331.

3332.

3333.

## Paradjie

Oct. 18. 1911.

3334. Black Weaver with yellow wing-coverts and rump. ♂  
Coulectes crossirostris.  
 testes much enlarged. Common about Paradjie.  
*Bradornis pallidus subalaris*
3335. Brown Fly catcher ♂. t. n. e.  
*Anomalospizixanthops butleri*
3336. Small yellow Weaver, ♂. testes slightly enlarged;  
 skull not completely ossified. Iris hazel; bill &  
 feet dull brown, rather light. Stom. cont.: seeds.  
 First I have ever seen; was sitting on a small  
 tree in the high grass.
- Campetherina palica*
3337. Woodpecker with red moustaches. ♂. t. n. e. Stomach  
 contents: little brown ants and small termites.
- Cinnyris sericeus* + +  
 3338. Tiny Sunbird. ♂ imm. Stom. cont.: remains of tiny insects.
- Cinnyris erythrocercus butleri*
3339. Widow-bird (*Vidua*) ♂ imm. (juvenile plumage)  
*aniceps*
- Oct. 19.
3340. Whitecrested Plantain-eater, ♀. ovary somewhat  
*Turacoides leucolophus*  
 enlarged. Stomach contents: a rather large snail,  
 and pieces of its shell.
3341. Snipe (*Gallinago*). ♂. t. n. e. Pretty numerous  
*media*  
 in open muddy places in swamps at the present  
 time.
- Oct. 20.
3342. Cuckoo (*Chrysococcyx*). ♂. testes somewhat enlarged,  
*Coprinus*  
 stomach contents: caterpillars. Rather common here.
3343. Large rufous-breasted Kingfisher (*Ceryle*) ♂ t. n. e.  
*maxima*.  
 Stomach contents: fish. Shot along the Dungee.  
 Its loud harsh notes are heard from time to time  
 at the post.

Faradje

Oct. 21. 1911.

direction, sometimes approaching to perform a dance in mid-air, and then alighting again at a little distance.

3348. Same species as preceding. ♀ immature.

3349. *Sitana taenioptera*.  
Yellow Weaver with black face & throat. ♂.  
t.e.

On Sept 16. 1911, 5 sets of eggs of this species were collected from nests in the tall reeds fringing the bank of the R. Dungu at Faradje, each set consisting of two. So the different sets exhibit very great variation in color, the two eggs from the same nest are in every case very much alike. (These eggs all labelled 3347). These weavers are still nesting (Oct 21); lately I have noticed a small tree close to the water, on the lower branches of which they were building, the their nests are more frequently placed in reeds & papyrus.

Oct. 23

3350. Honey Guide. ♂. t.m.e.

Faradjie

Oct. 25, 1911.

3358. *Ptilia phoenicoptera enni*. Hart.  
A Gray Weaver, with barred belly, and red wings & tail.

♂. t. e. Iris red; bill blackish, with a little blue at base, feet light brown. Crop contained grass seeds.

3359. *Ptilia phoenicoptera enni*.  
Mate of 3358. ♀. o. e. Eyes, bill & feet colored as in ♂. Crop contained grass seeds.

This pair of weavers, the first of their kind I have seen, were feeding on the ground on a road and in a field of sesame. The body of the female was larger than that of the male.

3360. *Campetherabalia*  
Woodpecker with red "moustaches". ♂. testes slightly enlarged. Stomach contents: small ants & their larvae.

3361. *Campetherabalia*  
Mate of 3360. ♀. ovary slightly enlarged. Iris pink, bill dusky, feet green. Stomach contents: small ants & their larvae.

3362. *Grenonela elegans*  
Small yellow-breasted Warbler with gray crown and white throat. ♂. immature.

3363. Red- & black Weaver (*Pyromelaena*). ♂. t. e.  
*Bucalopha clericalis* Glaucomelas

3364. Small Black Chat, with white forehead &

- Oct. 26. wing-coverts. ♂. t. e. Stom. cont.: insect remains.

3365. Tiny Sunbird. ♂. t. m. e. *Cinnyris osea böhmi*

During the greater part of the rainy season this

Paradjie.  
Oct. 26. 1911.

quite completely ossified. Stomach contents: seeds.  
S. ~~Same as~~ <sup>Same as (left)</sup> 3373 ~~State of~~ 3372. ♀. o. e. Stomach contents: seeds.

A nest of this species was found in a mimosa tree, at a height of only 10 feet. It contained 2 pure white eggs. Deserted nests are common in the small mimosas, and have long spouts, which are probably built during the period of incubation or the rearing of the young.

Oct. 27.

3374. Small Resplendent Starling. ♂. im.

Stomach contents: insect remains. *Lamprocolius*  
*lubricus* *leucus*  
*leucostethus* *chloropterus*

3375. Black Barbet with red face. ♀. o. n. e.

Stomach contents: fruit.

3376. Bee-eater. ♂ t. n. e. Five individuals seen today.  
3377. " " ♂ t. n. e. |

3377. " " ♂ t. n. e. | the only ones since Sept 12.

3378. " " ♀. o. n. e. |

3379. Rufous-crowned Swallow. ♀. o. n. e.

Oct. 29  
3380. Vulture (*Necrophorus*). ♂. testes slightly enlarged.  
*Necrosyrtes monachus*. Present during the whole year. In groups of 3 to 6 they are commonly seen high in the air, proceeding with slow, flapping flight, or circling about with thin gaze

Paradise

Nov. 2, 1911.

3387. Fruit Pigeon. ♂. t. e. Crop contained berries.

An egg of this species was taken today from a frail nest of small twigs placed on a bough projecting out over the river, and only about 7 feet above the water. The present specimen was sitting on an empty nest, in an adjoining tree.

Nov. 3.

3388. Wattled Lapwing. ♀. ovary slightly enlarged.

~~Africypus schlegelii~~

Yesterday this species was observed for the first time since June.

*Nyphargus monteiri*.

3389. <sup>Nov. 5.</sup> Weaver. ♂. t. e.

*Sylvietta brachyura*

3390. <sup>Nov. 5.</sup> Short-tailed Warbler. ♀. o. m. e.

3391. Large red & black Weaver with thick bill. ♂.

*Pyrenestes ostrinus*. Testes somewhat enlarged. Crop contained seeds.

Shot in small swampy wood.

3392. Small red-breasted Sunbird. ♂. t. e. <sup>Cinnyris</sup> *chloropygius*

<sup>Nov. 5.</sup> *Neopelmica goetzei poliocephalus*

3393. Red-rumped Woodpecker. ♂. testis slightly enlarged. Iris dark brown.

3394. <sup>Nov. 5.</sup> Mate of preceding. ♀. ovary slightly enlarged. Iris dark brown.

3395. Weaver. ♀. ovary slightly enlarged. Iris dark

<sup>Nov. 5.</sup> *Pyrenestes ostrinus*.

Faradje

Nov. 6. 1911.

grass and scrub. Now our feathered guide would fly noisily ahead of us till out of sight (50 yards or so) and then perch on the top of a bush till we came up; and then repeat the performance. After a few hundred yards we found that another honey guide, also a male, had arrived. We had gone thus about 600 yards when both birds stopped in a small tree, hardly worthy of the name, and plainly incapable of harbouring a hive. Nevertheless the indicators showed plainly, by taking short, <sup>aimless</sup> flights and returning always to the same spot, that this was the place. They were very tame, too, allowing an approach to within 10 or 15 feet. At first we were doubtful, but looking about, we heard the buzzing of bees, and found that they were going into a small hole in the ground, close to the base of a small scrubby tree ~~stood~~ to that on which the honey guides sat. My boy now went to fetch matches and

Faradje

Nov. 6, 1911.

I went to the spot and sat down to watch. One of the birds soon came silently to the tree, climbed cautiously down, and after a little pecking, seized a piece of the comb in his beak and flew off. After a short wait, his fellow arrived in the same timid fashion. That both birds were present is proved by the return of one after I had shot no. 3399.

This honey guide is rather common here, and said to be more so toward the north. Near Yakuluken, the Azande, who call it "Purubwa", told Mr. Lang that in former times this bird was very strictly protected, and that any native stupid enough to kill one would have been punished with the loss of an ear.

Para  
Nov.

34

34

Nov.

34

34

34

34

34

34

34

34

34

34

(M)

Nov.

34

Faradje  
Nov. 9. 1911.

3408. Widow-bird (*Stegannura* ?) ♂. t.e.; skull not completely ossified. Iris dark brown, bill black, feet dark brown. Stomach contents: seeds.

With the exception of a single individual seen at Niangara in November of last year, this is the first I have noticed. Unlike the two common widow-birds, this species makes long protracted flights, sometimes as far as  $\frac{1}{4}$  mile, and often attaining a height of 150 feet. Moreover they are usually alone, <sup>and silent,</sup> and alight on the tops of the trees, where they can be approached only with difficulty. The present example was secured in a bushy pasture, where another was observed a few days later.

Nov. 10.

3409. Wattled Lapwing with white forehead. ♂. t.w.e.  
*Africayx s. senegalensis*. Rather numerous at present, feeding in tilled ground and pastures, and becoming particularly active toward dusk.

Nov. 11.

3410. Small Replendent Starling. ♂. t.n.e.

Stomach contents: insects. *Lamprolaimus chalybaeus chloropterus*

3411. Same as preceding. ♀. ovary slightly enlarged.  
Stomach contents: insects. *L. chal. chloropterus*

Paradise

Nov. 13. 1911.

Paradise  
Nov.

3417. Starling (*Creatophora*,?) ♂. testes slightly enlarged. <sup>carunculata</sup>  
iris dark brown, bill pale pinkish white.  
naked cheeks yellow, naked skin of throat  
bluish gray, wattles on crown dark gray,  
feet brown. Stomach contents: berries  
and a tiny snail.

One of three or four found, <sup>for the first time</sup> in a pasture  
with resplendent starlings.

- Nov. 14. 3418. Gray Heron. ♀. o. n. e. Stomach contents:  
remains of grasshoppers and other insects.  
This heron seems to occur here only in  
the dry season, from November to about  
April, when it is common, often feeding  
on the ground in very dry spots, such  
as old fields, tho' it likes water too.  
The call is loud and harsh.

3419. Starling (*Creatophora*,?) ♂. testes slightly enlarged. <sup>carunculata</sup>  
Stomach contents: berries.

3420. Starling (*Creatophora*,?) ♀. o. n. e. Stomach contents: berries & a grasshopper. <sup>carunculata</sup>  
Irides dark brown, cheeks yellow, naked skin  
of throat gray, bill pale pink, feet brown.

3421. Starling (*Creatophora*,?) ♀. o. n. e. Stomach <sup>carunculata</sup>  
contents: berries.

Paradise

Nov. 15. 1911.

by a medley of harsh sounds. A single nasal "caw", strongly suggesting that of *Corvus ossifragus*, tho' not so loud, is also uttered.

*Yungipicus obsoletus*

3427. Small Gray Woodpecker. ♂ t. n. e. Irides dark red.

3428. Black Titmouse with white wing patches. ♂.  
*melanifacies niger leucomelas*  
t. n. e. Irides yellow. Stomach contents: some small caterpillars & beetle larvae.

3429. Black Titmouse with white wing patches. ♀.  
o. n. e. Irides yellow. Stomach contents: several small naked caterpillars.

3430. Resplendent Starling ♂. testes slightly enlarged  
5.3412. *Campephilus chalcopterus*

Stomach contents: a berry and pieces of insects.

3431. Starling (*Creatophora*?) ♂ imm.

Nov. 16.

3432. " " "

♀. ovaries slightly enlarged.  
Stomach contents: berries

3433. " " "

♀. o. n. e. Stom. cont. a little fruit & pieces of a cricket.

Members of a flock of 12 or 15, found in the same pasture as no. 3417. Their habits are like those of the resplendent starlings, with whom they mix, both kinds often alighting on the ground.

Faradj  
Nov. 16. 1911.

small. The gular wattles are also present in no. 3432, a female, tho practically absent in 3417, an adult male.

Nov. 17. 3437. Widow-bird (Steganura <sup>♂</sup> paradisea acutiprum) ♂ imm. testes somewhat enlarged. Crop contained grass-seeds. Iris dark brown, bill black, feet dark brown.

3438. Widow-bird. (Steganura <sup>♀</sup> paradisea acutiprum) slightly enlarged. Stomach contained seeds.

Members of a party of four, sitting on the dry bushes in a recently-burned area. All were in brown plumage. Since then I have seen birds in this plumage feeding on the ground.

3439. Black Titmouse with white wing-patches. ♀. Melaniparus niger leucomelas o. n. e.

3440. Pipit. Anthus trivialis ♀ imm. Stomach contained small seeds & insect remains.

3441. Hirundo smithii Rufous-crowned Swallow. ♂. testes slightly enlarged.

3442. Mate of 3441. ♀. imm. o. n. e.

3443. Small Resplendent Starling. ♂. t. n. e.  
L. chal. chlorosternus

Paradise

Nov. 20. 1911.

*Vidua serena*

3456. Black - white Widow-bird. ♂. t. e.; skull not quite completely ossified. Crop contained tiny seeds. (Post-nuptial molt just commencing)

3457. Blue-gray Falcon. ♀. ovary slightly enlarged.  
~~Dissodectes abdorsicrus~~.  
 Stomach contained a few pieces of butter. Iris dark brown, orbits cere & corners of mouth yellow, beak horn-blue with black tip, feet yellow, claws black with bluish gray bases.

3458. Weaver. ♀. ovary slightly enlarged; skull not completely ossified.

3459. Brown Fly catcher. ♂. t. n. e. *Bradornis pallidus subalaris*

3460. Nightingale. ♀. o. n. e. skull not quite  
~~Luscinia megarhynchos~~ completely ossified. Evidently a migrant from the north; arrives early in November, skulks in dense bushes, giving harsh notes and a somewhat broken song. *Leads in March*

Nov. 21.

3461. Hoopoe. ♂. testis slightly enlarged.

*Muppa somalensis* <sup>new</sup>

3462. Widow-bird (*Vidua*) ♂ im. testes somewhat enlarged. Not an adult molting out of breeding plumage, for the brown feathers are older & more worn than the black.

3463. Chat. ♂. t. n. e. Stomach contents: seeds & remains of insects. (*Saxicola rubetra*)

Paradise

Nov. 21

3464

Nov. 2

3465

See J. Ba  
Natural H

XI, 1913

p. 193

3466

3467

3468

No. 3

Paradise

Nov. 21. 1911.

*Campethera phoenicea*

3464. Barred Cuckoo-shrike. ♂ imm. Stomach contents:

small naked caterpillars and a bug.

Nov. 22.

3465. Roller (*Coracias*) <sup>♀</sup> o. n. e. Stomach contents:

green grasshoppers. These long, slender

grasshoppers are very fatty, and highly  
of genus *Homoconyphus*

See J. Bequaert,  
Natural History  
XXI, 1921,  
p. 193.

estimated by natives as food. Today they

were very numerous in an open grassy  
situation not far from the post, and

a number of natives, were engaged in their  
<sup>(title Dog)</sup> capture, while a flock of kites (*milvus*

<sup>sextans</sup> *aegyptius*) hovered about, swooping

down from time to time, catching the

flying insects in their talons, and

eating them on the wing.

*Sylvia subalpina?*

3466. Warbler. ♂ t. n. e.

*Pytilia phoenicopeplus emini*

3467. Weaver. ♂. Testes slightly enlarged. Iris red.

Stomach contents: grass-seeds.

*Cisticola lugubris*

3468. Warbler. ♀. o. e. Stomach contents: remains

of a small green grasshopper & other insects.

Shot in a grassy marsh

*carunculata*

- Nov. 23. 3469. Starling (*Creatophora*) <sup>t</sup> ♂. Testes slightly enlarged

Stomach contents: berries. (Gular wattle

small. Of the wattle on the crown, the anterior

Paradise

Nov. 23. 1911.

is placed transversely, the posterior longitudinally.

3470. Resplendent Starling. ♀ imm. Stomach contents.

S. 3472.

*Lamprocolius chalcous*

berries. (This species was observed at Aba in Dec.)

3471. Starling (*Creatophora*?) ♀ imm. Stomach contents: small termites & berries.

3472. Creeper. ♀. ovary slightly enlarged, but skull *Dalporus salvadorensis* not completely ossified.

S. 3473.

*Chalcotrichia senegalensis* *acuta*

Black Sunbird with large red breast patch. ♂. testes slightly enlarged. Stomach contents: a leaf-hopper, other tiny insects, and a small spider.

3474. Streak breasted Kingfisher. ♀. ovary slightly enlarged (*Halcyon chelicuti*)

Especially noisy at dawn and sunset.

its note being characteristically divided into two parts, and often repeated. In spite of considerable variation in its length, it may usually be represented as

"tirrrrrrr — rrubb!" Like *H. senegalensis*, when two individuals perch close together they may sit <sup>for a time</sup> with outspread wings and reiterated calls.

3475. Starling (*Creatophora*?) ♂. testes slightly en-

*coronata*

Paradise

Nov. 23. 1911.

larged. Stomach contents: fruit.

3476. Small Wood Hoopoe. ♂. testes slightly enlarged.  
*Scopetes sternulus cinnamomeus*.  
Stomach contents: a caterpillar.

Nov. 22.

3477. Hawk (*Melierax*) ♀. ovaries not enlarged.  
*m. metabates*.

Stomach contents: a lizard and several of its eggs. Occurs here in dry season. First seen today.

Nov. 24.

3478. Quail (*Excalfactoria*) ♀. ovary somewhat enlarged  
Iris reddish brown, maxilla dusky, mandible bluegray, feet yellow. Crop contained small hard seeds. Shot on the edge of a marsh where the grass had been burned almost 3 months previously, and a new growth of green grass, not yet waist high, had sprung up. It could be flushed only with difficulty, tho when on the wing it flew strongly and directly, with swift moving wings, very like *Coturnix*.

3479. <sup>Sarothrura (böhmii?)</sup> Red-headed rail. ♂ testes slightly enlarged.  
Stomach contents: small seeds. Shot near same spot as no 3478.

3480. Long-tailed Boatsucker. ♀. ovary slightly enlarged.  
*Scutomis elongatus*. Stomach contents: 13 small dung-beetles.

1.520

Paradise

Nov. 26. 1911.

*Young Picus obsoletus*

3481. Small Gray Woodpecker. ♀. ovary not enlarged.

Stomach contained a small caterpillar and remains of insects.

*Ptilia phoenicoptera enina*

3482. <sup>a-3358.</sup> Weaver. ♀. ovary somewhat enlarged. Crop contain small termites. Common at present about roads and cleared or burned places. (Also common at Aba in Dec., even on the hills.)

3483. Stone Curlew. ♂. t.m.e. feet grayish green. Stomach contents: dung beetles & remains of a few other insects. This bird first began to call again in the evening in early November. I had not seen one since June, and am not sure that they remain here during the rainy season. At present they are rather numerous and noisy about the post during the evening.  
(Mordjan)

3484. Least Bittern. ♀. ovary slightly enlarged.

*Ixobrychus pagei*  
Sitting on a tall grass-stalk in a swamp, at sunset.

*Dryocopus lineatus*

3485. Shrike. ♂. testes slightly enlarged. Stomach contents: a caterpillar and remains of insects.

Nov. 27.

3486. Large Black & white Hornbill. ♀. ovary slightly enlarged. *Buceros subcylindricus*. Iris medium brown, lighter around outer edge, orbits whitish; softer skin on feet

Paradjie (mordjan)

Nov. 27. 1911.

In Barnes' motion pictures from S. Africa, oxpeckers are seen on Hippos and on Oryxes. In Akeley's Chicago groups, I believe, they are shown on certain antelopes.

In 1937 I saw yellow-billed oxpeckers on buffalos in Rietshorn River - TPC

Iris red, bill black, feet purplish pink with dusky brownish claws. Stomach contents: a brown grasshopper and remains of other insects. A forest bird; the first example seen since medje. While perching it wags its tail from side to side.

Paradje (Mordjan)

Nov. 27. 1911.

blue gray, soles yellowish, large scales and claws blackish. Stomach contained fruit.

3487. Rhinoceros Bird. ♂. t. n. e. Iris orange, outer half of Beakage orange. Bill red, basal half yellow, feet dusky brown. Stomach contained many short hairs and some small slender seeds.

One of three that came back to a buffalo after it had been shot. These birds accompany both rhinos and buffaloes, and are said to be very wary, their cries at the approach of a man alarming their hosts. They do not come to cattle here, as far as I can ascertain.

3488. <sup>nicobar shrike</sup> Woody Shrike ♂. t. e. Stom. cont.: a grasshopper

3489. " " ♂ t. e.

3490. <sup>Megabyas atricapillus</sup> Fly catcher with black back and white rump and under surface. ♂. Testes somewhat enlarged. Iris red, bill black, feet purplish pink with dusky brownish claws. Stomach contents: a brown grasshopper and remains of other insects. A forest bird; the first example seen since Medje. While perching it wags its tail from side to side.

Paradjie (Mondjau)  
Nov. 27. 1911.

Sunday  
Nov. 27. 1911

3491. Same as preceding. ♀. ovary slightly enlarged. Iris red. The female also wags her tail from side to side.

3492. A brown thrush-like bird (Timaliidae?).  
Turdinus  
♂. t. e. Iris brown, maxilla dusky, mandible blue gray, feet flesh color. Stomach contained a spider.

3495.

A bird of the underbrush in forest. Its song, heard commonly during the very early morning, consists of 6 semi-musical notes uttered in rapid succession, the whole being repeated at short intervals.

Rather common in a wood midway between (in village of Marapili or Madrapili) Paradjie and Aba where it was heard in early October as well.

3496

Staphidae vulpina

3493. ♀. ovary slightly enlarged. Stomach contents: a small snail & insect remains. A forest bird; as might be expected from its broad bill and short feet, its habits are those of a flycatcher. The song is a whistled "tritretretretretre...."

Nov. 2

3497

3498

3499

3494. Cossypha <sup>notabilis</sup>. ♀ juv. Iris dark brown; bill dusky brownish, corners of mouth yellow.

Faradje (Mordjan)

Nov. 27. 1911.

feet flesh color. Stomach contents: insect remains and shell of a tiny snail.

Adults of this species have been heard to imitate the low "tut-wè" call note of the guinea fowl (*Numida*), and also the notes of the yellow-headed oriole such as no. 3267.

3495. <sup>Campetherina caroli</sup> Brown-cheeked Woodpecker. ♀. ovary somewhat enlarged. Stomach contents: small black ants and their larvae.

One of a pair found in the woods near the rest-house of Mordjan. This is a forest bird, and has never been noticed near the post of Faradje.  
<sup>Audubon's</sup>

3496. <sup>♀. o. n. e.</sup> skull not quite completely ossified. Iris red. Stomach contents: insects.

Nov. 28. 3497. Kingfisher. ♀. o. n. e. Stomach contents: 2 large <sup>Halcyon</sup> *sengalensis* mantis & a green beetle.

3498. <sup>*Nyphantopus nigricollis*</sup> Weaver. ♂. imm. Stomach contained insect remains.

3499. <sup>*Nyphantopus nigricollis*</sup> Weaver. ♂. imm. Stomach contents: insect remains

3500. <sup>*Nyphantopus nigricollis*</sup> " imm. " " : " "

Nos. 3498-3500 were members of a flock feeding in the undergrowth near a brook.

Faradje (mordjan)

Nov. 28. 1911.

in the forest.

*Diachlorophrys certanea*

3501. ♂ Small Black-r-white Fly catcher, with white over eye. ♂ testes slightly enlarged.

3502. Same as preceding. ♂. testes slightly enlarged. Collected in woods near rest-house of mordjan, where they were not uncommon. Both sexes fly noisily. The call is low, short, and hoarse. Mounted for Congo Forest group.

*Catoptrophorus sp.*

3503. ♂ testes slightly enlarged. Stomach contained a white coleopterous larva & pieces of insects. At present these birds make little or no noise.

Nov. 27.

*Andropadus sp.*

3504. ♀. o.m.e. Stomach contents: small insects. A common bird in the wood near mordjan.

Nov. 29.

3505. ♂. t.e. *Gurdinus*

Stomach contents: insect remains.

3506. ♀. ovary somewhat enlarged. *Gurdinus*

Stomach contents: insect remains.

3507. ♂ t.m.e. *Bleda syndactyla vocans*

Stomach contents: remains of a grass-hopper and other insects.

3508.

*Lamprolius splendidus*

Large Resplendent Starling, ♂ testes slightly enlarged. Stomach contents: fruit. Common in the wood near mordjan, also at Aba. Not yet observed at post of Faradje.

3509.

♀. o.m.e. Stomach contents: pieces of small insects.

*Phyllastrephus scandens orientalis*

Very common and noisy in wood near Mordjan not noticed near Faradje post.

Sarey

Nov. 30

35

Congo F.Z.  
Sarey  
Mordjan

35

35

35

35

35

35

35

35

35

## Faradje (Mordjan)

Nov. 30. 1911.

Burnesia richeniour

3510. Blue-gray Warbler with white throat. ♀ o.n.e.  
Stomach contents: insects. About the post at Faradje it occurs sparingly in bushes near water. They usually form parties of 5 or 6, which utter, in place of a song, excited "cheeping" notes, two or more birds together.
- American Museum 599041 Natural History.
- Diaphorophyia castanea. ♂  
Faradje. Nov. 28. 1911. Lang-Chaplin
- Stomach contents: pieces of beetles.
3512. Cosyphax. ♂ juv. Stomach contents: remains of a mantis & other insects.
3513. ♂. testes slightly enlarged. Stom. cont.: a spider. Fairly numerous in wood near Mordjan, not observed near post of Faradje.
3514. Small yellow bellied Sunbird. ♂. testes slightly enlarged. Stomach contents: insects. A forest bird from wood near Mordjan.
3515. ♂. testes slightly enlarged. Stomach contents: insects.
3516. ♀. o.n.e. Stomach contents: beetles.
3517. Kingfisher. ♀. o.n.e. Stomach contents: a mantis. Halcyon sinensis.
3518. Red Crested Plantain-eater. ♂. t.n.e. St. cont: fruit. Musofaga rosea  
One of a pair found in wood near Mordjan.
3519. White crested Plantain-eater. ♀. ovary slightly enlarged. Turaco leucophaeus

## Faradje (Mordjan)

Nov. 30. 1911.

Burnesia richmondi

3510. Blue-gray Warbler with white throat. ♀ o.n.e.  
Stomach contents: insects. About the post at Faradje it occurs sparingly in bushes near water. They usually form parties of 5 or 6, which utter, in place of a song, excited "cheeping" notes, two or more birds calling together.
3511. ♀. o.n.e. *Bela syndactyla weisseni*. Stomach contents: pieces of beetles.
3512. *Cossyphax* <sup>verticalis</sup> ♂ juv. Stomach contents: remains of a mantis & other insects.
3513. ♂. <sup>Audubonii sp.</sup> testes slightly enlarged. Stom. cont.: a spider. Fairly numerous in wood near Mordjan, not observed near post of Faradje.
3514. Small yellow bellied Sunbird. ♂. <sup>Anthrax collaris hypoxanthus</sup> testes slightly enlarged. Stomach contents: insects. A forest bird from wood near Mordjan.
3515. <sup>Audubonii sp.</sup> ♂. testes slightly enlarged. Stomach contents: insects.
3516. ♀. o.n.e. *Stigophorus virecino*. Stomach contents: beetles.
3517. Kingfisher. ♀. o.n.e. *Halcyon senegalensis*. Stomach contents: a mantis.
3518. Red Crested Plantain-eater. ♂. t.n.e. St. cont fruit. *Muscophaga rossae*. One of a pair found in wood near Mordjan.
3519. White crested Plantain-eater. ♀. ovary slightly *Turacæ leucophaeus*.

Faradjie (Mordjan)

Nov. 30. 1911.

enlarged. Stomach contents: fruit. Common in wood near Mordjan, where the notes (*Tamini* or a new ally) of the green-headed forest *Duracus*, were also heard today. We also heard the call of the large Corytheola, and have seen the gray Schizophis; so that the five species of plantain-eaters occurring at Miangara are also to be found here, tho about the post of Faradjie only the gray and white-crested species are to be seen.

3520. Hoplopterus <sup>spinosus</sup> ♀ ad. Was walking on the ground near the houses at the post. This is the first time we have seen it.

3521. Large Black & white Hornbill. ♂. t. n.e.  
Bucalites subcylindricus.  
Irides brown, orbits dusky, bill and anterior half of casque blackish, basal half of casque whitish above, soft skin on feet blue gray, larger scales & claws blackish, soles yellowish.

3522. Same as preceding. ♀. ovary slightly enlarged.  
B. subcylindricus. From wood near Mordjan, never noticed

about post.

Dec. 2. Neophasianusruficollis

3523. Red-rumped Woodpecker. ♀. ovary slightly enlarged.

3524. Weaver (Amblyospiza) <sup>capitata</sup> ♂. t. n.e. St. cont.: seeds.

## Faradje (Mordjan)

Dec. 2. 1911.

3525. *Campylorhynchus phoeniceus*  
Red-shouldered Cuckoo-Shrike. ♂ t. n.e.  
Stomach contained insect remains.

3526. *Nylobrax luciolaimus*  
Small yellow + black Barbet. ♂. t. n.e.

3527. *Glycophagus atricollis*  
Glycophagus. ♂. testes very small. Iris red.

3528. *Barbet (Tricholaema)* ♀. ovary slightly enlarged

3529. *Hoopoe*. ♀. ovary slightly enlarged. *Upupa*  
~~The hoopoe has a stiff tuft in legs & wings.~~ *somalensis*

3530. Blackish Sunbird, with throat & upper breast  
*Chalcomitra angolensis*  
metallic green. ♂. t. e. Collected in wood  
near Mordjan. A forest species, not yet  
seen near post.

3531. Large Black & white Hornbill. ♂. testes  
*Buceros subcylindricus*  
slightly enlarged. Stomach contents: fruit.

3532. Mate of no. 3531. ♀. ovary slightly enlarged.  
Stomach contents: fruit.

During a week's stay near Mordjan, we saw about 8 of these hornbills. They are forest birds, going usually in pairs, and perching in the high trees. The flight is noisy, and <sup>both sexes</sup> they utter a loud mournful note, slowly repeated (resembling that of *Ceratogymna*), or a hoarser, shorter, reiterated call.

Paradje (Mordjan)

Dec. 3. 1911.

*Dicrurus adsimilis divaricatus*

3533. Drongo. ♀ o.n.e. <sup>♀ ride red.</sup> *Dicrurus adsimilis divaricatus*

3534. " ♀. o.n.e. *D. a. divaricatus*

*Mesopisus griseus porcephalus*

3535. Red-rumped Woodpecker. ♀ ovary slightly enlarged. Stomach contained a white grub & insect remains.

3536. Red-crested Plantain-eater. ♂. t. n.e.

*Muscophaga rossae*

Stomach contents: fruit.

3537. Same as preceding. ♀. o.n.e. Stomach contents: fruit.

They still call, tho it is perhaps during June and July that they are most noisy.

3538. *Macrodipteryx macrodipennis*. ♀. o.n.e. *Stomach*  
<sup>= longipennis</sup> contents: 3 green grasshoppers & a moth.

This evening, from sunset till 10 o'clock we walked from Mordjan to ~~Paradje~~, coming back to Paradje. It was on an adjacent stretch of this same road that I counted 28 examples of *Macrodipteryx vexillarius* on July 11th last. Tonight the moon was full, and again I could count the night jars, which have of course entirely changed with the season. About 20 were seen in all, of which a dozen were the long-tailed *Scolopax*.

Faradje (mordjan)

Dec. 3. 1911.

3 adult males of *M. macropterus*, while the remainder were difficult to identify, tho probably females and immature males of *M. macropterus*. The season of activity of this latter species is just beginning, and the dry season has just set in in earnest. After the first two hours of darkness most of the nightjars seem to retire from the road, probably they have eaten their fill, but toward morning they are again to be seen there.

Nov. 30.

3539. Wood Owl (*Syrnium*) ♀. ovary much enlarged.

*Stix nicholisi*  
Stomach contents: remains of beetles & orthopterous insects. Shot in wood near Mordjan, as it was being mobbed by drongos and birds like No. 3509. (a young owl of this species, just well enough post by a native on Feb 14. 1912.)

Dec 4.

3540. Red- & black Weaver (*Pyromelaena*) ♂ t. n.e.  
(in post-nuptial molt.)

Now is the season when many of the weavers assume dull-hued plumages. The black widow-birds are fast losing their long

Parade

Dec. 4. 1911.

Aba  
Dec 10

rectrices and somber body plumage, while many of the related species have practically completed the molt. (But males of Hypochera are still completely black, and widows like No. 3408 must molt very late, for one was seen in full breeding plumage at Aba on Dec 13 th.)

Dec 7.

1545

3541. Ring Plover. ♀. o. n. e. Common here at present.  
*Charadrius dubius curonicus.* *Clyties tridactylus*

3544

3542. Black Barbet with red face. ♀. ovary slightly enlarged. Stomach contents: fruit.

3545

3543. Long-tailed Boatucker. ♂. testes somewhat enlarged. *Scotornis climacurus*  
(photo) Shot during the morning in a papyrus swamp, where two of them were spending the day. This is an unusual situation, but in places the papyrus had been cut and broken down by workmen in search of material for house building, the papyrus stalks, properly beaten and twisted, serving as cords, while they are also split to make baskets.

3546

3547

3548

3549

Aba  
(Between Mordjan & Dabiri)  
Dec. 10. 1911.

3550

3544. Small Wood Hoopoe. ♂ testes slightly enlarged.  
*Scopetus sterimus emini*  
Stomach contents: remains of insects + 2 small

3551

Aba (Mordjan to Djaliri)

Dec 10. 1911.

caterpillars.

3545. Hoopoe. ♂. testes slightly enlarged. Stomach  
*Upupa senegalensis*  
contained many ant-lions. Flew up from  
the road before us. Now that the rains have  
ceased, there is much dry sand on the road,  
and ant-lion pits are very abundant.
3546. Creeper. ♀. ovary slightly enlarged; but skull not  
*Selasphorus sasin*  
completely ossified. Stomach contents: small  
insects.

3547. Small yellow-breasted Finch. ♀ imm.  
Aba (bill. blow) Dec. 11.

3548. Kestrel. ♂. t. n. e. Stomach contents: a lizard  
*Cercus t. timoninus*  
and large grasshoppers. (Kestrels are very common  
at present both at Faradjie & Aba.)

- Dec. 12. 3549. Thrush (*Monticola*) ♂. t. n. e. Stom. cont.: insect remains.

3550. " " ♂. t. n. e. Iris dark brown,  
bill mostly blackish, but dull yellowish gray  
beneath base, feet dark brown. Stomach contents:  
fruit & remains of small insects.

In July we noticed none of these thrushes  
at Aba, but during the present visit counted  
about 8 in all. They perch quietly in the  
small trees on the hills, and are so tame  
as to be approached with ease. I heard no  
notes uttered.

Aba (Vill. 9 bu.)

Dec. 12. 1911.

3551. Swallow (*Riparia rufigula*). ♂. testes slightly enlarged

3552. " ( " " ). ♀ ovary "

Common about the hills at Aba, occasionally  
alighting on steeply sloping rocks. Silent.

3553. Swallow (*Hirundo puella*). ♀. o.w.e. Seen

frequently about the hills, often very high  
in the air. Seen not yet to be nesting.

3554. Crested Cuckoo. ♂ t.w.e. Stomach contents: hairy  
caterpillars. Not uncommon at Faradje and

Aba; its call, frequently heard during the rainy season,  
commences with a slowly uttered, hollow sounding "keer, keew,  
keer, keer ..." After this note has been repeated about a dozen  
times, it hesitates a moment, and changes abruptly to a  
rapid series of short, reiterated notes, "keu-keu-keu-keu ...."  
to the number of 20 or more, dying out gradually toward the close.

3555. Small Crown Bunting. ♂. t.e. Irides dark brown,  
*Trivellaria septentrionalis*  
maxilla mostly dusky brown, mandible yellow,  
feet buff. Crop contained grass seeds.

3556. Same as preceding. ♀. o.e. Irides dark brown.  
maxilla dusky brown, mandible & feet buff.

Dec. 13.

3557. Hoopoe. ♂. testes slightly enlarged. Stomach  
*Upupa somalensis*  
contents: several ant-lions and a small  
caterpillar.

3558. Hoopoe. ♀. o.w.e.  
*Upupa somalensis*

Aba (Vill. 9bu.)

Dec. 13. 1911.

slightly enchy.

" "

occasionally

Silent.

Seen

very high

nesting.

contents: hairy

Paradise and

rainy season,

singing "heer, hee

ed about a dozen

songs to a

few hours....

Toured the coast

dark brown.

dull yellow.

etc.

dark brown.

feet buff

Stomach

a small

3559. *Macrodipteryx macrodipterus*, ♂. t. n.e.  
= *longipennis*  
in the daytime

Often flushed, about the bases of the hills,  
frequently in places where the grass had been  
freshly burned. The more exposed rocky  
parts of the hills are occupied by the gray  
goatsuckers like no. 3589.

3560. Small Brown Bunting. ♂. t. e. Crop contained  
*Trinillusaria septentrionalis*  
grass seeds.

3561. Small Brown Bunting. ♂. t. e. Crop contained  
*Trinillusaria septentrionalis*  
grass seeds.

Not noticed in July, these buntings  
on the hills are abundant at present, hopping on the  
rocks, or perching in the small trees and  
brushes. The song is a simple repetition  
of a few short notes that can hardly be  
called musical.

3562. *Ochthoecetes badius* (Vicell.)  
Black-masked Weaver. ♂. t. n.e. Stomach  
contents: green seeds & a few insect remains.  
Irides dull yellow.

3563. *Dicrurus brunnineus* (Linné)  
Bee-eater. ♂. t. n.e. Shot in plantation at base of hill,  
2 more seen in same place on following day.

3564. Partridge. ♂. testis slightly enlarged. Iris  
*Ptilopechus petrosus* (Gmelin)  
medium brown, feet, orbits, and base of bill  
carmine, tip of bill brown. One of a pair

Aba (Vill. & bw)

Dec 14. 1911.

found on the top of a rocky hill where there was much high grass between the boulders, and many stunted trees. Another pair was seen on a neighboring hill. I have heard no call that I could attribute to these birds.

3565. <sup>Thamnolægus sp.</sup> Rock Thrush. ♀ o.n.e. With her mate she was sitting under a ledge at the top of a perpendicular cliff, some 50 feet high. The male was singing lowly. Sometimes the song of this thrush reminds me of the white-crowned Cossypha in the wooded swamps, at other times it seems to resemble that of the black thrushes with white wing patches that inhabit cultivated ground and other open places. I believe they are less musical and active than they were last July.

3566. Thrush (*Monticola*) ♂. t.n.e.

3567. " " ( ) ♀. o.n.e. Stomach contained many ants. As compared with thrushes like no. 3565, these birds are extremely tame.

Aba (Vill. Ibu.)

Dec 14. 1911.

*Sambagae* Alex.  
*Alouanahendeliola* Lombok

3568. Gray Fly catcher. ♂. testes slightly enlarged. Irides dark brown, maxilla dusky brown, mandible buff, feet dusky brown. Seen frequently on the hills during this visit to Aba.

3569. Quail (*Orecaudafactoria*) ♂. im. Caught in a pasture near Ibu's village.

3570. Small Crested Lark. ♂. testes slightly enlarged.

50% CZ in  
ex change.  
Gift 6/1928

*Calandrellus modestus*  
Stomach contents: small seeds. Fairly common about the hills. At this season its haunts are invaded by European pipits, buff-bellied pipits and yellow-breasted Wagtails.

3571. "Flycatcher-Shrike". ♀. o. n. e. Skull not quite completely ossified.

3572. Swallow (*Chelidon*). ♀. o. n. e. Two of these swallows were seen flying about a hill in company with many <sup>of the</sup> buff-throated species like no. 3573, and alighting on some sloping rocks warmed by the afternoon sun. (<sup>Martins were rather numerous at</sup> Banguna's in Apr. 1912. <sup>Seemed to be</sup> rather fond of flying about the hills.)

3573. Swallow (*Riparia rufigula*). ♂. im.

3574. Small Brown Bunting. ♀. ovary somewhat enlarged. *Prinellaria septentrionalis*

3575. Brown Warbler. ♀. o. n. e. *Cisticola petrophila*.

Aba (Vill. 3bu.)

Dec 14. 1911.

3576 Brown Warbler ♂ imm. Characteristic birds  
cinnicola petrophila of the hills, running actively over the rocks  
with tails cocked up like wrens.

3577 ♂. t.e.  
s. 2957.

3578. Same as 3577. ♀. t.w.e. Stomach contained  
remains of a grasshopper and other insects.  
Several were seen about the hills, perching  
in small trees, and wagging their tails.  
European redstarts are fairly numerous  
at this season, their tails quiver from  
time to time. Whinchat, so common  
here in the dry season, are also to be seen  
commonly on the hills.

3579. Kestrel, ♂ [Lim.] t. n. e. Stom. cont.: grasshoppers.  
Dec. 15. Circus t. terminalis.

3580. Lesser Kestrel, ♂ [Lim.] t. n. e. Irides dark  
Circus neumanni.  
brown; orbits, cere, and base of bill yellow;  
bill blue gray; feet yellow, claws light yellow.  
Stomach contained a grasshopper.

3581. Streaked Rufous Falcon. ♂. Testes slightly  
Circus albex albex enlarged. Iris pale buff; orbits and cere  
light greenish gray; bill bluish gray at base,  
black at tip; feet yellow; claws black.  
Stomach contents: grasshoppers.

Aba (Vill. Ibu.)

Dec. 16. 1911.

3582. Streaked Rufous Falcon. ♂. t. n. e.  
*Cerchnis a. slopex.*
3583. " " " ♂. t. n. e. Stom. cont. a grasshopper.
3584. " " " ♀. o. n. e. Stom. cont. a grasshopper.
3585. " " " ♀. ovaries slightly enlarged

These falcons seemed fond of the hills, but were shy except when attracted by our setting fire to the grass. Then in company with kites & European kestrels they, <sup>baldly</sup>, came to catch the grasshoppers put up by the flames. To eat their prey they would alight in a tree, tho the kites ate during flight. Mr Lang saw one alight on the bare rock. The flight and call are kestrel-like, tho the latter is hoarse. Today I saw about ten. [See note under No. 4529.]

3586. Kestrel. ♀. ovaries not enlarged.  
*Cerchnis t. tannimulus* *Hemundo puebla*
3587. Streak-breasted Swallow. ♀. o. n. e.
3588. Wood Hoopoe. ♀. o. n. e. *Phoeniculus purpureus* *melanotos*
3589. Large Gray Goatsucker. ♂. t. n. e. At this season too they may be heard calling at dusk and a little before dawn.  
*Caprimulgus trinotatus triotyma*

Dec. 17. 1911.

3590. Hawk (*Accipiter*) ♀. ovary slightly enlarged.  
*astur sparrow formatus*  
Irides bright yellow, orbits yellow, cere green.

Aba

Dec 17. 1911.

till black with bluish base; feet yellow, claws black.

Dec 18.

3591. Yellow Wagtail with black crown. ♂. t. n.e.

*Bidytes flavus melanoccephalus*

Shot in a small muddy swamp near the foot of Aba.

3592. Large Bustard. ♀. o.n.e. Stomach contents:  
*Otis derkheimi*. many large black ants & beetles, a few bugs

& caterpillars, and some red berries. A bird of the dry season, arriving in the latter part of November, and feeding while unseed grass, or in other open spots. Probably breeds here

Dec. 19. (near Frontier N.E. of Aba) "Libuque".

3593. Wheatear. ♂. testes very small. Several seen  
*Oenanthe oenanthe*. on hills today.

3594. Hawk (*Accipiter*). ♂. testes very small. Stomach  
*Accipiter sphenurus*. contents: remains of a lizard and a small bird. Irides yellow, rim of eyelids, cere, & corners of mouth yellow, till blackish, feet yellow, claws black.

3595. White-headed Barbet. ♂. t. n.e.

*Lybius leucocephalus*

3596. " " " ♀. ovary slightly enlarged  
Stomach contents: fruit. Members of a party of five that frequented some large trees at the foot of a hill. These barbets seldom go singly. The flight is well sustained and rapid, with swiftly beating

Aba (<sup>near frontier</sup> NE of post.)

Dec 19. 1911.

wings. The characteristic notes are somewhat metallic, but not short; being uttered in a series, starting strongly, but dying gradually out. (See note under 4101.)

3597. white-necked Stork. ♂ testes not enlarged.

Dissoraa microscelis

Stomach contents: a soft mass of vegetable (?) matter, enclosing remains of a large grasshopper and a few other insects & insect-larvae.

With one other of its kind, it alights nearly on top of a high rocky hill, where a little water oozed out over the face of a large sloping rock. Not very common; we have no record of their occurrence in the rainy season.

Dec. 20.

3598. Hoopoe. ♀. ovary slightly enlarged.

Upupa somalensis

Usually we have found hoopoes very shy, but on a freshly burned hillside there were three that showed but little fear of man. The males utter a soft, low, "hoo-poo-poo" and a hoarse "haaving" noise. The flight is often undulating like a woodpecker's, with depressed crest,

Aba (new Antice)  
NE part.  
Dec. 20. 1911.

while the latter may be erected while  
perching, or not.

*Lutreola leucophaea*  
3599. White-headed Barbet ♂ t. n. e. Stomach  
contents: fruit.

3600. Small yellow-bellied Sunbird with elongated rectrix.  
*Hedydipna platura*  
♂ t. e. Stomach contained a very small spider  
Iris dark brown, bill & feet black. First  
observed on Dec. 10. Does not seem to visit  
flowers, <sup>much</sup> but prefers the bushes and trees in  
freshly burned spots. 5 or 6 seen today over  
hillside.

3601. *Macropygia macropygia*. ♂ [Im.] t. n. e.

3602. Small Brown Bunting ♂ [im. <sup>longipennis</sup> (in juvenal pl.)]

3603. " " " " ♂ t. e.  
(photo)

The nest of no. 3603 was discovered by Mr. Lang  
on a flat rock of a barren stony hill. It had  
tho a few prostrate grasses lay around it,  
absolutely no shelter from above, and was composed  
of pieces of dry moss and other plant refuse, lined with  
fine dry grass-tips. The two eggs were pale greenish,  
reddish  
thickly covered with small brown spots, most numerous  
about the larger end. The ~~incubating~~ bird sat closely,  
but afterwards the parents approached only with  
great caution.

Aba (near frontier)

Dec 20. 1911.

*Hypargos dybowskii*

3604. <sup>A</sup> Small Gray Weaver with white-spotted breast & red back & rump. ♀. t. n. e.; skull not quite completely ossified. Crop contained grass seeds.  
Only a few seen.

3605. Brown Warbler. ♂. t. n. e. Stomach contents: *Cisticola petrophila* a grasshopper & small insects.

3606. Brown Warbler. ♂. t. n. e. Skull not quite completely ossified. *Cisticola semitorquatus* (?) Stomach contents: small insects.

Dec. 21. (near Port of Aba)

3607. Rufous-winged Hawk. ♂. t. n. e. Stomach contents: *Buteo rufipennis*.  
a few insect remains. A bird of the dry season, arriving about Nov. 1.  
Is attracted by grass fires.

*Hyliota flavigaster*

3608. Warbler. ♂. t. n. e. Stomach contents: some insect eggs & remains of insects.

3609. ♂ im. <sup>Hyliota flavigaster</sup> 5.2969. testes very small; skull almost completely ossified. Stomach contents: insect remains.

3610. ♀ im. <sup>Hyliota flavigaster</sup> 5.2969. ovary small; skull nearly completely ossified. Members of a party of about 5, presumably a single family. Birds of the small trees, whose voice I have yet to hear. They have been noticed twice near Faradje.

Aba

Oct 21-1911.



Dec 22. (Aba post to Djibiri) sometimes attr.



## Enneapterygius gubernator

3613. Shrike. ♀. ovary slightly enlarged. Stomach contents: remains of beetles.

- 3614 Finch ♂ testes slightly enlarged (singing).  
S2946 *Phainopepla elegans*.  
Crop contained seeds. Only example seen  
since last August.

3615. Shrike. ♀. o.n.e. Stomach contents: insect remains

3616. Yellow-crowned Oriole. ♂. t. e. Stomach  
*Oriolus auratus*  
contents: fruit of Irides red. Was singing,  
in brush on dry ground.

p. 119

BLANK

No. 3617 (cont'd). On Jan. 26, 1912, near our Rhine group  
Camp, about 16 miles N.W. of Faradje, I found a  
nest of this red-tailed Buzzard. It was placed  
at a height of 25 ft or so, <sup>between 4 or 5</sup> ~~in one of these trees~~  
common in this region, that bear large  
pendent, woody fruit, resembling in form  
a "French loaf" of bread <sup>(Kigelia)</sup>. It was rather bulky,  
and only slightly concave above, composed  
entirely of dry sticks, save for a lining of  
green leaves. The two eggs (which have been  
preserved) were greenish white, unspotted,  
and as they were slightly incubated I presume  
the set was complete. The old bird sat very  
close, and after being frightened off, came  
back and flew around uttering a call  
almost like that of *Buteo lineatus*.

On the same tree, just a little above the buzzards' nest, there were 12 or 15 nests of <sup>(S. itaga Lengini)</sup> yellow weavers. See nos 3649 & 3664.

Aba (Djabiri)

Dec. 22. 1911.

3617. Hawk (Buteo) ♀. o. e. Stomach contents : a striped rat. (See opposite page)

Faradje

Dec. 23.

3618. Bee-eater (M. nubicus) ♂. t. m. e. After Sept. 9, when I collected a single example, none were observed till about Oct. 25. At present they are common.

3619. Small yellow-bellied Sunbird with elongated <sup>tail</sup> ~~plumosa~~ rectrices. ♂. t. e.

3620. Same as preceding (middle rectrices just coming in). ♂. t. e.

Dec. 25.

3621. Large black & white hornbill. ♂. testes somewhat enlarged. Stomach contents : fruit.

3622. Same as preceding. ♀. ovary somewhat enlarged. Stomach contained fruit.

From wood near Mordjan.

3623. Owl (Scops) ~~leucopsis~~ ♂. testes slightly enlarged.

Dec. 28.

3624. Anastomus. ♂. testes very small. Iris dark brown most of ~~lambelligerus~~ but tip of mandible gray. Bill and toes black, naked skin of throat pale purple, feet black.

Paradise

Dec. 28. 1911.

*Chalcomitra senegalensis* ad.  
auct.

3625. Sunbird ♀. Ovary slightly enlarged. Stomach contents: a little ball of rubber and remains of a few tiny insects.

Dec. 30.

3626. Owl (*Scops*) <sup>♂</sup> <sub>*caffer*</sub>. Testes slightly enlarged. For nearly three months we have been able to hear them calling almost every night, from the post. Their vocabulary is not limited to the single "metallic" note, for at close range one may sometimes hear a fainter - but shriller ~~note~~ cry, and other short, low sounds. They haunt trees, both along the river-bank and off amid the high grass, and occasionally visit the groves of rubber-trees.

Dec. 31.

3627. Owl (*Scops*) <sup>♂</sup> t. n. c. Irides yellow, rim of eyelids greenish brown, bill dark green with blackish tip, feet light greenish gray with <sup>Tars</sup> claws dusky greenish. Stomach contained remains of an insect & a spider. Shot in same spot as No. 3626, among rubber trees near post. Uttered only a hoarse "haaa - haaa" -
3628. *Macrodipteryx macrodipterus*. <sup>*longipennis*</sup> ♂. Testes slightly enlarged. Stomach contents: 2 moths, 2 grasshoppers.

Paradjie

Dec. 31. 1911.

12 small beetles, 3 leaf hoppers, and 18 small bugs.

In the throat were a mosquito and an ant, the latter having fastened its mandibles firmly in the flesh. Common at present.

Aba

3629. Small black hornbill. *B. t. c.* Secured from natives, *Lophoceros fasciatus*, at Aba, where they are rather common.

Paradjie

Jan. 1. 1912.

3630. Stone Curlew. ♂. testes somewhat enlarged.

*C. semipalmarum*  
Stomach contained a beetle, a large earwig, and remains of other insects. Shot in the post at night. Three or four come every evening now, and are very noisy. They seem to spend the day somewhere up the river.

Jan. 2.

3631. Brown Eagle (*Aquila*) ♂. testis very small.

*Aquila rapax albicans*.

Stomach contained remains of a giant rat (*Cricetomys*). Killed with a stick.

Jan. 4.

3632. Marabout. ♀. ovary somewhat enlarged; oviduct enlarged, as tho the eggs had just been laid.

Plato. Bill greenish white, very dirty & irides grayish brown on inner edge, shading nearly to white on outer rim. Crown, front of lores and a

Faradje


Jan. 24. 1912.

patch at each side of the base of mandible covered with a rough, blackish-red, scab-like skin. Skin below eye orange-yellow, shading to red on back of head. Back of neck light blue, separated abruptly from the dirty pinkish white on the front of neck and the pouch. Base of neck above is bright brick red. On the head & neck are many spots of black, usually small, but forming a V-shaped mark about ear. Just below the feathers on the tibia is some soft naked skin, bright red. The scaly parts of the feet are black, but whitened with excrement.

Bangura says  
marabouts  
are seen there  
out the year,  
but nest only  
in the dry  
season.

Throughout the last rainy season neither Mr. Lang nor I saw a single marabout, but during the present month (Jan.'12), about 16 miles NW of Faradje, we found them rather numerous. Meat was purposely thrown out at a little distance from our camp, and thus we were attended most of the time by numbers of small vultures (*Neophron*) and sometimes marabouts and other vultures (*Croto* *gyps* & *Pseudogyps*). Marabouts were also attracted by grassfires. The largest number of these carrion-eating storks noticed at once was ten, flying

Faradje  
Jan. 14. 1912.



asions 5 were seen to alight on  
ajority were always birds of last  
about 1 in 6 being old birds. They  
pected, and with caution  
this shot-gun range. In  
by about so high as almost  
aving spied carrion on the  
circle around, descending  
far above the earth (perhaps 200 ft)  
an angle of 45°, the beak point  
spiral descent becomes more  
ulture already at work, it

departs abruptly. Having eaten its fill, the marabout  
mounts a tree, where it sits at peace with the  
vultures. Whether perching or walking, the attitude is  
usually erect, the head either drawn down when at rest, or  
reared when on the alert. the size of the neck pouch varying  
greatly.

Jan. 7. 3633. Vulture (Pseudogyps) ♂. [Imm] testis very small  
africanus. Iris dark brown, bill black, skin of face & throat  
dusky, that of neck very light green, feet  
blackish.

Jan. 8. 3634. Vulture (Pseudogyps). ♀ ovaries very small [Imm]  
africanus. Stomach contained carrion and pieces of dry  
grass.

Faradje  
Jan. 04. 1912.

over our camp. On two occasions 5 were seen to alight on a single tree, but the majority were always birds of last year (see no. 3645), only about 1 in 6 being old birds. They were not as wary as I expected, and with caution could be approached within shot-gun range. In general they seem to fly about so high as almost to escape attention. Having spied carrion on the ground, they begin to circle around, descending slowly, but when still far above the earth (perhaps 200 ft) their legs drop down to an angle of 45°, the beak points groundward, and the spiral descent becomes more rapid. If there is a vulture already at work, it departs abruptly. Having eaten its fill, the marabout mounts a tree, where it sits at peace with the vultures. Whether perching or walking, the attitude is usually erect, the head either drawn down when at rest, or raised when on the alert. The size of the neck pouch varying greatly.

Jan. 7.  
3633. Vulture (Pseudogyps) ♂. [Imm.] testes very small <sup>africinus</sup>. Iris dark brown, bill black, skin of face & throat dusky, that of neck very light green, feet blackish.

Jan. 8.  
3634. Vulture (Pseudogyps). ♀ ovaries very small [Imm] <sup>africinus</sup>. Stomach contained carrion and pieces of dry grass.

FaradjeJan. 8, 1912.

3635. Vulture (Lophogypa) ♂ [Imm.] t.n.e. Irides dark brown, top of cere light green, rest of cere and base of bill light blue, outer part of bill dull light red, bare skin of head & neck pinkish & bluish white, feet pinkish white with dusky claws.

Jan 14

3636. Large gray swift, with white throat & belly. (Cypselus) ♂. t.n.e. Irides very dark brown, bill black, feet dusky with slight purplish tinge. Stomach contents: many small grasshoppers and bugs, a few small beetles & a leaf hopper.

3637. Same as preceding. ♀. o.m.e. Stomach contents: many grasshoppers (mostly small), and a few bugs.

This morning, for the first time, I saw a few of these swifts flying high overhead, and later a flock of 50 or 60 whirling about in circles, but in the afternoon a grass-fire started by natives to scare game into the nets attracted them by hundreds. Their habits are like those

Paradiji

Jan. 14 1912.

of the other large swift already collected here (nos. 2553-2554.); they keep usually pretty high, but come flying along over the brush at terrific speed, or flocking where the fire burns hottest. Tho we stayed 2 weeks longer on the same spot, and saw several more fires, we did not observe these swifts afterward.

Jan 16.

3638. Bateleur Eagle. ♂. t. n.e. Irides dark brown,  
photo *Gierathipus esculatus*.  
lores, cere, and skin at base of mandible  
bright red, beak blackish at tip, yellow  
basally, feet red, claws black. (These are  
the colors during life; but the red of the  
cere, lores, and feet depends on the blood.  
In the present case, the cere, which is  
not hard as in other birds of prey, and  
the feet, which are covered with prominent  
rounded waxy scales, both turned orange  
soon after death, tho the cere regained  
its brilliant color when the bird was  
hung up by the feet.) Stomach contents:

Paradise

Jan. 16. 1910.

3 shrews and a piece of bone.

Tho the Bateleur Eagle is a common bird of this region, I never, till this morning saw one perch, for they seem to spend most of their life on the wing, scouring the country for food. The present example uttered a very loud "caw", which is sometimes repeated at intervals. These eagles have often sailed for some time over the spot where we placed meat to attract ~~vultures~~ vultures, and once I saw 2 immatures specimens fly up from the meat.

(On Jan. 25-12 we observed an adult Bateleur Eagle with white back & tail coverts, but I believe the red-backed kind are commoner.)

*Indicator indicator*

Jan. 20. 3639. Honey Guide, ♀. o. ex. Stomach contents: bee-wax. Irides grayish brown, orbits dark gray. Bill dirty brownish, with a tinge of pink. Feet dark gray slightly bluish in places. Was calling much like a male.

(The common note of the male may best

Paradise

Jan. 20. 1912.

be written "cutta-cutta-cutta-cutta . . . .",  
and is uttered very rapidly.

3640 Weaver. ♂ t.m.e. Tries dark brown,  
Toupleutes cossistriatus. maxilla brownish, mandible grayish white,  
feet light brown. Stomach contents: seeds & a  
few insect remains.

Jan. 22.

3641. Kite (Milvus) ♂ t.m.e.

3642 " M. a. percussus ♂ [imm] t.m.e.

At present, in the dry season, kites are  
abundant everywhere, flocking to  
a grass-fire to catch grass-hoppers on  
the wing, beating about over villages on  
the lookout for an odd scrap of meat.

Jan 24.

3643. Vulture (Lophogyps) bacpiculus. ♀ o.m.e.

Jan. 25.

3644 " Lophogyps " ♀. o.m.e. Stomach

contents: porcupine quills (longest one about  
4 in.) grass, and 2 pieces of bone.

This vulture was occasionally seen at  
the Rhino-group Camp, but never more than  
3 at a time. Adults, with white crown, crop

and belly were seen, but could not be secured. The habits seemed much like those of *Pseudogyps*. (As to the color of the head in Neophron — We often noticed that adults of Neophron, which can be distinguished by the whitish nape, had the skin of the head deep pink or light red, whereas the young were often bluish <sup>slate</sup> and very changeable there. That this color is merely due to the blood, I found when I broke the wing of a red-headed bird with the rifle. On catching the wounded bird, I was surprised to find the bare skin of its head had already faded to bluish white.)

Jan 24.

*Lycoptilus caeruleifrons*.

3645. Marabout <sup>I inv.</sup> I inv. ovary slightly enlarged. " Stomach contents: hair, a piece of hoof & a few insects remaining.

Jan. 25. *Bucanetes githagineus* <sup>Hartert</sup> 3646. Small gray Weaver with red rump, black cheeks

& throat, and small white spots on the sides.

J. t.m.s. Irides brown, rim of eyelids blue gray

bill blue ~~gray~~ at base, black at tip, feet dusky.

Flew up off the ground in a burned spot in the brush. The first specimen I have seen.

Leptoptilus crumeniferus

On April 15, 1912, I went with Sangura to visit a place where marabouts are accustomed to breed. He assured me that there were sometimes as many as 20 nests, and so I set out with good hopes. At 4 hours march to the westward of his village we came to a narrow, wooded valley, with a couple of very tall trees towering up above their fellows. One was very straight, with a smooth, branchless trunk, and among its leafy boughs, at a height of 55 or 60 feet we could discern two large flattish nests, rather roughly built of dry sticks, and supported by several more or less horizontal branches, well out from the main boughs. On top of one nest sat a single young marabout, <sup>and not easily seen from below</sup> of rather large size, the other was empty, but on the small twigs about it there hung two weaver-birds' nests (undoubtedly the same species, as no. 3649-51). Sangura was rather discouraged by the small number of nests, but said the main colony was over in the next swamp, so up we went over the intervening rise, covered with a growth

of coarse grass. The next depression, a quarter of a mile beyond, contained many small trees, and a few large ones, one of them a giant, with a huge trunk. Upon its branches there were 3 marabout nests, but all empty, and we quickly found that natives had already visited the spot and collected all the young, perhaps destroying many other nests of the marabouts. In this tree there were 3 weavers' nests, also placed close to a marabout's. The natives, who had camped in the hollow below, made the ascent by way of a smaller tree, passing over to the limbs of the larger one by means of vines they lashed to them. Young marabouts are considered choice eating; last year when marasha gave us two of them he said we would find them much better than ducks.

On our way back we passed by the nest containing the young bird, and

Jan 2  
364:  
Jan. 2  
364

saw one of its parents return to the nest alighting on the neighboring boughs, it approached its offspring with slightly opened wings, giving the low, reiterated grunting note we have already mentioned. This neighborhood is said to have a colony of breeding marabouts every year in the dry season, and is evidently selected by the birds because of its remoteness, there being no village closer than 4 hours, and the consequent abundance of big game, elephants, rhinoceroses and buffaloes.

In spite of Sangura's opinion that marabouts are resident here, I believe they are absent during most of the rainy season. Our hunters who were near Ayanga's in late August could not say that they had seen any, and after securing an elephant near the same place in October, <sup>they</sup> reported positively that no marabouts appeared, tho' vultures were numerous. (On Feb 18, 1913, Kasime sent 2 young Marabous to the post.)

(At Garamba, in March 1912, Mr Lang observed weaver birds' nests about a nest of Pseudogyps.)

p 554

po134u.tif

Blank  
Page

Total 485 npo

p. 135

BLANK

(end of Orinth 2)

